

70-256

PRECAUTIONARY STATEMENTS

CAUTION

Keep Out Of Reach Of Children

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Avoid breathing dust. May be absorbed through skin. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

CONTAINER DISPOSAL

Do not reuse container. Wrap container and put in trash collection.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SOIL APPLICATION: To control cutworms and wireworms prior to planting of the following vegetables:

Beans ✓	Cauliflower ✓	Lettuce	Peas ✓	Squash
Broccoli ✓	Corn	Melons	Peppers	Tomatoes
Cabbage ✓	Cucumbers	Onions ✓	Potatoes	
Carrots	Endive ✓	Parsnips	Spinach	

Apply evenly as a broadcast treatment over the surface of the soil. Immediately after spreading, mix thoroughly with the soil to a depth of 6-8 inches. Apply 1/2 lb. per 225 sq.ft.

FOLIAGE APPLICATION: Apply dust at first signs of insects. Repeat as necessary. For best results, dust in early morning or late evening when temperature is lower and air is still. Apply dust so that all parts of plants are contacted, especially upper and lower leaf surfaces. Number in parentheses, (), indicates days required between last application and harvest.

Plants	Insects
Beans (7)	Aphids, Mexican Bean Beetles, Leafhoppers
Peas (0)	Aphids
Broccoli (5), Cabbage (7), Cauliflower (5)	Aphids, Diamondback Moths, Imported Cabbageworm
Cucumbers (7), Melons: Canta- loupe, Muskmelon, Watermelon (3), Squash: Summer (7), Winter (3)	Aphids, Leafhoppers, Serpentine Leafminers, Mites, Thrips
Peppers (5)	Aphids, Serpentine Leafminers
Tomatoes (3)	Aphids, Dipterous Leafminers, Banded Cucumber Beetles, Fall Armyworms, Southern Armyworms
Sweet Corn (2)	Corn Earworm*
Strawberries (5)	Aphids, Twospotted Mites
Roses, Flowers, Shrubs, Ornamentals	Aphids, Mites, Leafhoppers, Mealybugs, Whiteflies, Scale Crawlers, Lacebugs

Note: Do not use on Ferns, Poinsettias, Hibiscus and Gardenias since injury may occur.

*Apply to silks as soon as they appear. Make application at 3 to 4 day intervals as infestation warrants. Reapply immediately after rain.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

- e. A Statement of Practical Treatment must be added to the labeling. We suggest the following statement, however, if your toxicological data suggests otherwise, please submit a draft Statement of Practical Treatment for prior approval.

"STATEMENT OF PRACTICAL TREATMENT

- If swallowed - Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.
- If inhaled - Remove victim to fresh air. Apply artificial respiration if indicated.
- If on skin - Remove contaminated clothing and wash affected areas with soap and water.
- If in eyes - Flush eyes with plenty of water. Call a physician if irritation persists."

- f. We suggest a statement be added on how to spread the dust for soil and foliage application. Is this a shaker can or are you suggesting use of dusting equipment?
- g. We note the zero days required between last application and harvest for peas. We are aware of some new varieties of peas where the pod is consumed along with the pea. We would like to suggest you add a 7 day preharvest interval for this type of pea.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosures