

U.S. ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (TS-767)
WASHINGTON, DC 20460

EPA REGISTRATION NO.
70-276

DATE OF ISSUANCE
SEP 30 1985

TERM OF ISSUANCE

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION
(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

NAME OF PESTICIDE PRODUCT

Rigo Aqueous Garden Spray

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Rigo Company
P.O. Box 39
Buchner, KY 40010

150 / 159,566
18 / 1

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others. This product is conditionally registered in accordance with FIFRA sec.

3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Add the phrase "EPA Registration No. 70-276" to your label before you release the product for shipment.
3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.


A stamped copy of the label is enclosed for your records.

Timothy A. Gardner
Product Manager (17)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL



DATE

9-30-85

R I G O

AQUEOUS GARDEN SPRAY

FOR INSECT CONTROL ON VEGETABLES AND ORNAMENTAL PLANTS

KILLS GYPSY MOTH CATERPILLARS AND JAPANESE BEETLES

MAY BE APPLIED ON DAY OF HARVEST

ACTIVE INGREDIENTS:

Pyrethrins.....	0.02%
*Piperonyl Butoxide, technical.....	0.20%
Petroleum Distillate	0.08%
INERT INGREDIENTS:.....	99.70%
TOTAL	100.00%

*Equivalent to min. 0.16% (butylcarbityl) (6-propylpiperonyl) ether and 0.04% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional precautionary statements

Accessed
with COMMENTS
in EPA Letter Dated:

SEP 30 1985

(EPA File Symbol 70-ETA)
EPA Reg. No. 70-
EPA Est. No. 70-KY-1

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.


70-276

Net Contents
Ounces

Manufactured by

RIGO COMPANY

BUCKNER, KENTUCKY 40010



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

CAUTION: Harmful if swallowed or absorbed through skin. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists. Remove pets, birds and cover fish aquaria before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams, ponds, tidal marshes and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where these are important resources.

Directions for Use


It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Do not transport or store under 32°F. If freezing occurs, thaw and agitate vigorously.

DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

TO OPERATE: SHAKE BEFORE USING. Turn nozzle to "spray" or "stream" position before using. Hold pump sprayer 18 to 24 inches from foliage or surfaces being treated. Direct the spray so that the upper and lower surfaces are contacted. Use caution on new growth and tender foliage. Repeat treatment as necessary. When using outdoors, spray when air is still.



RIGO AQUEOUS GARDEN SPRAY

Page 3

This insecticide may be used indoors, outdoors or in greenhouses throughout the growing season, or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

To control insects named, apply this spray to provide thorough coverage of upper and lower leaf surfaces. Repeat treatment weekly or as need to control infestations. May be applied on day of harvest.

For Use on Garden Crops:

CROP	INSECTS CONTROLLED
Asparagus	Asparagus Beetles
Beans	Aphids, Leafhoppers, Mexican Bean Beetles, 2-Spotted Cucumber Beetle, Whiteflies
Broccoli, Brussell Sprouts, Cabbage, Cauliflower	Aphids, Cabbage Looper, Crossstriped Cabbageworm, Diamondback Moth larvae, Flea Beetle, Harlequin Bug, Imported Cabbageworm, Stink Bug
Celery	Green Peach Aphid, Cabbage Looper, Leaf-tier
Cranberries	Fireworms, Leafhoppers
Eggplant	Green Peach Aphids, Blister Beetle, Colorado Potato Beetle, Flea Beetles
Lettuce	Green Peach Aphid, Cabbage Looper, Diamondback Caterpillar, Imported Cabbageworms
Collards Kale, Mustard Greens, Turnips	Aphids, Cabbage Looper, Imported Cabbageworms, Diamondback Caterpillar
Peppers	Green Peach Aphid
Potatoes	Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers
Radishes	Aphids, Flea Beetles
Spinach	Aphids, Cabbage Looper, Webworm
Tomatoes	Green Peach Aphid, Colorado Potato Beetle, Flea Beetles, Stink Bugs, Whiteflies

For Use on Ornamentals such as: African Violets, Azaleas, Begonias, Camellias, Dahlias, Dogwood, Roses, Wandering Jew, Asters, Carnations, Chrysanthemums, Geraniums, Gladiolis, Pepperonia, Spirea, Philadendron, Euonymus, Sedum, Ferns, Crabapple Rose, and Wisteria. To Control insects such as: aphids, flea beetles, spider mites, leaf hoppers, whiteflies, mealybugs, Japanese beetles, southern army worms, gypsy moth caterpillars and adults, Eastern and forest tent caterpillars.

To kill gypsy moth caterpillars and adults, adult Japanese beetles and Eastern and forest tent caterpillars on contact:

On Trees and Shrubs: Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.

RIGO AQUEOUS GARDEN SPRAY
Page 4

On Roses and in Flower Gardens: Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

INDOORS AS A SPACE SPRAY

To control house flies, fruit flies, mosquitoes, gnats and small flying moth : apply using a good space sprayer adjusted to deliver a fine mist. Cover or remove exposed food and food handling surfaces. Close windows and doors and shut off ventilating systems. Direct spray toward upper portions of the enclosure filling the room with mist. Use 4 to 5 ounces per 1000 cubic feet of space. Keep area closed for at least 30 minutes. Vacate treated areas and ventilate before reoccupying. Repeat treatment as necessary.

OUTDOORS, ADULT MOSQUITO CONTROL

Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery and vegetation where mosquitoes may rest particularly around the perimeter of mosquito breeding sites such as swamps, marshes, stagnant pools and ponds. Apply at a rate of 5 to 6 quarts per acre (4 ounces per 1000 square feet).

To control mosquito larvae, apply at the rate of 7.75 gallons per acre or 24 fl. oz. per 1000 square feet. Apply as a coarse wetting spray to the border of, or on shallow waters of wetlands, swamps and marshes where mosquito larvae are known to breed.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling are contrary to label instructions.

