

ATTACHMENT 1

A. ACTIVE INGREDIENTS:

Tetramethrin [(1-Cyclohexene-1, 2-dicarboximido) methyl 2, 2-dimethyl-3-(2-methylpropenyl) cyclopropanecarboxylate]	0.200%
Cyano 93-phenoxyphenyl) methyl 4-chloro-alpha-(1- methylethyl) benzeneacetate	0.400%
Inert Ingredients:	99.400%
Total	<u>100.000%</u>

B. Rigo Company
 A Division of Wilbur-Ellis Company
 Buckner, Kentucky 40010

R I G O

AQUEOUS GARDEN SPRAY

FOR INSECT CONTROL ON VEGETABLES AND ORNAMENTAL PLANTS
KILLS GYPSY MOTH CATERPILLARS AND JAPANESE BEETLES
MAY BE APPLIED ON DAY OF HARVEST

ACTIVE INGREDIENTS:

Pyrethrins.....	0.02%	} A
*Piperonyl Butoxide, technical.....	0.20%	
Petroleum Distillate	0.08%	
INERT INGREDIENTS:.....	99.70%	
TOTAL	100.00%	

*Equivalent to min. 0.16% (butylcarbityl) (6-propylpiperonyl) ether and 0.04% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional precautionary statements

Not Reviewed. Registrant claims to be in accordance with registration of Pesticide Product Guidance for Phase 2 response, Page 2.12, Active to Inert Change in Status.

EPA Reg. No. 70-276
EPA Est. No. 70-KY-1

Net Contents
24 Ounces

Manufactured by

RIGO COMPANY

BUCKNER, KENTUCKY 40010 } B

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

C A U T I O N

CAUTION: Harmful if swallowed or absorbed through skin. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists. Remove pets, birds and cover fish aquaria before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams, ponds, tidal marshes and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where these are important resources.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Do not transport or store under 32°F. If freezing occurs, thaw and agitate vigorously.

DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

TO OPERATE: SHAKE BEFORE USING. Turn nozzle to "spray" or "stream" position before using. Hold pump sprayer 18 to 24 inches from foliage or surfaces being treated. Direct the spray so that the upper and lower surfaces are contacted. Use caution on new growth and tender foliage. Repeat treatment as necessary. When using outdoors, spray when air is still.

RIGO AQUEOUS GARDEN SPRAY

Page 3

This insecticide may be used indoors, outdoors or in greenhouses throughout the growing season, or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

To control insects named, apply this spray to provide thorough coverage of upper and lower leaf surfaces. Repeat treatment weekly or as need to control infestations. May be applied on day of harvest.

For Use on Garden Crops:

CROP	INSECTS CONTROLLED
Asparagus	Asparagus Beetles
Beans	Aphids, Leafhoppers, Mexican Bean Beetles, 12-Spotted Cucumber Beetles, Whiteflies
Broccoli, Brussell Sprouts, Cabbage, Cauliflower	Aphids, Cabbage Looper, Crossstriped Cabbageworm, Diamondback Moth larvae, Flea Beetle, Harlequin Bug, Imported Cabbageworm, Stink Bug
Celery	Green Peach Aphid, Cabbage Looper, Leaf-tier
Cranberries	Fireworms, Leafhoppers
Eggplant	Green Peach Aphids, Blister Beetle, Colorado Potato Beetle, Flea Beetles
Lettuce	Green Peach Aphid, Cabbage Looper, Diamondback Caterpillar, Imported Cabbageworms
Collards Kale, Mustard Greens, Turnips	Aphids, Cabbage Looper, Imported Cabbageworms, Diamondback Caterpillar
Peppers	Green Peach Aphid
Potatoes	Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers
Radishes	Aphids, Flea Beetles
Spinach	Aphids, Cabbage Looper, Webworm
Tomatoes	Green Peach Aphid, Colorado Potato Beetle, Flea Beetles, Stink Bugs, Whiteflies

For Use on Ornamentals such as: African Violets, Azaleas, Begonias, Camellias, Dahlias, Dogwood, Roses, Wandering Jew, Asters, Carnations, Chrysanthemums, Geraniums, Gladiolis, Pepperonia, Spirea, Philadendron, Euonymus, Sedum, Ferns, Crabapple Rose, and Wisteria. To Control insects such as: aphids, flea beetles, spider mites, leaf hoppers, whiteflies, mealybugs, Japanese beetles, southern army worms, gypsy moth caterpillars and adults, Eastern and forest tent caterpillars.

To kill gypsy moth caterpillars and adults, adult Japanese beetles and Eastern and forest tent caterpillars on contact:

On Trees and Shrubs: Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.

RIGO AQUEOUS GARDEN SPRAY
Page 4

On Roses and in Flower Gardens: Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

INDOORS AS A SPACE SPRAY

To control house flies, fruit flies, mosquitoes, gnats and small flying moths: apply using a good space sprayer adjusted to deliver a fine mist. Cover or remove exposed food and food handling surfaces. Close windows and doors and shut off ventilating systems. Direct spray toward upper portions of the enclosure filling the room with mist. Use 4 to 5 ounces per 1000 cubic feet of space. Keep area closed for at least 30 minutes. Vacate treated areas and ventilate before reoccupying. Repeat treatment as necessary.

OUTDOORS, ADULT MOSQUITO CONTROL

Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery and vegetation where mosquitoes may rest particularly around the perimeter of mosquito breeding sites such as swamps, marshes, stagnant pools and ponds. Apply at a rate of 5 to 6 quarts per acre (4 ounces per 1000 square feet).

To control mosquito larvae, apply at the rate of 7.75 gallons per acre or 24 fl. oz. per 1000 square feet. Apply as a coarse wetting spray to the border of, or on shallow waters of wetlands, swamps and marshes where mosquito larvae are known to breed.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling are contrary to label instructions.