


72-7

7/2/70

NET WEIGHT 3 POUNDS

P9183

Best Available Image



V-75 DUST

INSECTICIDE

**V-75 DUST**

INSECTICIDE

CONTAINS 0.75% DIFENTHALIN

FOR USE ON VEGETABLES, FRUIT, AND NURSERY PLANTS

**ACTIVE INGREDIENTS:**

Difenthenal	0.75%
Other Extractives (Cube)	1.50%
<b>INERT INGREDIENTS:</b>	<b>97.75%</b>

**CAUTION:** KEEP OUT OF REACH OF CHILDREN  
SEE ADDITIONAL CAUTIONS ON BACK PANEL

Manufactured By  
**MILLER CHEMICAL & FERTILIZER CORPORATION**  
Hanover, Pennsylvania 17331/An Alco Standard Company

V-75 DUST

MILLER

283

V-75 DUST

INSECTICIDE

**DIRECTIONS**

Dust the under surface of leaves as well as top surface. Apply this dust when insects first appear. Repeat applications weekly, or as often as necessary, throughout the growing season.

**BEANS (SNAP AND LIMA):** For Control of Mexican Bean Beetle, Flea Beetle, Cucumber Beetle, Leaf Beetle and Aphid: Apply 1 lb. of Dust per 1,500 sq. ft. when insects are first noted and repeat at 7 to 10 day intervals as needed.

**ASPARAGUS:** For Control of Asparagus Beetle and Japanese Beetle: Apply 1 lb. per 1,500 sq. ft. when insects are first noted and repeat in 7 to 10 days as needed.

**ORNAMENTAL TREES, SHRUBS AND FLOWERS:** For Control of Japanese Beetle: Make first application when insects are noted. Repeat as often as necessary to control insects. Use 1/2 to 3/4 lb. per 1,500 sq. ft. of planted area.

**STRAWBERRIES:** For Control of Spittlebug, Flea Beetle and Leaf Beetle: Apply 1 lb. per 1,500 sq. ft. Make first application early in Spring for Spittlebug control, and repeat every 7 to 10 days as necessary. For Flea Beetle and Leaf Beetle, apply when insects are noted and repeat in 7 to 10 days, as necessary, for control.

**APPLES AND PEACHES:** For Control of Japanese Beetle: Dust foliage lightly, using 1/2 lb. per 1,500 sq. ft. Apply when insects are first noted and repeat as needed.

**GRAPES, BLUEBERRIES, RASPBERRIES, BLACKBERRIES AND DEWBERRIES:** For Control of Japanese Beetle, Raspberry Fruit Fly and Sawfly: Apply when insects are first noted and repeat in 7 to 10 days as needed.

For Control of Raspberry Cane Borer: Apply to foliage just before plants bloom and repeat in 7 to 10 days. Use 1 lb. per 1,500 sq. ft.

**CAUTION**

Do not apply to food crops within 1 day of harvest.  
Keep out of reach of children and pets.

Do not contaminate feed or foodstuffs. (2)

Toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply in any manner or to any area not specified on this label.

The use of this material in a manner or at a time other than in accordance with directions may produce plant injury. Federal Food and Drug Administration has set official tolerances for pesticide residues. Crops having residues in excess of these tolerances should not be marketed without removal of such excess.

The use of this material being beyond our control and involving elements of risk to human beings, animals and vegetation, we do not make any warranty, express or implied, as to the effects of such use, when this product is not used in accordance with the directions as stated on this label.

Manufactured By

**MILLER CHEMICAL & FERTILIZER CORPORATION**

Hanover, Pennsylvania 17331/An Alco Standard Company

119M35A F. N. 1955 USDA Reg. No. 72-7 BATCH NO.

**BEST COPY AVAILABLE**

MILLER

383

ACCEPTED

JUL 2 1970

UNDER THE PROVISIONS OF THE  
FUNGICIDE ACT, 1958 (C. 117)  
FOR FOLIOLOGICAL SOCIETY REGISTERED  
ED UNDER NO. 28-7 SUBJECT  
TO ATTACHED COMMENTS.