

Reg no: 72-156

PM-16

1974

ACCEPTED

AUG 23 1974

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PEST CONTROL REGISTERED
UNDER NO. 72-156 SUBJECT
TO ATTACHED COMMENTS.

MILLER KWIK-TOX INSECT DUST (4% Malathion)

ACTIVE INGREDIENT:

Malathion 4%

INERT INGREDIENTS:

96%
O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate

DIRECTIONS

Application of MALATHION DUST should follow the recommendations of State Agricultural Experiment Stations. Rate of Application: Apply 4 MALATHION DUST at the rate of 20-35 lbs. per acre, depending upon the size of plants and degree of infestation. Repeat application as needed. Cover plants thoroughly. **Apply when foliage is dry or plant injury may occur.**

TIME TO APPLY

Snapbeans and Lima Beans: Apply as directed above for use against Mexican Bean Beetle, Red Spider Mite, Leafhopper, and Aphid. Apply when insects first appear and repeat as needed. Do not apply within 24 hrs. of harvest.

Tomatoes and Potatoes: Apply as directed above for use against Aphid and Red Spider. Begin applications when plants are small (before insects appear) and repeat at 10-14 day intervals through the growing season. Do not apply within 1 day of harvest on tomatoes. No time limit for potatoes. **Peas:** Apply as directed for use against Pea Aphid when insects first appear. Repeat as needed. Do not apply within 3 days of harvest.

Cucumbers, Squash and Melons: For use against Aphids and Spider Mites, apply as directed when insects first appear and repeat as needed. Do not apply within 1 day of harvest.

CAUTION

Harmful if swallowed. Avoid breathing dust. Avoid contact with skin. Wash thoroughly. Avoid contamination of feed and foodstuffs. Avoid contamination of feed, feeding troughs and water fountains.

44MINUS

Use of this material at a time or in a manner other than in accordance with directions may produce plant injury. Federal Food and Drug Administration has set official tolerances for pesticide residues. Crops having excess of these tolerances should not be marketed without removal of such excess.

Do not apply within the specified time limit of harvest. Wash edible portions of fruit after harvest to remove any remaining residues.

This product is toxic to fish. Do not contaminate any body of water. Apply this product only as specified on this label.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning (Keep out of smoke).

Malathion is a cholinesterase inhibitor and can cause symptoms similar to those caused by other organic phosphate compounds, although much larger exposures are required. If poisoning should occur, call a physician. Atropine is antidotal.

The use of this material being beyond our control and involving elements of risk to human beings, animals and vegetation, we do not make any warranty, express or implied, as to the effects of such use, when this product is not used in accordance with the directions on this label.

CAUTION: KEEP OUT OF REACH OF CHILDREN
NET WEIGHT 50 POUNDS

Manufactured By
MILLER CHEMICAL & FERTILIZER CORPORATION
Hanover, Pennsylvania 17331, U.S.A. • An Alco Standard Company

E.N. 2315

EPA Reg. No. 72-156

BATCH No.

EPA Est. No. 12-ME-2



WEATHERION DUST at the rate of 1.0 to 2.0 lbs. per acre depending upon the
degree of infestation. For detailed information needed, Cover
sheet to this page. **Apply when foliage is dry or plant injury may occur.**

TIME TO APPLY

Snapbeans and Lima Beans: Apply when infested by Mexican
Bean Beetle, Leaf Spider, Mite, or thrips. Apply when insects
first appear and repeat as needed. Do not apply within 24 hrs. of harvest.

Tomatoes and Potatoes: Apply as directed at first infestation against Aphid and
Leaf Spider. Best application when plants are small before insects appear.
Repeat at 10-14 day intervals through the growing season. Do not apply
within 1 day of harvest on tomatoes. No time limit for potatoes.

Peas: Apply
as directed for use against Pea Aphid when insects first appear. Repeat as
needed. Do not apply within 3 days of harvest.

**Cucumbers, Squash and
Melons:** For use against Aphid and Spider Mite, apply as directed when
insects first appear and repeat as needed. Do not apply within 1 day of
harvest.

CAUTION

Do not inhale or swallow. Avoid contact with skin. Avoid contact with eyes. Wash
hands after use. Avoid contact with food and feed stuff. Avoid contact with
children and pets. Wash thoroughly with water after use.