

72-157

NET CONTENTS 8 FLUID OUNCES



KWIK-TOX[®] INSECT SPRAY

WARNING: KEEP OUT OF REACH OF CHILDREN
SEE ADDITIONAL CAUTIONS ON SIDE PANEL

Use on plants used for food or forage Use according to label directions.

WARNING: KWIK-TOX INSECT SPRAY is poisonous. Poisoning may occur as a result of exposure. It may be fatal if swallowed. May be absorbed through the skin. Avoid contact with skin, eyes, and clothing. Wash hands thoroughly after handling. Wash contaminated clothing before re-use. Do not breathe spray. Do not contaminate feed or food. Do not contaminate drinking water or fish ponds. Keep out of reach of children.

NOTE: Atropine sulfate and or 2-PAM. KEEP CONTAINER CLOSED AND AWAY FROM HEAT AND OPEN FLAME. Do not re-use empty container. Bury or incinerate in a safe place.

Amount of this material being beyond our control and involving elements of risk to human beings, animals and vegetation, we do not make any warranty, express or implied, as to the effects of such use, when this material is not used in accordance with the directions printed on this label.

FA SYSTOX-R[®] is a registered trademark of Farbwerke Bayer A.G., Chemagro Corporation licensee. DIBANER[®] registered trademark of Rohm & Haas.

Reg. No. 100-1007 Est. No. 150003

5NLS

Read entire label before using. Use strictly in accordance with the directions.

A portion of this insecticide combination is deposited on the plant surfaces to give general contact and residual insecticidal action. Another portion gives systemic action by being absorbed through the plant surfaces into the sap system. This gives longer residual effect and controls many insects not exposed to regular contact sprays.

FOLIAR SPRAY: For Roses, Azaleas, Taxus, Rhododendrons and other flowers, evergreens and ornamental fruit and shade trees, mix 2 tablespoonfuls (1 oz.) in each gallon of water. When insects appear, apply spray to hit as many as possible, and to thoroughly wet all plant surfaces. Repeat weekly on flowers or as necessary on other plants. Early May is the average time for spraying LEAF MINERS on birches, boxwood and holly. Repeat in 10 days.

Plant Insects Controlled on Flowers & Ornamentals:

Aphids	Lace Bug	Certain Scale
Red Spider Mites	Plant Bugs	Crawlers
Leaf Hoppers	Japanese	Mites
Rose Chafers	Beetle	Thrips

LAWN SPRAY: For ANTS, CLOVER MITES and certain other lawn insects, water thoroughly after treating to wash insecticide into soil. Dilution: 2 tablespoonfuls per gallon of water applied to each 35 sq. ft. Keep pets and children off treated area until spray dries.

PATIO SPRAY: For TICKS, CHIGGERS and to aid MOSQUITO CONTROL around the patio and in picnic areas, spray hiding places, lawns, shrubs and surrounding grass and brush. Do not spray patio surfaces. Keep children and pets out of treated area until spray dries. Dilution: 3 tablespoonfuls per gallon of water applied to each 100 sq. ft.

NET CONTENTS 8 FLUID OUNCES

