



FLY AWAY SPRAY

READY-TO-USE

ACCEPTED
SEP 23 1985
U.S. Environmental Protection Agency
as amended, for the pesticide
registered under
EPA Reg. No. 72-279

ONLY FOR SALE TO, USE AND STORAGE BY SERVICE PERSONS
(Statement For Sizes Over One Gallon)

For Insect Control In Food Processing Plants, Meat Packing Plants, Dairy Barns
And Grain Storage Bins.

ACTIVE INGREDIENTS:	BY WEIGHT
Pyrethrins.....	0.1%
Piperonyl Butoxide, Technical*.....	1.0%
Isoparaffinic Petroleum Hydrocarbons.....	98.9%
TOTAL.....	100.0%

*Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% of other related compounds.

EPA Reg. No. 72-279

EPA Est. No. 72-PA-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

CAUCIÓN

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED: Call a physician or Poison Control Center. Do not induce vomiting. Vomiting may cause aspiration pneumonia. Do not induce vomiting or give anything to an unconscious or convulsing person.
- IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.
- IF ON SKIN: Remove contaminated clothing. Immediately wash affected area with soap and water. Launder clothing before reusing. Get medical attention if irritation persists.
- IF INHALED: Remove victim to fresh air. Apply respiration if indicated.
- NOTE TO PHYSICIAN: Empty stomach if material was taken internally. Gastric lavage may be indicated. If symptoms of cholinesterase inhibition are present, atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

MILLER CHEMICAL & FERTILIZER CORPORATION
Hanover, Pennsylvania 17331

NET CONTENTS: 1 GALLON LIQUID
5 GALLONS LIQUID

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with eyes, skin and clothing. Avoid breathing vapors - spray mist. Wash thoroughly after handling. Wash hands, face and arms with soap and water before eating and smoking. Wear gloves and long-sleeved work clothing. Change to clean clothing daily. Wash all clothing with soap and water before reuse. Do not remain in treated area. Ventilate the area thoroughly before re-entry. Do not apply directly on very young animals. Do not use in poultry houses. Remove pets, birds and cover fish aquariums before spraying. Do not apply in a food processing plant while plant is in operation. When applied in a food processing plant, remove or protect food during treatment. Facilities and equipment must be thoroughly washed with an acceptable detergent solution and then rinsed with potable water to remove all traces of contamination.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARD

Do not use or store near heat or open flame. Keep Container Closed.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL (for one gallon size)

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store in tightly closed, original container in a locked area away from children and domestic animals.
FOR DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash. Do not reuse empty container.

STORAGE & DISPOSAL (for sizes over one gallon)

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store in tightly closed, original container in a locked area away from children and domestic animals.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not reuse empty container.

DIRECTIONS

FOOD PROCESSING PLANTS: For Use as a Space Spray for Control of Fruit Flies, House Flies, Gnats, Flying Moths, Hornets and Skipper Flies: Use 4 oz. per 1,000 cubic feet, using an Electric Sprayer which produces a FINE SPRAY. This method of application is excellent for the treatment of the following large enclosed areas: breweries, warehouses, food plants, bottling plants, dairy barns and flour mills. Do not remain in treated areas and ventilate the area after treatment is completed. Spray will be more effective if plant walls and floors are clean. For Control of the following CRAWLING INSECTS: Ants, Crickets, Roaches, Spiders and Cheese Mites: Apply as a COARSE SPRAY, directed into infested cracks and hiding places of these crawling insects and contacting the insects directly with the spray whenever possible. Spray will be more effective if plant walls and floors are clean. One Gallon of spray should be applied to each 750 sq. ft. of wall or floor surface. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatments or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation.

MEAT PACKING PLANTS: Same conditions and amounts as given above. Can be used where residual-type sprays are unlawful.

DAIRY MILKROOMS AND DAIRY BARNs: For Control of House Flies, Stable Flies, Gnats and Mosquitoes: For best results close doors and windows and use as space spray between milkings. When using in the milk-room, care should be taken not to contaminate milk.

LIVESTOCK SPRAY: Dairy & Beef Cattle, Swine and Sheep: To protect against Horn Flies, Mosquitoes, Gnats and Midges: Spray dairy cattle before each milking. For other types of livestock, spray daily or as needed. Use approximately 2 ounces per head. Apply only as a light mist, spray over the animals' coat. Avoid wetting the hides of animals. Do not exceed a dosage of 2 ozs. of spray per animal per application.

FOR SPRAYING STORAGE BINS: For Control of exposed stages of Rice Weevil, Granary Weevil, Confused Flour Beetle, Lesser Grain Borer and Saw-Toothed Grain Beetle: Clean bins thoroughly before filling with grain and thoroughly wet walls, ceilings, partition boards, etc. One gallon covers about 500 to 800 sq. ft. of surface.

LIMITED WARRANTY: The use of this material being beyond our control and involving elements of risk to human beings, animals and vegetation, we do not make any warranty, express or implied, as to the effects of such use, when this product is not used in accordance with the directions as stated on this label.