



NET CONTENTS
1 GALLON LIQUID

STRAWBERRIES: For Control of Cyclamen Mite, Meadow Spittlebug and Strawberry Aphid: Spray Treatment: 100 gallons of water, apply 40 gallons of spray per acre. Spray plants thoroughly. This spray will control the insects when insects first appear and repeat as required. For multiple applications during the fruiting period, do not apply less than 35 days. Do NOT apply within 4 days of picking.

TOBACCO: (Other than Shade Grown): For Control of Flea Beetle, Hornworm, Budworm and Green Peach Aphid: Apply when insects first appear and repeat as required. Thorough coverage is essential to good control. **Seed Bed:** Use 1 quart per 100 gallons of water and apply about 6 gallons of spray per 100 sq. yds. **Field:** Use 1 to 2 quarts per acre and apply in sufficient water for uniform coverage. Do not apply within 5 days of harvest.

COTTON: For Control of Boll Weevil, Cotton Stink Bugs, Lygus Bug, Bollworm, and Cabbage Looper: For Boll Weevil, use 1 to 2 1/2 quarts per acre, and for other insects, use 2 quarts per acre, in sufficient water to cover. Begin applications when an infestation reaches the point where control is required. Apply at 3 to 5 day intervals, depending upon the severity of the infestation and local conditions. Thorough coverage is essential for good control. Do not apply after bolls begin to open. Do not feed treated plants to meat or dairy animals. Do not graze meat or dairy animals in treated fields.

ORNAMENTALS (Greenhouse and out-of-doors): For Control of Aphids and Cyclamen Mite: Use 1 quart in 100 gallons of water, or 2 teaspoons in 1 gallon of water. Apply when insects are first noted and repeat at 7 to 10 day intervals as necessary. May cause severe injury on birch trees. **NOTE:** In Greenhouse applications, wear a mask or respirator passed by U. S. Department of Agriculture for Thiodan protection.

WARNING

Hazardous if Swallowed, Inhaled, or Absorbed Through Skin! Do not breathe spray mist. Do not get in eyes, on skin or on clothing. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking. Wear clean clothing. During commercial or prolonged exposure in spray mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type passed by the U. S. Department of Agriculture for Thiodan protection. Do not apply or allow to drift to occupied by unprotected humans or beneficial animals.

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning. (Keep out of smoke.) Apply this product only as specified on this label. Do not apply when weather conditions favor drift from areas treated.

CAUTION

Workers entering treated areas within 24 hours of application should wear protective clothing.

WARNING! FLAMMABLE!

KEEP AWAY FROM HEAT AND OPEN FLAME. KEEP CONTAINER CLOSED. LEAKING PACKAGES SHOULD BE REMOVED TO A SAFE PLACE. DO NOT STORE AT TEMPERATURES BELOW 15° F. DO NOT USE IN UNDILUTED FORM.

The use of this material in a manner or at a time other than in accordance with directions may produce plant injury. Federal Food and Drug Administration has set official tolerances for pesticide residues. Crops having residues in excess of these tolerances should not be marketed without removal of such excess.

The use of this material being beyond our control and involving elements of risk to human beings, animals and vegetation, we do not make any warranty, express or implied, as to the effects of such use, when this product is not used in accordance with the directions as stated on this label.

THIODAN E.C. #2

**WARNING: KEEP OUT OF REACH OF CHILDREN
SEE ADDITIONAL WARNING ON SIDE PANEL**