

SPECIMEN LABEL
NOT FOR RESALE

BACK RUBBING DEVICES: For reduction of Lice and Hornflies:
Beef Cattle: Apply 5 pints per 33 gallons of Fuel Oil to Back Rubbing Devices (One Back Rubbing Device to each 35 or 41 head of cattle, these rubbers should be continuously accessible). Do NOT use on dairy animals.
SWINE: For Control of Lice. Use 5 pints per 100 gallons of water. Apply as a complete coverage to animals, pens and litter. One treatment may be sufficient. Repeat application only if needed.
POULTRY: CHICKENS, DUCKS, GEESE AND TURKEYS: Direct application, for Control of Northern Fowl Mite, Poultry Lice, and as a supplement to premise treatments for Chicken Red Mite. Mix 1½ tablespoonsful per 1 gallon, or 3½ ounces per 5 gallons, or 4 pints per 100 gallons of water. Spray 1 gallon of dilute mixture per 100 to 150 birds. Repeat in 4 to 6 weeks if necessary.
POULTRY HOUSES: SPRAYS: For Control of Northern Fowl Mite, Chicken Red Mite, Poultry Lice and Flies. Use 2½ tablespoonsful per 1 gallon, or 6 ounces per 5 gallons, or 1 gallon per 100 gallons of water. Spray walls, ceilings, roosts, nests and adjacent areas liberally, taking care to force the spray into all cracks and crevices. For control of Poultry Ticks. Use 3½ tablespoonsful per 1 gallon, or 3 gallons per 100 gallons of water, apply liberally to walls, ceilings, and adjacent areas, taking care to force the spray into all cracks and crevices. Repeat as necessary for control. Use a high pressure for surface sprays.
ROOST PAINT: For the Control of Chicken Red Mite and Poultry Lice. Use 2½ tablespoonsful per gallon of water. Brush on this mixture as a roost paint at the rate of 1 pint per 150 feet of roost. Repeat as necessary. Avoid contamination of feed, feeding troughs and water fountains with spray.
MOSQUITOES, FLIES AND SMALL FLYING INSECTS: For the Control of Mosquitoes, Flies and Small Flying Insects in and around buildings which house domestic animals. Apply a spray containing 1½ gallons in 100 gallons, or 2½ tablespoonsful per gallon of water. Apply the spray at the rate of 1 gallon per 1000 sq. ft. on painted surfaces and 2 gallons per 1000 sq. ft. on unpainted surfaces. Repeat application as necessary. Do NOT apply LIQUID 87 MALATHION within 14 days to freshly whitewashed surfaces. Do NOT use in milkrooms. Remove milking animals and calves under one month of age from buildings before treating.
MOSQUITO LARVAE IN STANDING WATER: Use ½ pint per acre in sufficient water or oil to obtain uniform coverage when applied by air or ground equipment. Repeat application as necessary. Avoid applying oil sprays to valuable ornamental plants, as injury may occur. Malathion may be toxic to certain species of fish, particularly in shallow water.
DOG AND CAT QUARTERS: For Control of Fleas in kennels, pens, yards, lawns and under house. Remove manure or debris. Use 3½ ounces per gallon of water and spray at the rate of 1 gallon per 1000 sq. ft. Use garden sprayer. Avoid contamination of feed, feeding and water containers.
DOGS AND CATS: For Control of Fleas. Use 1½ tablespoonsful in 1 gallon of water. Apply with hand sprayer. Wet animal thoroughly by spraying. Re-treat every three weeks if dogs or cats become re-infested.


CAUTION

Do NOT apply to Beans within 24 hours of harvest, Apples, Asparagus, Broccoli, Peas and Tomatoes within 3 days; Corn within 5 days of harvest or forage, Peaches, Brussels Sprouts, Cabbage, Cauliflower, Kale and other Cole Crops within 7 days, or to Alfalfa within 7 days of harvesting or pasturing.
Harmful if swallowed. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly with soap and water after using. Avoid contamination of feed, foodstuffs, feeding and water containers. Keep out of reach of children and domestic animals. Malathion is a cholinesterase inhibitor and can cause symptoms similar to those caused by other organic phosphate compounds, although much larger exposures are required. If poisoning should occur, CALL A PHYSICIAN. Atropine is antidotal.
The use of this material in a manner or at a time other than in accordance with directions may produce plant injury. Federal Food and Drug Administration has set official tolerances for pesticide residues. Crops having residues in excess of these tolerances should not be marketed without removal of such excess.
Toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not use where fish are an important resource.
The use of this material being beyond our knowledge and control and involving elements of risk to human beings, animals and vegetation, we do not make any warranty, express or implied, as to the effects of such use, whether or not in accordance with directions or claimed so to be.

98M1NLS

USDA Reg. No. 72-313

56



BEST DOCUMENT AVAILABLE



NET CONTENTS
1 GALLON LIQUID

MALATHION "8" E.C.

**CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE ADDITIONAL CAUTIONS ON SIDE PANEL**

For Small Amounts: 1 tsp. per gal. is equal to 1 pt. per 100 gals. of water. Use 1 g.

DIRECTIONS

MILLER MALATHION "8" contains approximately 0.13% of Malathion per gallon. For

1 fluid ounce
1/2 pint = 1 fluid ounce
1 pint = 2 fluid ounces

Mix MALATHION "8" into spray tank by pouring a measured amount into the full tank. Make first application when insects are first noted and repeat application in 7 to 10 days. State may issue recommendations as to proper timing of sprays to control specific insect pests.

APPLES AND PEACHES: For Control of Spider Mites, European Red Mite, Two-spotted Spider Mite and Williamsite. Use at least two applications 10 to 12 days apart during summer.

APPLES: For Control of Felted Scale, Woolly Apple Aphid, Green Aphid and Rosy Aphid. Apply 1 to 1 1/4 pints in 25 or more gallons of water per acre for uniform coverage. Repeat harvest or pasturing.

ALFALFA AND CLOVER: For Control of Aphid, Potato Leafhopper, Spittlebugs, Spider Mites and European Red Mite. Apply 1 1/4 pints in 25 or more gallons of water per acre for uniform coverage. Repeat harvest or pasturing.

ASPARAGUS: For Control of Asparagus Beetle. Apply 1 1/4 pints in 25 to 100 gallons of water per acre. Repeat as needed.

BEANS: Snap and Lima: For Control of Mexican Bean Beetle, Spider Mites and Leafhoppers. Repeat as needed.

CABBAGE, CAULIFLOWER, BROCCOLI, KALE AND OTHER COLE CROPS: For Control of Cabbage Worm. Apply 1 1/4 pints per 100 gallons of water per acre. Repeat as needed.

CORN (Field and Sweet): For Control of Sap Beetle and Thrip. Apply 1 1/4 pints in 25 to 100 gallons of water per acre. Do not apply within 5 days of harvest or forage use.

PEAS: For Control of Pea Aphid. Apply 1 pint per 100 gallons of water per acre. Repeat as needed.

POTATOES: For Control of Leafhopper and Aphid. Make one or more applications as needed. Apply 1 1/4 pints per 100 gallons of water per acre.

TOMATOES: For Control of Aphid, Russett and Spider Mites. Make one or more applications as needed. Apply 1 1/4 pints per 100 gallons of water per acre.

ORNAMENTALS: AZALEAS: For Control of Spider Mite and White Flies. Use 1 pint of Aphid, Japanese Beetle, Tarnished Plant Bug and Rose Leafhopper. Apply as a spray when insects appear and repeat at 7 to 10 day intervals if necessary.

FOR CONTROL OF SOFT BROWN SCALE, OYSTER SHELL SCALE, JUNIPER SCALE, OAK SCALE AND OTHER SCALE INSECTS: Use 2 1/2 pints per 100 gallons of water. Make one or more applications of a spray containing 1 1/4 pints per 100 gallons of water. Make applications as necessary, when crawlers have settled on foliage.

NOTE: Injury may occur on certain ferns, including Boston, Maiden Hair and Pteris.

DIRECT APPLICATION TO LIVESTOCK

CATTLE: For Control of Lice on Beef Cattle. Use 5 pints per 100 gallons or 1 1/2 tables per acre. Repeat application in 14 days if necessary. Do NOT apply to dairy animals. Do not contaminate feed, food containers and watering troughs when applying spray.

F.N. 3071

BATCH NO.

**TENTS
LIQUID**

For small amounts of spray use 1 quart to 1 pt. per 100 gals. of water. Use 1 gal. more spray for 100 gals. of water.

APPLES AND PEACHES: Two or three applications of spray are necessary. The first application should be made when the fruit is about half grown. The second application should be made when the fruit is about three-quarters grown. The third application should be made when the fruit is about full grown. The fourth application should be made when the fruit is about full grown. The fifth application should be made when the fruit is about full grown. The sixth application should be made when the fruit is about full grown. The seventh application should be made when the fruit is about full grown. The eighth application should be made when the fruit is about full grown. The ninth application should be made when the fruit is about full grown. The tenth application should be made when the fruit is about full grown.

ALFALFA AND CLOVER: Apply 10 to 20 gals. of spray per acre. Repeat application as necessary.

ASPARAGUS: For control of Asparagus beetle. Apply 10 to 20 gals. of spray per acre. Repeat application as necessary.

BEANS: Snap and Lima. For control of bean beetle. Apply 10 to 20 gals. of spray per acre. Repeat application as necessary.

CABBAGE, CAULIFLOWER, BROCCOLI, KALE AND OTHER COLE CROPS: Apply 10 to 20 gals. of spray per acre. Repeat application as necessary.

CORN: Field and Sweet. Apply 10 to 20 gals. of spray per acre. Repeat application as necessary.

PEAS: For control of pea aphid. Apply 10 to 20 gals. of spray per acre. Repeat application as necessary.

POTATOES: For control of Colorado potato beetle. Apply 10 to 20 gals. of spray per acre. Repeat application as necessary.

TOMATOES: For control of tomato worm. Apply 10 to 20 gals. of spray per acre. Repeat application as necessary.

ORNAMENTALS: AZALEAS: For control of azalea scale. Apply 10 to 20 gals. of spray per acre. Repeat application as necessary.

ROSES: For control of rose beetle. Apply 10 to 20 gals. of spray per acre. Repeat application as necessary.

FOR CONTROL OF SOFT BROWN SCALE, OYSTER SHELL SCALE, UNIPER SCALE, OAK KERMES, FLETCHER SCALE AND BAGWORMS: Apply 10 to 20 gals. of spray per acre. Repeat application as necessary.

FOR CONTROL OF PINE NEEDLE SCALE: Apply 10 to 20 gals. of spray per acre. Repeat application as necessary.

NOTE: Injury may occur on certain trees, shrubs and plants. Use with caution.

F.N. 3071 BATCH NO.

BEST DOCUMENT AVAILABLE