

PRECAUTIONARY STATEMENT
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

ENVIRONMENTAL HAZARD

PHYSICAL HAZARD

DIRECTIONS FOR USE

STORAGE & DISPOSAL



MALATHION

An Emulsifiable Liquid Agricultural Insecticide Containing 50% Malathion

ACTIVE INGREDIENTS:
Malathion (O, O dimethyl dithiophosphate of diethyl malathion)
INERT INGREDIENTS

KEEP OUT OF REACH OF
CAUTION

See side panel for additional cautions

EPA Reg. No. 72-313

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or poison control center immediately. Do not induce vomiting unless directed to do so by a physician. Give nothing by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. If breathing is difficult, give oxygen.

IF ON SKIN: Remove contaminated clothing and wash affected area thoroughly with soap and water.

IF IN EYES: Flush with plenty of water. Call a physician.

NOTE TO PHYSICIAN: Malathion may cause cholinesterase inhibition. Give adjunct to atropine.

ONLY FOR SALE TO, STORAGE, AND USE BY PROFESSIONAL SERVICE PERSONNEL

Manufactured By


MILLER CHEMICAL & FERTILIZER CO.
Hanover, Pennsylvania 17331/An Alco Station

NET CONTENTS 1 5 55

APPLES AND PEACHES
APPLES
ALFALFA AND CLOVER
ASPARAGUS
BEANS (Snap and Lima)
BLUEBERRIES, BLACKBERRIES, RASPBERRIES, STRAWBERRIES
GROUND APPLICATION
RASPBERRIES AND STRAWBERRIES
AERIAL APPLICATION

AERIAL APPLICATION

Blackberries, Blueberries



BEST DOCUMENT AVAILABLE



ALATHION "8" E.C.

Agricultural Insecticide Containing Approximately 8 lbs. Malathion Per Gallon

DIESTERS.	
O-dimethyl dithiophosphate of diethymercaptosuccinate	60
DIESTERS	20
Total!	100

KEEP OUT OF REACH OF CHILDREN
CAUTION

See side panel for additional cautions

13 EPA Est. No 72 PA 1

STATEMENT OF PRACTICAL TREATMENT

...an or poison control center immediately. Do not induce vomiting. Do not give any
...to an unconscious person.
...fresh air.
...nated clothing and wash affected area with soap and water. Launder clothing before

of water. Call a physician

on may cause cholinesterase inhibition Atropine is antidotal 2 PAM may be an effec

DO NOT USE BY PROFESSIONAL SERVICE PERSONS OR FOR AGRICULTURAL USES

Manufactured By

ER CHEMICAL & FERTILIZER CORPORATION
Mer, Pennsylvania 17331/An Alco Standard Company

NET CONTENTS	1	5	55	GALLONS
1				
2				
3				
4				
5				
6				
7				
8				
9				
10				
11				
12				
13				
14				
15				
16				
17				
18				
19				
20				
21				
22				
23				
24				
25				
26				
27				
28				
29				
30				
31				
32				
33				
34				
35				
36				
37				
38				
39				
40				
41				
42				
43				
44				
45				
46				
47				
48				
49				
50				
51				
52				
53				
54				
55				
56				
57				
58				
59				
60				
61				
62				
63				
64				
65				
66				
67				
68				
69				
70				
71				
72				
73				
74				
75				
76				
77				
78				
79				
80				
81				
82				
83				
84				
85				
86				
87				
88				
89				
90				
91				
92				
93				
94				
95				
96				
97				
98				
99				
100				

CABBAGE CAULIFLOWER BROCCOLI BRUSSELS SPROUTS KALE AND OTHER COLE CROPS
CORN (field and Sweet)
MUSHROOMS PHORID SCARID FLIES
PEAS
PEAS PEA APPL. 3 APPL. 3
POTATOES
TOMATOES
ORNAMENTALS* AZALEAS
ROSES

DIRECT APPLICATION TO LIVESTOCK

CATTLE - The cattle population in the United States is estimated at 97 million head as of January 1, 1968. The average annual slaughter rate is approximately 10 million head. The total number of cattle slaughtered in 1967 was 10,000,000 head. The total number of cattle slaughtered in 1968 was 10,000,000 head.

BACK RUBBING DEVICES Forming all types of Back Rubbing Devices per Volume 1 of Rule 6 - Back Rubbing Device, each 15¢ plus shipping and handling charges. \$4.95 per dozen. U.S. Patent 2,870,000.

SWINE
Release: 01/01/01

[illegible][illegible][illegible][illegible]

MOSQUITO LARVAE IN STANDING WATER Use DDT spray in sufficient amount to cover the entire surface of the water and adjacent shore. Repeat applications as necessary. Avoid spraying during the day. **CAUTION:** Do not spray on plants, animals, or people. Do not spray on water used for drinking.

DOGS AND CAT QUARTERS. For Comfort Feeds Use 1 quart of water and 1/2 cup of feed for each dog and 1/4 cup of water and 1/8 cup of feed for each cat. Water and feed should be given at the same time. For Comfort Feeds Use 1 quart of water and 1/2 cup of feed for each dog and 1/4 cup of water and 1/8 cup of feed for each cat. Water and feed should be given at the same time.

The use of this material is intended for the purpose of providing a general overview of the subject matter and is not intended to be used as a substitute for professional advice or as a basis for any legal or other action. The user assumes all responsibility for the use of this material and for the accuracy and completeness of the information contained herein.