

MILLER 10% CHLORDANE GRANULAR

FOR CRABGRASS SUPPRESSION AND FOR CONTROL OF LARVAE OR GRUBS OF JAPANESE BEETLE, ORIENTAL BEETLE, EUROPEAN CHAFER AND MAY BEETLE IN LAWNS AND FLOWER BEDS

(This product is formulated on 15/30 Attaclay Granules)

CORN (Field, Sweet and Popcorn): For Control of Corn Rootworms (Northern and Southern), Corn Field Ants, Corn Root Aphid, Seed Corn Beetle Larvae, and Seed Corn Beetle Maggot: Use only at pre-planting or planting time. Can be used on corn intended for use as silage. Apply pre-plant incorporated: 25-40 lbs./ A broadcast. For band application at planting time, use 25 lbs. /Acre.

ACTIVE INGREDIENT:

Technical Chlordane*10%

INERT INGREDIENTS

.....90%

*Equivalent to 6% Octachloro-4, 7-methanotetrahydroindane and 4% related compounds

DIRECTIONS

CRABGRASS SUPPRESSION ON LAWNS AND ORNAMENTAL TURF: Apply 600 lbs. per acre, or 15 lbs. per 1,000 sq. ft. of Lawn or Turf area to suppress crabgrass seedlings. Application should be made before crabgrass seed germinate. This product is not effective against established crabgrass. Do not use on newly seeded lawns.

Time of Application for Crabgrass Seedling Control: Apply early Spring, during period April 15 to May 1; apply with fertilizer spreader. After applying, use sprinkler or hose to wash granules off grass into the soil.

GRUB-PROOFING LAWNS, ORNAMENTAL TURF AREAS, NURSERY AND FLOWER BEDS: For Control of Larvae or Grubs of Japanese Beetle, Oriental Beetle, Masked Chafer, European Chafer, May Beetle, and Rose Chafer and for Controlling Wire Worms and Ants: Apply 100 lbs. per acre, or approximately 2½ lbs. per 1,000 sq. ft. After applying, use sprinkler or hose to wash off grass and into the soil.

When treating Nursery and Flower Beds, apply 10% Chlordane Granular, and work into the soil at depth of 2 to 4 inches prior to planting.

FOR CONTROL OF TICKS, CHIGGERS, ANTS, SPIDERS, CRICKETS AND ARMY WORMS AND FOR REDUCTION IN ANNOYANCE FROM MOSQUITOES ON LAWNS, ORNAMENTAL TURF AND AROUND SHRUBS: Apply 50 lbs. per acre, or approximately 1 lb. per 1,000 sq. ft. For best results, apply to grass when it is wet from dew, rain or sprinkling. Repeat treatment at 7 to 10 day intervals, as necessary, to control insects. CAUTION: Do not use on newly seeded lawns.

44MINLS

E.P.A. Reg. No. 72-330
E.P.A. Est. No. 72-MD-2

F.N. 25079


CAUTION

Store out of reach of children and pets. Wear natural rubber gloves when handling this product. Contact with skin can cause toxic symptoms. Avoid inhalation and skin contact. In case of spillage on skin, wash immediately with soap and water. Avoid contamination of feed and foodstuffs. Keep children and pets off treated areas until this material has been thoroughly washed into the soil and the grass has completely dried.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

The use of this material being beyond our control and involving elements of risk to human beings, animals and vegetation, we do not make any warranty, express or implied, as to the effects of such use, when this product is not used in accordance with the directions as stated on this label.

CAUTION: KEEP OUT OF REACH OF CHILDREN



NET WEIGHT 50 POUNDS

Manufactured By
MILLER CHEMICAL & FERTILIZER CORPORATION
17331, U.S.A. • An Alco Standard Company

BATCH No.

72-330