

DIRECTIONS CONTINUED

BEEF CATTLE, NON-LACTATING DAIRY CATTLE -- Horn Flies, Lice; SWINE -- Lice: Apply as dip or spray .5% Solution (2 gallons of **METHOXYCHLOR** \approx 2 E. C. per 100 gallons of water). Spray 2 quarts per head on large animals and proportionally less for smaller animals. Make application every 3 weeks or as needed. Do Not apply to Dairy Cattle during lactation or within 2 weeks of freshening. Do Not dip calves under one month of age.

SHEEP AND GOATS: Lice: Apply as a .25% Dip or .5% Spray. (To make a .25% Dip Solution, mix 1 gallon of **METHOXYCHLOR** \approx 2 E. C. per 100 gallons of water; for .5% Spray Solution, mix 2 gallons of **METHOXYCHLOR** \approx 2 E. C. per 100 gallons of water). When spraying, wet animals thoroughly. Do Not apply to lactating goats.

For ground application on vegetable and forage crops, apply recommended rate in 50 to 100 gals. of water per acre, depending on the amount of crop growth. For aerial application, apply in 3 to 10 gals. of water per acre, depending on crop growth.

CAUTION

Avoid Contact with Skin. In case of contact, wash with soap and water. Avoid contamination of feed and foodstuffs. Do Not use on humans. Harmful if swallowed. Avoid breathing spray mist.

Do not use, pour, spill, or store near heat or open flame.

The use of this material being beyond our control and involving elements of risk to human beings, animals and vegetation, we do not make any warranty, express or implied, as to the effects of such use, when this product is not used in accordance with the directions as stated on this label.

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift of spray from areas treated. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes. This product is toxic to bees and should not be applied when bees are actively visiting the area.

FLAMMABLE LIQUID

Flash Point 78° F. Tag Closed Cup Method

GENERAL DIRECTIONS

MILLER METHOXYCHLOR #2 E. C. is an Emulsifiable Concentrate Insecticide containing 2 pounds Actual Methoxychlor per gallon. It is designed to be mixed with water for control of certain insects. Use sufficient water to give thorough coverage of plants when insects first appear.

DIRECTIONS

Alfalfa, Clover: Alfalfa Caterpillar, Clover Leaf Weevil, Potato Leaf Hopper: Apply 1½ to 2 quarts per acre; **Alfalfa Weevil Larvae:** Apply 2 to 3 quarts per acre; **Meadow Spittlebug:** Fall Treatment -- Apply 3 quarts per acre. Spring Treatment -- Apply 2 quarts per acre. Do Not apply within 7 days of harvest.

Apples -- APPLE MAGGOT; Cherries -- CHERRY FRUIT FLIES, CHERRY FRUITWORM; Peaches, Pears, Plums -- TENT CATERPILLARS, CODLING MOTH, JAPANESE BEETLE, PLUM CURCULIO: Apply 2 to 3 quarts per 100 gallons of water. Do Not apply to Apples, Cherries, Peaches and Plums within 21 days of harvest. Do Not apply to Pears within 7 days of harvest.

Grapes -- GRAPE BERRY MOTH; Strawberries -- STRAWBERRY WEEVIL; Blackberries, Blueberries, Raspberries -- JAPANESE BEETLE, LEAFHOPPERS, PLUM CURCULIO, ROSE CHAFER, SPITTLEBUG: Apply 2 to 3 quarts per 100 gallons of water per acre. Do Not apply within 14 days of harvest.

Soybeans: Velvetbean Caterpillar: Apply 2 quarts per acre. **Mexican Bean Beetle:** Apply 3 quarts per acre. Do Not apply within 7 days of harvest.

Beans: Corn Earworm, Flea Beetle, Japanese Beetle, Mexican Bean Beetle, Potato Leafhopper: Apply 3 quarts per acre. Do Not apply within 3 days of harvest.

Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Eggplant, Kale, Peppers: Flea Beetle: Apply 3 quarts per acre. Do Not apply within 14 days of harvest to Broccoli, Brussels Sprouts, Collards, Kale and Peppers; within 3 days of harvest to Cabbage; and within 7 days of harvest to Cauliflower and Eggplant.

Cantaloupe, Cucumber, Pumpkin, Squash, Watermelon: Cucumber Beetle: Apply 3 quarts per acre. Do Not apply within 1 day of harvest.

Corn: Japanese Beetle: Apply 3 quarts per acre. Do Not apply within 7 days of harvest.

Peas: Alfalfa Looper: Apply 4 quarts per acre. **Pea Weevil:** Apply 2 to 3 quarts per acre. Do Not apply within 7 days of harvest.

Potatoes: Potato Leafhopper: Apply 4 quarts per acre.

Spinach: Alfalfa Looper: Apply 5 pints per acre. Do Not apply within 21 days of harvest.

Tomatoes: Blister Beetle: Apply 3 quarts per acre. Do Not apply within 7 days of harvest.

15M2NLS

EPA Reg. No. 72-341

F.N. 2365

BATCH NO.

EPA Est. No. 72-PA-1

MILLER

NET CONTENTS
5 GALLONS LIQUID

METHOXYCHLOR #2 E. C.

**CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE ADDITIONAL CAUTIONS ON SIDE PANEL**

or spray
er head
or as
calves

mix 1
lons of
Do Not

gals. of
gals. of

ed and

beings
ch use,

nditions
y bury-
oles in
ees are