

**ACCEPTED**  
**JAN 2, 1969**

UNDER THE FEDERAL INSECTICIDE  
 ACT OF 1947, AS AMENDED BY ACT  
 OF OCTOBER 3, 1958, IN REGISTRATION  
 NUMBER **72-500** SUBJECT  
 TO ATTACHED COMMENTS.

hoppers, Lace Bugs, Flea Beetles, Aphid, Bagworm, Rose-slug, Scale Insects, Thrips (exposed). Apply 4 standard teaspoonfuls of concentrate per gallon of water when insects or their damage appears. Re-apply as necessary when dried powder no longer is apparent. Completely cover all parts of plants, including underside of leaves and branches. Repeat application in 7 to 10 days, as necessary.

**VEGETABLES:** Beans, Corn, Pepper, Tomato and Potatoes. To control Potato Beetle, Mexican Bean Beetle, Flea Beetles, Japanese Beetle, Army Worms, Tomato Hornworm, and Tomato Fruit Worm. Mix 4 to 6 standard teaspoonfuls in each gallon of water. Use 1 gallon of dilute spray to 400 sq. ft. Apply when insects or damage appears and repeat in 7 to 10 days, as necessary. **POTATOES:** Foliage application only.

**KEEP OUT OF REACH OF CHILDREN AND ANIMALS. HARMFUL IF INHALED OR SWALLOWED!** Do Not Breathe Spray Mist. Do Not Take Internally. Avoid Prolonged or Repeated Breathing of Spray. Skin Contact May Be Harmful. Avoid Unnecessary Contact. Wash Hands Before Eating. Take Shower or Bath After Work. Wear Regular Long-Sleeved Work Clothing. **CHANGE TO CLEAN CLOTHING DAILY.**

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

**Avoid contamination of foodstuffs.** Avoid contamination of feed, feeding troughs and watering receptacles.

The use of this material being beyond our control and involving elements of risk to human beings, animals and vegetation, we do not make any warranty, express or implied, as to the effects of such use, when this product is not used in accordance with the directions as stated on the label.

**SMALL MEASUREMENTS**

One ounce = 2 standard tablespoons or 6 standard teaspoons.  
 One cup = 8 oz. or 16 standard tablespoons.

FORMULAS: USDA Reg. No. 72-500  
 E.N. X-250 BATCH NO. 119003

**CONTAINS 2 LBS. SEVIN®  
 INSECTICIDE PER GALLON**



**NET CONTENTS  
 6 FLUID OUNCES**

**ACTIVE INGREDIENT:**  
 Carbaryl (1-Naphthyl Methylcarbamate) ..... **21.50%**  
**INERT INGREDIENTS:** ..... **78.50%**

**Do not store below 32° F.**

Manufactured By  
**MILLER CHEMICAL & FERTILIZER CORPORATION**  
 Lancaster, Pennsylvania 17301, U.S.A. • Subsidiary of The Standard Corporation

© Trade Mark of Union Carbide Corp. for active ingredient  
 Carbaryl (1-Naphthyl N-methylcarbamate).  
 U.S. Pat. No. 2,903,478

SEVIN® #2 AQUA, a new and different insecticide. This product mixes instantly in water and is ideal for use through home garden sprayers. It will not clog spray nozzles. SEVIN® #2 AQUA controls most insects that damage Turf, Vegetables, Strawberries, Ornamental Trees and Shrubs including several insects that have developed resistance to other pesticides, such as Colorado Potato Beetle, Flea Beetle, Mexican Bean Beetle, Japanese Beetle and Leatherhopper. SEVIN® #2 AQUA can be used on many Fruit and Vegetable Crops close to harvest.

SEVIN® #2 AQUA can be mixed with most of the Wettable Powder insecticides and fungicides. It can also be mixed with many other liquid insecticides, however, before using with other liquid materials, make a small batch and test to make sure they are physically compatible. SEVIN® #2 AQUA does not control mites; use with a suitable miticide containing Kelthane. This product is unstable when mixed with lime, Bordeaux, Lime and Sulphur, Casein emulsifiers or other alkaline materials. Some injury may occur when used on tender foliage when prolonged wet periods or during periods of high humidity. Do not use on Boston Ivy.

For protection of honey bees, avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application. Avoid spray drift to adjoining food and forage crops or traps in bloom.

SHAKE WELL BEFORE USING. CLOSE CONTAINER IMMEDIATELY AFTER USE TO PREVENT MATERIAL FROM EVAPORATION OR DRYING. PROTECT FROM TEMPERATURES BELOW 32° F. Pour recommended amount of this material into nearly filled spray tank, or pre-mix in a bucket. Agitate during filling and spraying.  
**ORNAMENTAL PLANTS, FLOWERS, SHRUBS AND SHADE TREES:** Rose, Chrysanthemum, Arbor Vitae, Azalea, Juniper, Elm, Maple, Oak, Dogwood and Pines. To control Japanese Beetle, Rose Sawfly, Periodical Cicada, Mealy Bugs, Leaf

RECEIVED  
JAN 2, 1969  
72-500

CONTAINS 2 LBS. SEVIN  
INSECTICIDE PER GALLON

NET CONTENTS  
6 FLUID OUNCES



VEGETABLES

ACTIVE INGREDIENT  
Carbaryl (1-Naphthyl Methylcarbamate) 21.50%  
INERT INGREDIENTS 78.50%

Do not store below 32° F

NOTE FOR PHYSICIAN  
Avoid contamination of foodstuffs

Manufactured By  
**MILLER CHEMICAL & FERTILIZER CORPORATION**  
Hanover, Pennsylvania 17331, U.S.A. • Subsidiary of Alco Standard Corporation

SMALL MEASUREMENTS

® Trade Mark of Union Carbide Corp. for active ingredient  
Carbaryl (1-Naphthyl N-methylcarbamate)  
U.S. Pat. No. 2,903,478

ORNAMENTAL PLANTS, FLOWERS, SHRUBS AND SHADE  
TREES