

This Turf Food is a balanced plant food designed to provide the grass plants with necessary nutrients to aid in the development of a luxurious turf.

The Organic Nitrogen, Ureaform, is a Long Lasting Nitrogen combined with Quick Release Nitrogen. The result is immediate feeding plus continued even feeding.

The other plant foods in this product will help the plants to withstand drought and winter injury by aiding the development of an adequate root system.

TURF FOOD PLUS 2 will kill the following broadleaved weeds:

Ragweed
Buckhorn

Carpet Weed
Dandelion

Plantain
Wild Garlic
Wild Onion

And certain other broadleaved weeds sensitive to 2,4-D.

READ ENTIRE LABEL BEFORE USING

FOR USE ON ESTAB.

Apply contents of this bag to 5,000 sq. ft. Make applications any time weeds are actively growing. For best results, application of this product should be made in late spring (May).

	RATE NO. 1 50 Lbs./5,000 Sq. Ft.	RATE NO. 2 25 Lbs./5,000 Sq. Ft.
CROSS COUNTRY	7- ¹ / ₄	.6
SCOTT 35-8	6	.4 ¹ / ₂
JACKSON 24-26	7- ¹ / ₄	.6
JACKSON 20CP-26	7- ¹ / ₄	.6
GARDEN MARK	10	.8 ¹ / ₂
LAWN BEAUTY	7- ¹ / ₄	.5 ¹ / ₂
CYCLONE	4	.3 ¹ / ₄

When using RATE NO. 1, go over lawn once. To insure more even coverage, use RATE NO. 2 and go over lawn twice - North to South and then East to West.

72-509

Overlap spreader wheel marks so as not to leave untreated areas. Shut off spreader when stopping or turning around.

NOTE: Due to variabilities in spreader equipment and accuracy in machine adjustment, the application rates may vary slightly. No warranty is made as to the uniformity of coverage obtained from suggested spreader settings listed.

Do Not apply when grass is wet or on hot, humid days.

For Spot Treatments: Use 1 standard level cupful to treat 60 sq. ft.

46

PRECAUTIONS

Rain, sprinkling, mowing or heavy traffic on treated areas within 24 hours will remove material from grass, reducing effectiveness.

New Lawns may be treated with TURF FOOD PLUS 2 after 4 mowings.

Do Not reseed within 6 weeks after applying TURF FOOD PLUS 2.

Bent grasses, White Clover and other legumes are damaged by 2,4-D.

Tall grass should be mowed to a height of approximately 2 inches. Remove clippings if they are heavy. Best results will be obtained if lawns are not mowed for several days before or after using this material, and weeds have started to make new growth.

Re-treatment is usually necessary in 3 to 4 weeks to control resistant weeds such as Knotweed or Doorweed (*Polygonum aviculare*), Wild Garlic (*Allium vineale*), Wild Onion (*Allium canadense*), Red or Sheep Sorrel (*Rumex acetosella*) and certain 2,4-D semi-tolerant weeds. "For most effective control of Wild Garlic and Wild Onion, application should be made in the spring (April) and in the fall (October) for two or more consecutive growing seasons."

2,4-D sensitive weeds will show distortion, or twisting, within a few days, however, it may take 2 to 4 weeks to obtain complete kill of 2,4-D sensitive weeds. Action is slower on mature plants during cool weather or on plants growing slowly due to dry weather. MILLER'S TURF FOOD PLUS 2 will not control established grassy weeds, such as Crabgrass, Nutgrass, Quackgrass, Bermudagrass that may be a problem.

CROSS COUNTRY	7-1/4	6
SCOTT 35-8	6	4 1/2
JACKSON 24-26	7-1/4	6
JACKSON 20CP-26	7 1/4	6
GARDEN MARK	10	8 1/2
LAWN BEAUTY	7-1/4	5 1/2
CYCLONE	4	3 1/4

When using RATE NO. 1, go over lawn once. To insure more even coverage, use RATE NO. 2 and go over lawn twice - North to South and then East to West.

72-509

Overlap spreader wheel marks so as not to leave untreated areas. Shut off spreader when stopping or turning around.

NOTE: Due to variabilities in spreader equipment and accuracy in machine adjustment, the application rates may vary slightly. No warranty is made as to the uniformity of coverage obtained from suggested spreader settings listed.

Do Not apply when grass is wet or on hot, humid days.

For Spot Treatments: Use 1 standard level cupful to treat 60 sq. ft.

46

BE CAREFUL

Rain, sprinkling, mowing or heavy traffic on treated areas within 24 hours will remove material from grass, reducing effectiveness.

New Lawns may be treated with TURF FOOD PLUS 2 after 4 mowings.

Do Not reseed within 6 weeks after applying TURF FOOD PLUS 2.

Bent grasses, White Clover and other legumes are damaged by 2,4-D.

Tall grass should be mowed to a height of approximately 2 inches. Remove clippings if they are heavy. Best results will be obtained if lawns are not mowed for several days before or after using this material, and weeds have started to make new growth.

Re-treatment is usually necessary in 3 to 4 weeks to control resistant weeds such as Knotweed or Doorweed (Polygonum aviculare), Wild Garlic (Allium Vineale), Wild Onion (Allium canadense), Red or Sheep Sorrel (Rumexacetosella) and certain 2,4-D semi-tolerant weeds. "For most effective control of Wild Garlic and Wild Onion, application should be made in the spring (April) and in the fall (October) for two or more consecutive growing seasons."

2,4-D sensitive weeds will show distortion, or twisting, within a few days; however, it may take 2 to 4 weeks to obtain complete kill of 2,4-D sensitive weeds. Action is slower on mature plants during cool weather or on plants growing slowly due to dry weather. MILLER'S TURF FOOD PLUS 2 will not control established grassy weeds, such as Crabgrass, Nutgrass, Quackgrass, Bermudagrass that may be a problem.

Excessive applications may be harmful. Avoid drift onto, or use on desirable plants, vegetables, flowers, trees and shrubs. Do Not use on lawns, golf greens, or tees containing Bent grass, St. Augustine grass, Carpet grass and Dichondra. Clover may be affected by this product and may be killed with repeated application. To control Clover, use MILLER'S IMPROVED LIQUID WEEDAWAY. Do not store near seed, insecticides, fungicides or fertilizers. Clean spreader thoroughly after using. Avoid contact with skin, eye or clothing. Keep bag tightly closed when not in use and destroy empty bag. "Store in a safe, dry place away from children and domestic animals."

The use of this material being beyond our control and involving elements of risk to human beings, animals and vegetation, we do not make any warranty, expressed or implied, as to the effects of such use, when this product is not used in accordance with the directions as stated on this label.

72-509

Manufactured by
MILLER CHEMICAL & FERTILIZER CORPORATION
 Hanover, Pennsylvania 17331/Subsidiary of Alco Standard Corporation

47