J

TRI-FENE EXTRA TM

Net Contents U.S.D.A. Registration No	_
A selective combination herbicide for control of seedling Johnsongrass and annual broadleaf weeds and grasses.	
ACTIVE INGREDIENTS:	
2,4-Dichlorophenoxyacetic acid, dimethylamine salt* 12.5% 2,3,6-Trichlorophenylacetic acid, dimethylamine salt* 18.5%	
INERT INGREDIENTS: 69.0%	
*2,4-Dichlorophenoxyacetic acid equivalent 10.4% by weight	

TRI-FENE is a registered trademark of Tenneco Chemicals, Inc.



**2,3,6-Trichlorophenylacetic acid equivalent ... 15.6% by weight

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANEL FOR OTHER CAUTIONS

or 12 lbs. per gallon.

t ...

TENNECO CHEMICALS, INC. 280 Park Avenue New York, N. Y. 10017 A Major Component of Tenneco, Inc.

ال درية والمالية

TRI-FENE EXTRA TM herbicide is a dual purpose herbicide for killing weeds in sugarcane, both preemergence and early post emergence.

Weeds controlled by TRI-FENE EXTRA herbicide include the following:

GRASS WEEDS

BROADLEAF WEEDS

Alexander grass	chickweed
barnyard grass	dock
crabgrass	henbit
foxtail	morningglo
goosegrass	nightshade
seedling johnsongrass	purslame

ragweed
spiny amaranth
t sow thistle
ngglory and other weeds
shade susceptible to 2,4-D

Consult your Extension Service Weed Control Specialist for current recommendations on rates, timing, and combinations with other weed killer products labeled for use on sugarcane.

DIRECTIONS

Sugarcane

LOUISIANA

Fall Application

Treat after planting but before seedling weeds are more than 2 inches tall.

Apply on a band 30" wide on 72" rows using 12-18 gallons of water per banded acre.

A. August 1 - September 15

Weed Problem

AMOUNT TRI-FENE EXTRA PER BANDED ACRE

Light Textured Soils Dark, Heavy Textured Soils

1. Seedling johnsongrass 3 quarts

2-3 quarts

2. Annual broadleaves and grasses 2-3 quarts 2-3 quarts

B. From September 15 until planting is completed.Apply 2 to 3 quarts per banded acre regardless of soil texture.

Spring Application

Apply in the spring after shaving, off-barring and redrawing the rows of Fall planted cane, and after all danger of freezing is past. Make applications to rows free of weeds and grasses.

- A. Where TRI-FENE EXTRA herbicide was used in the Fall, repeat treatments in the Spring on the same area using 3 to 4 quarts on light textured soils where seedling johnsongrass is a problem. For all other situations, use the same rate as was applied in the Fall.
- B. When TRI-FENE EXTRA herbicide was not applied in the Fall, use 3 to 4 quarts in 12 to 18 gallons of water per acre, applied on a band 30 inches wide. Use the higher rate on light textured soils or where johnsongrass seedlings are a major problem.

FLORIDA

Apply 1 gallon TRI-FESE EXTRA herbicide in 50 gallons water per acre as an over-all

Treat Full planted cane shortly after planting, but before weeds are 4 inches high. Depending on the response of the weeds, a second application using 1 gallon TRI-FENE EXTRA per broadcast agre may be required 3 weeks after the first treatment.



CAUTION

- ---- Avoid contact with skin, eyes, or clothing.
- ---- Harmful if swallowed.
- ---- KEEP OUT OF REACH OF CHILDREN.
- ---- Keep out of any body of water.
- ---- Apply, TRI-FENE EXTRA herbicide as specified on this label.
- --- Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes. Do not burn containers.
- --- Do not use bagasse for feed or forage.
- ---- Avoid spray drifts. TRI-FENE EXTRA herbicide may injure susceptible desirable broad leaf plants such as tobacco, tomatoes, soybeans, cotton, most vegetable crops, ornamentals, etc.

 Coarse sprays are less likely to drift.
- ---- Do not use same sprayer for other purposes.
- ---- Do not contaminate water used for irrigation or domestic purposes.
- --- Do not store near feed, fertilizers, seeds, insecticides, or fungicides.

NOTICE

TENNECO CHEMICALS, INC. MAKES NO WARRANTY BEYOND THE DESCRIPTION OF THE PRODUCT HEREON AND DIRECTIONS FOR USE UNDER NORMAL CONDITIONS OF USE, NOR ANY WARRANTY OF MERCHANTABILITY AND DISCLAIMS ANY LIABILITY FOR CONSEQUENTIAL OR INCIDENTAL DAMAGES ARISING OUT OF THE SALE OR USE OF THIS MATERIAL. BUYER'S SOLE REMEDY SHALL BE REFUND OR REPLACEMENT.

T	D	T	

Net	Contents	

A selective combination herbicid and annual broadleaf weeds and g

ACTIVE INGREDIENTS:

2,4-Dichlorophenoxyacetic ac 2,3,6-Trichlorophenylacetic

INERT INGREDIENTS:

*2,4-Dichlorophenoxyacetic ac or 1 1b. per gallon.

**2,3,6-Trichlorophenylacetic or 1½ lbs. per gallon.

TRI-FENE is a registered



CAUTION: KEEP OUT OF REACH OF C

TRI-FENE EXTRA TM herbicide is a dual purpose herbicide for killing weeds in sugarcane, both preemergence and early post emergence.

Weeds controlled by TRI-FENE EXTRA herbicide include the following:

GRASS WEEDS

BROADLEAF WEEDS

Alexander grass	chickweed	ragueed
barnyard grass	dock	spiny amaranth
crabgrass	h <u>enbi</u> t	sow thistle
foxtail	morningglory	and other weeds
goosegrass	nightshade	susceptible to 2,4-D
seedling johnsongrass	purslane	•

Consult your Extension Service Weed Control Specialist for current recommendations on rates, timing, and combinations with other weed killer products labeled for use on sugarcane.

DIRECTIONS

Sugarcane

LOUISIANA

Fall Application

Treat after planting but before seedling weeds are more than 2 inches tall.

Apply on a band 30" wide on 72" rows using 12-18 gallons of water per banded acre.

- A. August 1 September 15

 Weed Problem

 Light Textured Scils Dark, Heavy Textured Soils
 - 1. Seedling johnsongrass 3 quarts 2-3 quarts
 - 2. Annual broadleaves and grasses 2-3 quarts 2-3 quarts
- B. From September 15 until planting is completed.
 Apply 2 to 3 quarts per banded acre regardless of soil texture.

Spring Application

Apply in the spring after shaving, off-barring and redrawing the rows of Fall planted cane, and after all danger of freezing is past. Make applications to rows free of weeds and grasses.

- A. Where TRI-FENE EXTRA herbicide was used in the Fall, repeat treatments in the Spring on the same area using 3 to 4 quarts on light textured soils where seedling johnsongrass is a problem. For all other situations, use the same rate as was applied in the Fall.
- B. 'TRI-FENE EXTRA herbicide was not applied in the Fall, use 3 to 4 quarts in 12 18 gallons of water per acre, applied on a band 30 inches wide. Use the higher rate on light textured soils or where johnsongrass seedlings are a major problem.

FLORID.

Apply 1 gailon TRI-FENE EXTRA herbicide in 50 gallons water per acre as an over-all spray.

Treat Fall planted cane shortly after planting, but before weeds are 4 inches high. Depending on the response of the weeds, a second application using 1 gallon TRI-FFS LYDA per broadcast acre may be required 3 weeks after the first treatment.

CAUTION

- ---- Avoid contact with skin, eyes, or clothing.
- ---- Harmful if swallowed.
- ---- KEEP OUT OF REACH OF CHILDREN.
- ---- Keep out of any body of water.
- ---- Apply TRI-FENE EXTRA herbicide as specified on this label.
- ---- Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes. Do not burn containers.
- --- Do not use bagasse for feed or forage.
- ---- Avoid spray drifts. TRI-FENE EXTRA herbicide may injure susceptible desirable broad leaf plants such as tobacco, tomatoes, soybeans, cotton, most vegetable crops, ornamentals, etc.

 Coarse sprays are less likely to drift.
- ---- Do not use same sprayer for other purposes.
- ---- Do not contaminate water used for irrigation or domestic purposes.
- --- Do not store near feed, fertilizers, seeds, insecticides, or fungicides.

NOTICE

TENNECO CHEMICALS, INC. MAKES NO WARRANTY BEYOND THE DESCRIPTION OF THE PRODUCT HEREON AND DIRECTIONS FOR USE UNDER NORMAL CONDITIONS OF USE, NOR ANY WARRANTY OF MERCHANTABILITY AND DISCLAIMS ANY LIABILITY FOR CONSEQUENTIAL OR INCIDENTAL DAMAGES ARISING OUT OF THE SALE OR USE OF THIS MATERIAL. BUYER'S SOLE REMEDY SHALL BE REFUND OR REPLACEMENT.