

**NET WEIGHT 40 POUNDS
WATKINS SWAT®
MEDICATED**

EPA Reg. No. 99-85

EPA Est. No. 99-Mn-1

Aids in preventing the breeding of horn flies and face flies in the manure of treated cattle, when used as directed. Aids in the control of stomachworms (*Haemonchus*, *Ostertagia*, and *Trichostrongylus* spp.), nodular worms (*Oesophagostomum* spp.), large-mouth bowelworms (*Chabertia* spp.), and hookworms (*Bunostomum* spp.) in cattle when fed in accordance with directions for use.

Pesticidally Active Drug Ingredient, Permethrin	95.6%
Pesticidally Inert* Ingredients	4.4%
Total	100.0%

*Refers only to ingredients which are neither insecticides nor drugs.

GUARANTEED ANALYSIS

MINERALS (Percent Guarantee)

Calcium (Ca), not more than	18.9000%	Iodine (I), not less than	0.0075%
Calcium (Ca), not less than	15.8000%	Copper (Cu), not less than	0.0075%
Phosphorus (P), not less than	9.5000%	Manganese (Mn), not less than	0.0075%
Magnesium (Mg), not less than	4.0000%	Zinc (Zn), not less than	0.0075%
Iron (Fe), not less than	1.7000%	Cobalt (Co), not less than	0.0062%
Sulfur (S), not less than	0.9000%		

VITAMINS (Minimum Guarantee Per Pound)

Vitamin A	125,000 U.S.P. Units	Riboflavin	75 mg.
Vitamin D ₃	90,000 I. C. Units	d-Pantothenic Acid	30 mg.
Niacin	120 mg.		

INGREDIENTS

Dicalcium Phosphate, Calcium Carbonate, Dried Cane Molasses, Magnesium Oxide, Iron Sulfate, Iron Oxide, Mineral Oil, Sulfur, Vitamin A Acetate in Gelatin (Stability Improved), Vitamin D₃ Supplement (Stability Improved), Riboflavin Supplement, Niacin, Calcium Pantothenate, Cobalt Carbonate, Potassium Iodide (Stability Improved), Manganous Oxide, Copper Oxide, Zinc Oxide.
See feeding directions on back of tag.

WARNING: Keep out of reach of children!
See back panel for additional warnings.

WATKINS PRODUCTS, INC., WINONA, MINN. 55987


WARNING: Do not administer to lactating dairy animals. Occasional sensitivity to phenothiazine has been reported in cattle. Consult a veterinarian before using in severely debilitated animals. Certain parasites may become resistant to phenothiazine therapy. This treatment will not affect population of existing adult flies.

FEEDING DIRECTIONS

of salt per day. However, because of salt content of water and natural feedstuffs, salt consumption will vary. Two ounces of Swat are needed for a 1,000 lb. animal. In using this product it is necessary to determine the amount of salt consumed daily per 1,000 lbs. of herd weight. If salt consumption is not known, this should be determined during a 1 month period prior to using Swat.

To aid in the selection of the proper mix use the following guide:

For Each 1,000 lbs. of Herd Weight	
Daily consumption of salt in ounces	Pounds of Salt to add to 40 lbs. of Swat
1/4	5 lbs.
1/2	10 lbs.
3/4	15 lbs.
1	20 lbs.
1 1/4	25 lbs.
1 1/2	30 lbs.
1 3/4	35 lbs.
2	40 lbs.



Mix the Swat and salt thoroughly. Feed no other mineral or salt.

Self-feed in protected feeders located near watering or other gathering places. For best results, the self-feeding of Watkins Swat should begin one month before the face fly and horn fly season starts and be fed continuously for internal parasite control. To control existing flies, cattle should be sprayed at start of fly season with residual insecticide. Because face flies may migrate from neighboring herds, an area wide control program will assist in the reduction of the adult face fly population.

STORE IN A DRY PLACE

WATKINS PRODUCTS, INC., WINONA, MINN. 55987

PRINTED IN U. S. A.