NET WEIGHT 24 OZ. (1.5 LBS.)

Crickets-Dust infested area in ornamental plant beds (6 oz. per 500 sq. ft.), preferably after a rain or watering. Sowbugs, Millipedes—In the garden, apply to soil around plants at the time of planting or transplanting (beans, peas, tomatoes). Use 1 lb. per 250-300 square feet. White Grubs, Sod Webworms, Bill Bugs, Wireworms, Japanese Beetle Grubs, (radish, onion, cabbage Root Maggots)-Apply dust to the seed bed soil at the time of planting or transplanting for radishes, onions and cabbage (11b. to 600-700 square feet) and work into top 2 or 3 inches of soil. No additional treatments are needed during the season. Pests in Lawns—For Ants and Chiggers dust area,  $\frac{1}{2}$  lb. to 250-300 sq. feet. For Japanese Beetle Grubs, White Grubs, Sod Webworms, use 1 lb, per 250-300 sq, ft, and soak area with hose; keep children or pets away from treated area until the area is dry. Fests on Flowers and Ornamentals-(Grasshoppers, Thrips, Plant Burs)—Dust plants thoroughly, both sides of foliage. CAUTION: /This material, if taken internally, can cause toxic symptoms. Avoid inhalation of dust and avoid contamination of foodsturfs or cooking utensils. In case of skin contact wash with soap and water immediately. This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed.

Keep out of any body of water. Apply this product only as

**Active Ingredient:** Technical Chlordane\*. Inert Ingredients

\*Equivalent to 3.0% Octachloro-4, 7-Meth and 2.0% of related compounds.

## U.S.D.A. Registration No.



specified on this label.

. · ·



## 

	USES AND DIRECTIONS: Except where indicated, repeat
	treatments when necessary to keep pests under control. This
	shaker-type can can be used in many situations; for treating
	large areas, a hand dust gun is needed.
	<b>INDOORS:</b> Dust only areas inaccessible to children and pets.
	Roaches (Cockroaches, "Waterbugs") Apply as a spot treat-
	ment to cracks, crevices and other hiding places or runways.
	AntsApply to runways. Dusting the outside walls and founda-
	tion may help keep ants from entering the nome. Givers
100.0%	Apply as spot treatment to corners, baseboards, under cabinets,
	in closets, etc. Silverfish, Firebrats—Apply to infested areas,
thanotetrahydroindane	corners in closets, behind radiators, around pipes. Spiders,
, , , , , , , , , , , , , , , , , , ,	Centipedes, Scorpions-Dust infested area along baseboards,
	door sills, behind shelving, damp basement corners, storage
	areas, and around pipes where pests are noticed or may hide.
No. 99-86	Brown Dog Ticks—Treat sleeping quarters of pets and replace
	brown bog ricks real sleeping quarters of pers and replace
	bedding. Do not treat the animals. Apply as a spot treatment
	to cracks, crevices, and hiding places.
	OUTDOORS (Lawn, Garden, Ornamental Plants): EarwigsDust
	infested area around flower gardens as soon after young earwigs
	come out of the soil in the spring, again in midsummer if neces-
AVINONIA MANI 55007	sary. Use 6 ounces of dust for 500 square feet. Thorough
WINONA, MN. 55987	treatment may make indoor treatments unnecessary. Mole
	PRINTED IN U.S.A