

ACCEPTED

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the products registered under EPA Reg. No. 99-114

99-114

WEIGHT 25 POUNDS

SWAT FOR DAIRY CATTLE

(Contains RABON Oral Larvicide)

To prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle.

ACTIVE INGREDIENT	BY WEIGHT
2-chloro-1-(2, 4, 5-trichlorophenyl) vinyl dimethyl phosphate	.70%*
INERT INGREDIENTS	99.30%**
TOTAL	100.00%

*Each pound contains 3175 mg. RABON
**Refers only to ingredients which are not larvicidal.

GUARANTEED ANALYSIS

Crude Protein, Not less than	14.00%
Crude Fat, Not less than	2.50%
Crude Fiber, Not more than	10.00%

FEED INGREDIENTS

Wheat Middlings, Iron Oxide.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See reverse side of tag for use directions and other precautions.

EPA Reg. No. 99-114
EPA Est. No. 99-MN-1
U. S. Patent No. 3,102,842
Made in U. S. A.

Manufactured by
WATKINS INCORPORATED
WINONA, MN 55987

FD7759

Printed in U. S. A.

BEST DOCUMENT AVAILABLE

66

ACCEPTED
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, and the authority
thereunder
EPA Reg. No. 99-114

99-114

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION
KEEP OUT OF REACH OF CHILDREN**

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning equipment or disposal of wastes.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Feed Swat for Dairy Cattle at the rate of 1/4 pound per animal per day to animals weighing between 400-1,100 pounds by incorporating it into the daily ration, or top dress on the grain or roughage. Cattle weighing between 1,100-1,800 pounds should be fed 6 ounces per day.

Start feeding early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.

Swat for Dairy Cattle can be fed to dry cows, replacement animals and lactating cows. Swat for Dairy Cattle prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but it is not effective against existing adult flies.

In some cases supplemental fly control measures may be needed in and around dry lots, calf pens and barns to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, Swat for Dairy Cattle should be used in conjunction with other good sanitation and management practices.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of the container in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

FD7759

Printed in U. S. A.

BEST DOCUMENT AVAILABLE

4-24