

NET WEIGHT 40 POUNDS

WATKINS SWAT FOR BEEF CATTLE

(Contains RABON Oral Larvicide)

To prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle.

ACTIVE INGREDIENT	BY WEIGHT
2-chloro-1-(2, 4, 5-trichlorophenyl) vinyl dimethyl phosphate	0.800%*
INERT INGREDIENTS	99.200%**
TOTAL	100.000%

*Each pound contains 3628 mg RABON.

**Refers only to ingredients which are not larvicidal.

MINERALS (Percent Guarantee)

Calcium (Ca), not more than	12.520%
Calcium (Ca), not less than	10.440%
Salt (NaCl), not more than	10.800%
Salt (NaCl), not less than	9.000%
Phosphorus (P), not less than	5.000%
Fluorine (F), not more than	0.050%

VITAMINS (Minimum Guarantee Per Pound)	
Vitamin A	250,000 IU
Vitamin D	100,000 IU

FEED INGREDIENTS

Dicalcium Phosphate-Monocalcium Phosphate, Wheat Midds, Ground Corn, Calcium Carbonate, Salt, Magnesium Oxide, Sulfur, Mineral Oil, Iron Oxide, Iron Sulfate, Potassium Chloride, Zinc Oxide, Vitamin A Acetate in Gelatin (Stability Improved), Vitamin D₃ Supplement (Stability Improved), Copper Oxide, Vitamin E Supplement, Vitamin B₁₂ Supplement, Riboflavin, Niacinamide, Cobalt Carbonate, Manganous Oxide, Potassium Iodide (Stability Improved), d-Calcium Pantothenate.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See reverse side of tag for use directions and other precautions.

EPA Reg. No. 99-118
EPA Est. No. 99-MN-1
U. S. Patent No. 3,102,842
Made in U. S. A.

Manufactured by
WATKINS INCORPORATED
WINONA, MN 55987

7760

Printed in U. S. A.

BEST DOCUMENT AVAILABLE

ACCEPTED

Under the Federal Insecticide, Fungicide, and Rodenticide Act
LFA Reg. No. 99-118

99-118

ACCEPTED

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS**

**CAUTION
KEEP OUT OF REACH OF CHILDREN.**

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust.
Wash thoroughly with soap and water after handling and before eating or smoking.
If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

**DIRECTIONS FOR USE
GENERAL CLASSIFICATION**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Feed Watkins Swat for Beef Cattle at the rate of 2 ounces per animal per day to animals weighing up to 500 pounds, by incorporating it into the daily ration, or top dress on the grain or roughage. Cattle weighing 500 to 1,200 pounds should be fed 4 ounces per day. Cattle weighing 1,200 to 1,900 pounds should be fed at least 6 ounces per day. Start feeding early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.

Swat for Beef Cattle prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but it is not effective against existing adult flies.

In some cases supplemental fly control measures may be needed in and around dry lots, calf pens and barns to control adult house flies and stable flies which can breed not only in cattle manure, but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, Swat for Beef Cattle should be used in conjunction with other good sanitation and management practices.

No withdrawal time required before slaughter.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of the container in an incinerator or landfill approved for pesticide containers, or bury in a safe place. Pesticide that cannot be used should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

7760

Printed in U. S. A.

BEST DOCUMENT AVAILABLE