

100 5 2019-118

NET WEIGHT 40 POUNDS

WATKINS SWAT FOR PASTURE CATTLE

(Contains RABON Oral Larvicide)

A mineral-vitamin supplement containing RABON for control of fecal flies in manure of treated cattle. Prevents development of face flies, horn flies, house flies, and stable flies in the manure of treated cattle.

ACTIVE INGREDIENT	BY WEIGHT
2-chloro-1-(2, 4, 5-trichlorophenyl) vinyl dimethyl phosphate	0.848%*
INERT INGREDIENTS	99.152%**
TOTAL	100.000%

*Each pound contains 3849 mg RABON.

**Refers only to ingredients which are not larvicidal.

MINERALS (Percent)

Calcium (Ca), not more than	14.16%
Calcium (Ca), not less than	11.80%
Salt (NaCl), not more than	10.80%
Salt (NaCl), not less than	9.00%
Phosphorus (P), not less than	5.00%
Sulfur (S)	2.64%
Potassium (K)	1.60%
Magnesium (Mg)	1.50%
Iron (Fe)	0.55%
Manganese (Mn)	0.44%
Zinc (Zn)	0.44%
Copper (Cu)	0.11%
Iodine (I)	0.0055%
Cobalt (Co)	0.0030%
Selenium (Se)	0.0022%

VITAMINS (Per Pound)

Vitamin A	350,000 IU
Vitamin D ₃	145,000 IU
Vitamin E	100 IU
Niacin	53 mg
Riboflavin	32 mg
d-Pantothenic Acid	16 mg
Vitamin B ₁₂	500 mcg

FEED INGREDIENTS

Dicalcium Phosphate-Monocalcium Phosphate, Wheat Midds, Calcium Carbonate, Salt, Potassium Sulfate, Magnesium Sulfate, Sodium Selenite, Sodium Bentotenite, Mineral Oil, Ferrous Carbonate, Magnesium Oxide, Sulfur, Manganous Oxide, Zinc Oxide, Potassium Chloride, Copper Oxide, Vitamin A Acetate in Gelatin (Stability Improved), Vitamin D₃ Supplement (Stability Improved), Vitamin E Supplement, Vitamin B₁₂ Supplement, Riboflavin Supplement, Niacin Supplement, Calcium Iodate, Cobalt Carbonate, d-Calcium Pantothenate.



KEEP OUT OF REACH OF CHILDREN
CAUTION
 See reverse side of tag for use directions
 and other precautions.

EPA Reg. No. 99-118
 EPA Est. No. 99-MN-1
 U.S. Patent No. 3,102,842
 Made in U.S.A.

Manufactured by
WATKINS INCORPORATED
WINONA, MN 55987

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS**

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Feed Watkins Swat for Pasture Cattle by incorporating it into the daily ration, or top dress on the grain or roughage so that the intake per day is 0.07 gm of Rabon for each 100 pounds of body weight. This amounts to 0.018 pound of Watkins Swat for Pasture Cattle for each 100 pounds of body weight.

Some guidelines are as follows:

Average body Weight (pounds)	Pounds of Swat per head per day
200	0.036
500	0.09
800	0.146
1100	0.200
1400	0.255

In some cases supplemental fly control measures may be needed in and around dry lots, calf pens, and barns to control adult house flies and stable flies which can breed not only in cattle manure, but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, Swat for Pasture Cattle should be used in conjunction with other good sanitation and management practices.

No withdrawal time required before slaughter.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE — Store in a dry place not accessible to children or pets.

PESTICIDE DISPOSAL — Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL — Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

BEST AVAILABLE COPY