

FEED FOR CATTLE AND SWINE
Contains RABON Oral Larvicide

To prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies
in the manure of treated cattle and House Flies in the manure from treated Swine.

| ACTIVE INGREDIENT | WEIGHT |
|---|----------|
| 2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate | 6.65* |
| Inert Ingredients | 93.35*** |
| Total | 100.00 |

*Each pound contains 29.2 grams RABON. (R)
***Refers only to ingredients which are not larvicidal

GUARANTEED ANALYSIS

| | |
|------------------------------------|--------|
| Crude Protein, not less than | 13.00% |
| Crude Fat, not less than | 2.50% |
| Crude Fiber, not more than | 9.00% |

FEED INGREDIENTS

Processed Grain by-Products

KEEP OUT OF REACH OF CHILDREN

CAUTION

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a dry place not accessible to children or pets.

PESTICIDE DISPOSAL: Pesticide, concentrate or rinseate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Dispose of bags according to approved Federal, State or local procedures under the Resource Conservation and Recovery Act.

See reverse side of tag for use directions and other precautions.

RABON (R) is a Registered Trademark of Shell Oil Company
EPA Reg. No. 29-101
EPA Act. No. 29-51-1
U.S. Patent No. 3,100,112
Made in U.S.A.

ACCEPTED
with COMMENTS
in EPA Label Dec 8

DEC 28 1982

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, this product is registered under EPA Reg. No.

99-121

Manufactured by
MATHIAS CORPORATION
WINONA, MN 55987

NET CONTENTS 25 POUNDS

ENVIRONMENTAL STATEMENT
HATKING 2000 WAX
CAUTION
KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARD

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning equipment or disposal of wastes.

DIRECTIONS FOR USE
GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MIXING DIRECTIONS FOR CATTLE
(Per Ton Concentrate Mix)

| Pounds of concentrate consumed per animal per day | RANGE | | Amount of HATKING 2000 WAX FOR CATTLE and SWINE Premix per ton of concentrate | |
|---|--------|--------|---|-----|
| | lb/ton | oz/ton | lbs. | oz. |
| 0.5 | 188 | 0.25 | 105 | 13 |
| 1.0 | 94 | 0.12 | 52 | 6 |
| 1.5 | 62 | 0.08 | 35 | 4 |
| 2.0 | 47 | 0.06 | 26 | 3 |
| 3.0 | 31 | 0.04 | 17 | 2 |
| 4.0 | 23 | 0.03 | 12 | 1 |
| 5.0 | 19 | 0.02 | 9 | 1 |
| 10.0 | 9 | 0.01 | 4 | 0.5 |
| 15.0 | 6 | 0.007 | 3 | 0.4 |
| 20.0 | 4 | 0.005 | 2 | 0.3 |
| 25.0 | 3 | 0.004 | 1.5 | 0.2 |
| 30.0 | 2 | 0.003 | 1 | 0.1 |

MIXING DIRECTIONS FOR SWINE
(See Appropriate Measuring Device)

Feed HATKING 2000 WAX for cattle and swine at the rate of 1 to 1.5 ounces per animal per day to animals weighing between 400-500 pounds by incorporating it into the daily ration, or top dress on hay or roughage. Cattle weighing between 100-200 pounds should be fed 0.5 to 1 ounce per animal per day. Cattle 200-400 pounds should be fed 1 to 1.5 ounces per animal per day.

Start the fly control early in the spring before flies begin to appear and continue feeding HATKING 2000 WAX until the fall until cold weather restricts fly activity.

Cattle containing HATKING 2000 WAX PREMIX may be fed up to slaughter and to lactating dairy cows without withholding milk from market during or after treatment.

HATKING 2000 WAX for cattle and swine can be fed to dry cows, replacement animals and lactating cows.

HATKING 2000 WAX for cattle and swine prevents the development of horn flies, face flies, house flies and stable flies in the pasture of treated cattle, but it is not effective against existing horn flies.

In some cases, supplemental fly control measures may be needed in and around dry lots, calf pens and barns to control adult house flies and stable flies which can breed not only in cattle manure, but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, Swat for Cattle and Swine should be used in conjunction with other good sanitation and management practices.

DIRECTIONS FOR SWINE

No preslaughter withholding period is required.

Pigs: (Weaners to market weight) Mix 1.5 lbs of Swat for Cattle and Swine per ton of meal type feed and offer free choice. This is equivalent of 22.4 mg of Pabon per pound of feed.

Cows, boars, and breeding gilts: Mix 3 lbs of Swat for Cattle and Swine per ton of meal type feed and offer 4.6 lbs of feed per animal per day. This is equivalent of 44.8 mg of Pabon per pound of feed.

Start feeding Swat for Cattle and Swine early in the spring before flies begin to appear and continue feeding throughout the summer and the fall until cold weather restricts fly activity.

All Swine should be treated. Swat for Cattle and Swine prevent development of house flies in the manure from treated swine, but is not effective against existing adult flies.

In some cases, supplemental fly control measures may be needed in and around the area to control adult flies which come in from other areas or which breed on decaying vegetable matter elsewhere on the premises.

In order to achieve optimum fly control, Swat for Cattle and Swine should be used in conjunction with other good management and sanitation practices.

300/79444
17/2

310/79491
18/3

DRS
12/27/82

DEC 28 1982

Mr. Lloyd Ferdinandsen
Watkins Incorporated
100 Liberty Street
Winona, MN 55937

Dear Mr. Ferdinandsen:

Subject: Amendments - Add Swine and Change Product
Name
Watkins Swat for Dairy Cattle
EPA Registration No. 99-121
Your Application Dated September 9, 1982

The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable, provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
 - a. The percentage of Rabon declared should show as 6.00%.
 - b. Please refer to the enclosed type size guide for assistance in selecting the correct print size for the front panel signal word, child hazard warning, and the heading "Storage and Disposal."
 - c. Delete "for 15 minutes" from the "If in eyes" advice.
 - d. ~~Delete "GENERAL CLASSIFICATION" since the added used on swine has not been classified.~~ *filed 12/27/82*
 - e. Correct the spelling of "adult" in the paragraph beginning "In some cases," in cattle feeding directions.
 - f. Revise the heading "DIRECTIONS FOR SWINE" to "FEEDING DIRECTIONS FOR SWINE."

The registration file has been amended to show the change in product name to "Swat for Cattle and Swine."

3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(c). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

George T. Larocca
Product Manager (15)
Insecticide-Scienticide Branch
Registration Division (TS-767C)

Enclosure

RB:GTLAROCCA:DCR-39860:WANG-1709C:CBP:Raven:479-2013:12/20/82