

Princep 4G
brand of simazine

Granular Herbicide

For weed control in
certain crops, orna-
mental plantings, on
industrial sites, and
around-the-farm

_____ Pounds
Net Weight

Active Ingredient:	
Simazine (2-chloro-4, 6-bis-(ethylamino)-s- triazine)	4%
Inert Ingredients:	96%
Total	100%

Princep 4G is a 4%
granular herbicide


Caution: Keep out of reach of
children. Harmful if swallowed.
Avoid inhalation of dust, and
contamination of food and feed.

Do not contaminate domestic or
irrigation water supplies or
lakes, streams or ponds.

Do not reuse container. Destroy
when empty.

DIRECTIONS FOR USE AND CONDI-
TIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire
Directions For Use and the
Conditions Of Sale And Warranty
before using this product.



4.

Caneberries, i.e. blackberries, boysenberries, loganberries, raspberries

Broadcast a single application of 50-100 lbs. per acre in the spring before bud break and weed emergence.

To control quackgrass, broadcast 100 lbs. per acre in the fall or split the application, broadcasting 50 lbs. per acre in the fall and 50 lbs. per acre in the early spring, when quackgrass is actively growing.

Note: 1. On new plantings (less than 6 months old), use one-half the above rates. 2. Do not apply when fruit is present.

Cranberries

In Massachusetts: Broadcast up to 100 lbs. per acre prior to weed emergence. Apply in the spring prior to new cranberry growth or in the fall after harvest.

In other areas: Broadcast 50 lbs. per acre in the spring prior to weed emergence.

Grapefruit and Oranges

In Florida: Broadcast 160-240 lbs. per acre in the spring (March or April) prior to weed emergence. Repeat annually.

In Texas: Broadcast 100-125 lbs. per acre in the spring prior to weed emergence. Repeat annually.

Precautions: 1) Do not use in citrus nurseries. 2) Do not apply to bedded citrus. 3) Use only in groves where trees have been established for one year or more. 4) Do not apply to trees under stress from freeze damage for one year following the freeze to avoid possible injury. 5) Avoid contact with fruit.

Nonselective Weed Control on Noncrop Land

Apply Princep at rates recommended below. Use the higher rates on heavy clay and muck soils. Best results are obtained when Princep 4C is applied before weeds emerge.