

Sutan+®: Use in a preplant incorporated tank mixture for control of all weeds claimed on both the Princep and Sutan+ labels including partial control (suppression) of shattercane (wild cane). Fluid fertilizer may replace all or part of the water in the spray. Check the physical compatibility of mixture with fertilizer before use. Refer to the General Information section of this label for a compatibility test procedure. Use a minimum of 20 gals. of spray volume per acre. Refer to the Sutan+ label for incorporation directions. Use the higher rate of Princep on fine-textured soil and where heavy broadleaf infestations are expected. For partial control (suppression) of shattercane, broadcast and incorporate immediately before planting 1.25-2.5 lbs. of Princep 80W plus 7.33 pts. of Sutan+ 6.7E per acre. For control of the other weeds claimed on both labels, broadcast and incorporate 1.25-3.75 lbs. of Princep 80W plus 4.75 pts. of Sutan+ 6.7E per acre. Observe all precautions and limitations on the Princep 80W and Sutan+ labels.

✓ Sugarcane - *See label (rather use)*

Apply 2.5-5 lbs. per acre at planting or after harvest if ratoon crop is to be grown, but before weeds or sugarcane emerge. Princep 80W may also be applied interline-directed or over the sugarcane to provide continuing control of germinating weeds.

Precautions: 1) Do not apply more than one 2.5-5 lb. application over the plants or more than two-interline directed applications. 2) Do not apply after "close-in." 3) Total application should not exceed 12.5 lbs. per acre crop cycle.

Note: Do not feed or graze treated forage to livestock.

Grasses Grown for Seed (Pacific Northwest Only)

Perennial ryegrass, Highland or Astoria bentgrasses, orchard-grass, Wall fescue, and fine fescues such as Illahee, Rainier, Pennlawn, Chewings, and related species. For the control of broadleaf weeds and annual grasses including annual ryegrass, rattail fescue, silver hairgrass, and downy brome (cheat), apply 2.5 lbs. Princep 80W in a minimum of 15 gals. of water per acre as soon as fall rains start. Early application is important.

Precaution: Apply only to grasses from which at least one seed crop has been cut.

Note: Allow 30 days between last application and grazing of dairy and beef cattle and sheep, and allow 60 days between the last application and cutting for hay.

Nursery Stock
✓ Nurseries, ✓ Christmas Tree Plantings, ✓ Shelterbelts

Nurseries (see list below)

Apply 2.5-3.75 lbs. in at least 25 gals. of water per acre in fall or spring at least one year after transplanting.

✓ Christmas Tree Plantings and Shelterbelts (see list below)

Remove weed growth before application. Apply 2.5-5 lbs. in at least 25 gals. of water per acre after transplanting. Use the same rate for annual maintenance applications.

For quackgrass control apply 5 lbs. per acre in the fall or apply a split application of 2.5 lbs. per acre in the fall plus 2.5 lbs. per acre in early spring, after quackgrass begins growth.

Precautions: 1) Do not use Princep 80W on seedbeds or cutting beds. 2) In CA, OR, and WA, do not apply to Christmas trees or shelterbelts sooner than one year after transplanting. In other areas, do not apply to Christmas trees or shelterbelt transplants less than three years of age. 3) Do not apply more than once a year, except as directed for quackgrass control.

Apply Princep 80W to these species of trees and shrubs, as recommended above:

Conifers

- ✓ Arborvitae (Sh, S) (Pl, S) (N.S.)
- ✓ Austrian pine (N.S.), (Sh, S), (Pl, S)
- ✓ Bishop pine
- ✓ Knobcone pine (N.S.)
- ✓ Lodgepole pine (shore pine)
- ✓ Monterey pine (Sh, S) (Pl, S) (N.S.)
- ✓ Mugho pine (N)
- (N.S.) (Sh, S) (N) ← ✓ Red pine (Norway pine)
- ✓ Scotch pine (N) (Sh, S) (Pl, S)
- ✓ White pine (Sh, S) (Pl, S)
- ✓ Blue spruce (Sh, S) (Pl, S)
- ✓ Norway spruce (Sh, S) (Pl, S) (N)
- ✓ Red spruce (N) (Pl, S) (Sh, S)
- ✓ White spruce (N)
- ✓ Balsam fir
- ✓ Douglas fir
- ✓ Fraser fir (Sh, S) (N) (Pl, S)
- ✓ White fir
- ✓ Hemlock (Sh, S) (N) (Pl, S)
- ✓ Juniper (Sh, S), (N), (Pl, S)
- ✓ Red cedar
- ✓ White cedar (Pl, S) (N) (Sh, S)
- ✓ Yew (Taxus spp.) (Sh, S) (N)

✓ Deciduous Trees and Woody Ornamentals

- ✓ American elm (N)
- ✓ Siberian elm (Sh, S) (N)
- ✓ Barberry (N)
- ✓ Boxelder (Sh, S) (N)
- ✓ Bush honeysuckle (N) (Sh, S)
- ✓ Caragana (Sh, S)
- ✓ Cotoneaster (Sh, S) (N)
- ✓ Dogwood (N)
- ✓ Eucalyptus (N)
- ✓ Holly (Ilex spp.) (N) (Sh, S) (max. 4 lbs./A)
- ✓ Honey locust (Sh, S) (N)
- ✓ Oregon grape (Mahonia spp.) (N)
- ✓ Pieris spp. (max. 4 lbs./A) (N) (Sh, S)
- ✓ Red oak (Sh, S) (N)
- ✓ Russian olive (Sh, S) (N) (Pl, S)
- ✓ Oleander (N) (Sh, S)
- ✓ Palm (2.5 lbs./A) (N)
- ✓ Bottle brush (N) (Sh, S)
- ✓ Carob (N.S.) (N)

*For California only

Surflan Tank Mix: On Christmas tree plantings, use this tank mix for preemergence control of weeds listed on the Surflan 75W (or Surflan A.S.) and Princep 80W labels. Use on field grown conifer species listed on the labels for each herbicide plus grand fir, alpine fir, Engelmann spruce, black spruce, Colorado blue spruce, Coulter pine, giant redwood, and Veitchi fir. Broadcast the mixture as a directed spray to the soil surface or as an overtop spray using 2.5-5 lbs. of Princep 80W and 2.67-5.33 lbs. of Surflan 75W (or 2-4 qts. of Surflan A.S.). Apply in sufficient water per acre to uniformly treat the area. Follow overtop sprays with sprinkler irrigation to move the herbicide from leaf surfaces to the soil. Remove weed growth before application. Mix weed residues, prunings or trash into the soil, or remove them before treatment. Soil should be in good tilth and free of clods at time of application. Shallow cultivation (1-2 inches) after treatment will not reduce weed control. Observe all precautions and limitations on the Princep 80W and Surflan labels.

Dual® 8E Tank Mix: On field and liner grown woody ornamentals, use this tank mix for preemergence control of yellow nutsedge, barnyardgrass, crowfootgrass, fall panicum, giant foxtail, green foxtail, yellow foxtail, goosegrass, large crabgrass, smooth crabgrass, redroot pigweed, galinsoga, Venice mallow, lambs-quarters, purslane, ragweed, smartweed and other weeds listed on the Dual 8E alone label. Control should last about 60 days. Apply 1-1.25 lbs. of Princep 80W plus 2-3 pts. of Dual 8E in a minimum of 15 gals. of water per acre. For band applications use a proportional amount of herbicide. Direct the spray toward the base of established ornamentals or ornamentals transplanted at least 10 days. Apply before weeds emerge or after removing existing weeds. Use the higher rates on fine-textured soil and where broadleaf weeds are expected to form a significant part of the infestation. Use the lower rates on coarse-textured soil and where light infestations of broadleaf weeds are expected. Apply posttransplant to the following field and liner grown ornamentals:

Arborvitae (<u>Thuja</u> spp.)	Leucothoe (<u>Leucothoe</u> spp.)
Cotoneaster (<u>Cotoneaster</u> spp.)	Osmanthus (<u>Osmanthus</u> spp.)
Crabapple (<u>Malus</u> spp.)	Weigela (<u>Weigela</u> spp.)
Holly (<u>Ilex</u> spp.), Boxleaf	Yew (<u>Taxus</u> spp.), Dwarf
Japanese, Japanese	Japanese
Juniper (<u>Juniperus</u> spp.), Andorra,	
Creeping, Shore	

Note: Length of weed control may be reduced when continuous wet soil conditions follow herbicide application.

Precautions: Do not use on seedbeds or on unrooted cuttings. Do not use in greenhouses or other enclosed areas.

✓ Tree Plantations for Timber

Apply before weeds emerge or following removal of weed growth. Apply before or after seedling transplanting. If applying after transplanting, application may be made over the top of tree seedlings before new growth appears. If new growth is present, direct the spray to avoid tree foliage contact. Do not use on seed nurseries or injury may occur. Apply only once per year.

The following rates are based on broadcast treatment in 20-100 gals. of water per acre. For band or spot applications, reduce the broadcast rate in proportion to the area actually sprayed.

✓ Black Walnuts (Plant, Soil)

Apply 2.5-5 lbs. per acre.

✓ Slash Pine and White Pine

Use for establishment in the South only. Apply 5-7.5 lbs. per acre.

✓ Loblolly Pine

Use for establishment in the South only. Apply either 5-7.5 lbs. of Princep 80W per acre or a tank mix of 4-6 lbs. of Princep 80W plus 1-2 lbs. of AAtrex 80W (or 0.9-1.75 lbs. AAtrex Nine-O or 0.8-1.6 qts. AAtrex 4L or 4LC) per acre. The tank mix provides broad spectrum annual weed control (see list in General Information section of this label) and better weed control in limited rainfall areas than Princep 80W alone.

✓ Tree Seedlings in North Central U.S. *Princep 80W*

Apply 2.5-5 lbs. per acre on the following species of trees:

American beech	red pine (Norway pine)
balsam fir	red spruce
→ black locust	Russian olive
caranga	Scotch pine
dogwood	*sweet gum
*Eastern red cedar	*white ash
honey locust	white cedar
maple	*white oak
Norway spruce	white pine
→ pecan	white spruce
*red oak	

*Instead of Princep 80W alone, on these species a tank mix of 2.5 lbs. of Princep 80W + 2.5 lbs. of AAtrex 80W (or 2.2 lbs. AAtrex Nine-0 or 2 qts. AAtrex 4L or 4LC) per acre may be applied in a minimum of 25 gals. of water.

Precautions: Apply only once per year. On sandy soil, do not use more than 2.5 lbs. Princep 80W per acre and do not use the tank mix. Do not apply the tank mix directly to seedlings after the start of bud break.

Turfgrasses for Sod

St. Augustinegrass, Centipedegrass, and Zoysia Grass

Apply 2.5-5 lbs. per acre, according to soil texture as indicated below.

Muck or peat	5 lbs.	Old beds	within 2 days after lifting of sod
		New beds	3-4 days after sprigging or plugging
Sandy soil	2.5 lbs.	Old beds	within 2 days after lifting of sod
		New beds	7-10 days after sprigging or plugging

If weeds regrow, apply an additional 2.5 lbs. on muck or peat, or 1.25 lbs. on sandy soil.

Precautions: 1) Do not apply within 30 days before cutting or lifting. 2) Do not use north of NC.

Turfgrass for Fairways, Lawns, and Similar Areas

Bermudagrass, Centipedegrass, St. Augustinegrass, and Zoysia Grass

Apply Princep 80W after October 1 before emergence of winter annual weeds for control of annual bluegrass, burclover, carpet burweed, chickweed, corn speedwell, henbit, hop clover, and spurweed. Princep 80W will control annual bluegrass even if it is emerged at time of treatment. For control of summer annual weeds listed in the General Information section of this label, also apply Princep 80W in late winter before the weeds emerge. Apply in a minimum of 15 gals. of water per acre or 1 gal. per 1,000 sq. ft.

Where annual bluegrass is the major weed, use 1.25 lbs. of Princep 80W per acre (0.5 oz. per 1,000 sq. ft.). Use 2.5 lbs. per acre (1 oz. per 1,000 sq. ft.) for control of the other weeds named above. Do not exceed 1.25 lbs. per acre per treatment on newly sprigged turfgrass or on hybrid bermudagrass such as Tiflawn, Tifway, and Ormond.

For continued summer annual weed control, apply another 1.25 lbs. per acre at least 30 days after the previous application, but not after April 15. However, do not make more than two applications of this product per year.

Precautions: To avoid injury, use only on turfgrass reasonably free of infestations of insects, nematodes, and diseases. Do not use on golf greens. Do not use north of NC or west of the high rainfall areas of eastern OK and eastern TX. Do not use on muck or alkaline soils. Do not apply over the rooting area of trees or ornamentals not listed on this label. On newly sprigged turfgrass, temporary slowing of growth may follow application. On hybrid bermudagrass, temporary yellowing may occur. Do not overseed with desirable turfgrass within 4 months before or 6 months after treatment.

Vegetable Crops

✓ Asparagus (Established)

Apply 2.5-5 lbs. per acre to beds established one year or more. Apply once per year either after disking in the spring before weeds emerge, or following harvest after removal of existing weed growth. Spring application must be made at least 3 days before first cutting.

Precautions (WA and OR only): 1) Do not apply on gravelly, sand, or loamy sand soil or crop injury may result. 2) Do not allow irrigation water to stand or to form pools in the field or injury may occur.

✓ Artichokes

Apply 2.5-5 lbs. per acre as a directed spray after the last fall tillage.

Precautions: 1) Do not apply on gravelly, sand, or loamy sand soil. 2) Apply only once per year.

Nonselective Weed Control on Noncrop Land

Best results are obtained when Princep 80W is applied before weeds emerge. If weed growth has begun and removal is not feasible, include a contact or translocated herbicide in the spray.

Princep 80W provides long-term weed control on industrial sites, highway medians and shoulders, railroad rights-of-way, lumberyards, petroleum tank farms, and in ~~noncrop areas~~ on farms, such as around buildings, equipment and fuel storage areas, along fences, roadsides, and lanes. Do not use this product for nonselective weed control on land to be cropped, near adjacent desirable trees, shrubs, or plants, or in greenhouses, as injury may occur. Do not use in areas accessible to livestock or allow livestock to graze treated weed foliage.

Aerial applications may be made where it is feasible. Use at least 1 gal. of water for each 1 lb. of Princep 80W for both ground and aerial applications. Use more water if practical.

To control annual broadleaf and grass weeds (including barnyard-grass, cheat, crabgrass, common lambsquarters, foxtail, ragweed, puncturevine, and turkey mullein) apply 6-12.5 lbs. per acre.

To control most annual and many perennial broadleaf and grass weeds (including quackgrass, bluegrass, redtop, burdock, Canada thistle, orchardgrass, purpletop, smooth brome, dogfennel, and plantain) apply 12.5-25 lbs. per acre.

To control hard-to-kill perennial weeds (including bull thistle and sowthistle) apply 25-50 lbs. per acre.

For longer residual control in regions of high rainfall and a long growing season, apply 25-50 lbs. per acre.

Tank Mixture with Roundup

This tank mixture will provide control of emerged annual weeds and residual control of weeds listed on the Princep label. The combination also will partially control emerged perennial weeds listed on the Roundup label. Add to the spray tank the appropriate rate of Princep noted in the above section for Princep alone. Then add 1-5 qts. of Roundup per acre depending on weeds present and their growth stage. Also add an agriculturally approved nonionic surfactant at 0.5% by volume of spray solution. Apply the mixture in a 10-40 gals. of water per acre as a postemergence spray at the appropriate weed growth stage given on the Roundup label. Provide constant agitation during mixing and application to keep the mixture in suspension. Refer to the Application Procedures section of this label for further directions. Refer to the labels of both herbicides for specific noncrop sites, rates, weeds controlled, and further directions, precautions, and limitations.

Tank Mixture with Oust®

Use this tank mixture to control the weeds listed below with either preemergence, early postemergence, or postemergence applications. Use the higher rates within the ranges given under conditions of heavy weed growth, on soil with over 2 1/2% organic matter, on areas with heavy decaying plant residues, or on high moisture areas such as along road edges or railroad shoulders. To improve wetting and/or contact activity on emerged weeds, add a nonionic surfactant such as DuPont Surfactant WK or X-77 at 1 qt. per 100 gals. of spray. Do not apply by aircraft. Provide constant agitation during mixing and application to keep the mixture in suspension. Refer to the labels of both herbicides for specific noncrop sites and further directions, precautions, and limitations.

Preemergence to Early Postemergence: To control the weeds listed below apply 5-10 lbs. of Princep 80W plus 3-5 ounces of Oust in a minimum of 25 gals. of water per acre.

Broadleaf Weeds (Up to 3-4" in Height or Diameter)

bouncingbet	ox-eye daisy
buckhorn plantain	pigweed
burclover	prickly lettuce
Carolina geranium	puncturevine
common chickweed	purple starthistle
common dandelion	ragweed
common speedwell	Russian thistle
common yarrow	sowthistle
crimson clover	St. Johnswort
dogfennel	sunflower
filaree	sweet clover
hoary cress	tansymustard
horsetail	tansy ragwort
kochia	tumble mustard
lambquarters	turkey mullein
little mallow	vetch
mustard	wild carrot
	yellow rocket

Grasses (Up to 6-12" in Height)

Alta fescue	Italian ryegrass
annual bluegrass	johnsongrass (short-term control)
annual ryegrass	little barley
bahia grass	red brome
barnyardgrass	red fescue
cheatgrass	reed canarygrass
crabgrass	ripgut brome
downy brome	ryegrass
fall panicum	smooth brome
fescue	sprangletop
foxtails	witchgrass
Indian grass	

Postemergence: To control actively growing weeds listed below apply 10-20 lbs. of Princep 80W plus 6-12 ounces of Oust in a minimum of 25 gals. of water per acre.

Actively Growing Weeds

bedstraw	Jerusalem artichoke	johnsongrass
Canada thistle	kudzu	(season-long control)
curly dock	mayweed	yellow nutsedge
dewberry	musk thistle	
fiddleneck	poison ivy	
fleabane	puncturevine	
goldenrod	Spanishneedles	
hemlock	turkey mullein	
hemp dogbane	Virginia pepperweed	
honeysuckle	wild blackberry	

Storage and Disposal

Store in a dry place. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited. Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable federal, state, or local procedures. Completely empty bag by shaking, and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by state and local authorities.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Causes eye irritation. Harmful if inhaled or absorbed through the skin. Avoid breathing dust or spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

First Aid: If in eyes, flush with plenty of water. Get medical attention if irritation persists. If on skin, wash with soap and water. If inhaled, remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth; get medical attention.

Environmental Hazards

Do not contaminate domestic, livestock or irrigation water supplies, or lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

AAtrex® trademark of CIBA-GEIGY for atrazine

Compex® trademark of Kalo Laboratories, Inc.

Dual® trademark of CIBA-GEIGY for metolachlor

Eradicane® trademark of Stauffer Chemical Co.

Evik® trademark of CIBA-GEIGY for ametryn

Princep® trademark of CIBA-GEIGY for simazine

Nine-O® trademark of CIBA-GEIGY

U.S. ENVIRONMENTAL PROTECTION AGENCY
 OFFICE OF PESTICIDES PROGRAMS
 REGISTRATION DIVISION (PH-567)
 WASHINGTON, D.C. 20460

EPA REGISTRATION NO.	DATE OF ISSUANCE
TERM OF ISSUANCE	
NAME OF PESTICIDE PRODUCT	

NOTICE OF PESTICIDE: REGISTRATION
 REREГИSTRATION
 (Under the Federal Insecticide, Fungicide
 and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

<input type="checkbox"/>	<input type="checkbox"/>	161 37	102397 1
<input type="checkbox"/>	<input type="checkbox"/>	161 37	102332 1
<input type="checkbox"/>	<input type="checkbox"/>	161 15	102333 1

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

ATTACHMENT IS APPLICABLE


SIGNATURE OF APPROVING OFFICIAL	DATE
---------------------------------	------

Permanent Label
(Booklet)

Princep® 80W

Herbicide

For weed control in certain
crops, ornamental plantings,
on industrial sites, and
around-the-farm



Active Ingredient:

Simazine: 2-chloro-4,6-bis(ethylamino)-s-triazine	80%
Inert Ingredients:	20%
Total:	100%

EPA Reg. No. 100-437

Princep 80W is a
wetable powder

Keep Out of Reach of Children.

Caution

See additional precautionary statements
at end of booklet.

See directions for use inside booklet.

CIBA-GEIGY



DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of CIBA-GEIGY or the Seller. All such risks shall be assumed by the Buyer.

CIBA-GEIGY warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. CIBA-GEIGY makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall CIBA-GEIGY or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. CIBA-GEIGY and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of CIBA-GEIGY.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR WEED CONTROL, AND/OR ILLEGAL RESIDUES.

General Information

Apply Princep 80W Herbicide before weeds emerge or after removal of weed growth. Princep controls a wide variety of annual broadleaf and grass weeds when used at selective rates in agricultural crops and ornamental plantings. When used at higher, nonselective rates in noncrop areas, it also controls many perennial broadleaf and grass weeds.

Where a range of application rates is given, use the low rate on coarser textured soil and soil lower in organic matter; use the high rate on finer textured soil and soil higher in organic matter.

Since Princep enters weeds mainly through their roots, moisture is needed to move it into the root zone. Very dry soil conditions and lack of rainfall following application may necessitate shallow cultivation.

Princep 80W is noncorrosive to equipment, nonflammable, and has low electrical conductivity.

Annual Weeds Controlled

alyssum	common lambsquarters
annual bluegrass	nightshade
annual morningglory	pepperweed
annual ryegrass	pigweed
barnyardgrass (watergrass)	pineappleweed
burclover	prickly lettuce
carelessweed	common purslane
carpetweed	ragweed
common chickweed	rattail fescue
crabgrass (<i>Digitaria</i> spp.)	redmaids
downy brome (cheatgrass)	Russian thistle
fall panicum	shepherdspurse
fiddleneck	signalgrass
filaree	(<i>Brachiaria</i> spp.)
fireweed	silver hairgrass
fivehook bassia	smartweed
Flora's paintbrush	spanishneedles
Florida pusley	speedwell
foxtails	tansymustard
goosegrass	wild mustard
groundsel	wild oats
henbit	witchgrass
junclerice	yellow flower pepperweed
knawel (German moss)	

Following many years of continuous use of this product and chemically related products, biotypes of some of the weeds listed on this label have been reported which cannot be effectively controlled by this and related herbicides. Where this is known or suspected and weeds controlled by this product are expected to be present along with resistant biotypes, we recommend the use of this product in registered combinations or in sequence with other registered herbicides which are not triazines. If only resistant biotypes are expected to be present, use a registered non-triazine herbicide. Consult with your State Agricultural Extension Service for specific recommendations.

Application Procedures

Ground application: Use conventional ground sprayers equipped with nozzles that provide accurate and uniform application. Be certain that nozzles are uniformly spaced and the same size. Calibrate sprayer before use and recalibrate at the start of each season and when changing carriers. Unless otherwise specified, use a minimum of 20 gals. of spray mixture per acre.

Use a pump with capacity to (1) maintain 35-40 psi at nozzles, (2) provide sufficient agitation in tank to keep mixture in suspension, and (3) to provide a minimum of 20% bypass at all times. Use centrifugal pumps which provide propeller shear action for dispersing and mixing this product. The pump should provide a minimum of 10 gals./minute/100 gal. tank size circulated through a correctly positioned sparger tube or jets.

Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on the suction side of the pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles. Check nozzle manufacturer's recommendations.

For band applications, calculate amount to be applied per acre as follows:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \frac{\text{broadcast rate}}{\text{per acre}} = \frac{\text{amount needed}}{\text{per acre of field}}$$

Aerial application: Use aerial application only where specified in the use directions. Apply in a minimum of 1 gal. of water for each 1-1.5 lbs. of wettable powder applied per acre. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Avoid application to humans or animals. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin, and should wash thoroughly before eating and at the end of each day's operation.

Mixing procedures - all uses: (1) Be sure sprayer is clean and not contaminated with any other materials, or crop injury or sprayer clogging may result. (2) Fill tank 1/4 full with clean water, nitrogen solution, or complete fluid fertilizer. (3) Start agitation. (4) Be certain that the agitation system is working properly and creates a rippling or rolling action on the liquid surface. (5) Pour product directly from bag into tank. (6) Continue filling tank until 90% full. Increase agitation if necessary to maintain surface action. (7) Add tank mix herbicide(s) after this product is thoroughly suspended. (8) Finish filling tank. Maintain agitation to avoid separation of materials. (9) Empty tank as completely as possible before refilling to prevent buildup of emulsifiable concentrate residue from possible tank mix herbicides. (10) If an emulsifiable concentrate film starts to build up in tank, drain it and clean with strong detergent solution or solvent. (11) Clean sprayer thoroughly immediately after use by flushing system with water containing a detergent.

before addition, or (B) add 1/2 of the compatibility agent to the fertilizer and the other 1/2 to the emulsifiable concentrate or flowable herbicide before addition to the mixture. If still incompatible, do not use the materials mixed in the same spray tank.

Rotational Crops and Perennial Crop Replanting

To avoid crop injury, observe the following precautions: 1) If rotating treated land the year following application, plant only corn, unless otherwise stated in this label. 2) If replanting perennial crops or if rotating land to crops other than corn, do not apply Princep in the year preceding planting these crops.

Fruit and Nut Crops

Apply the spray to the orchard or vineyard floor avoiding contact with fruit, foliage, or stems. Apply only to orchards or groves where trees have been established one year or more unless specified differently. Make only one application per year, except as noted otherwise. Recommended rates are based on broadcast treatment. For band applications or spot applications around trees in fruit or nut plantings, reduce the broadcast rate of Princep 80W and water per acre in proportion to the area actually sprayed.

Almonds and Peaches (CA only)

Apply 1.25-2.5 lbs. of Princep 80W per treated acre in a 2-4 ft. band on each side of the tree row. Apply before weeds emerge in late fall or early winter. Weeds controlled by 1.25 lbs. of Princep 80W include burclover, common chickweed, wild mustard, and shepherdspurse. Apply only once per year.

Precautions: 1) Do not treat trees established in the grove less than 3 years. 2) Do not treat the Mission (Texas) variety of almonds. 3) Do not apply to almond trees propagated on plum rootstocks. 4) Do not replant almonds or peaches in treated soil for 12 months after treatment. 5) Do not apply on gravelly, sand, or loamy sand soil or on soil with less than 1% organic matter. 6) Do not treat areas where water will accumulate.

Apples, Pears, Sour Cherries

Apply 2.5-5 lbs. per acre.

✓ Avocados

CA and FL only: Apply 2.5-5 lbs. per acre after final preparation of grove.

Precaution: Do not apply on gravelly, sand, or loamy sand soil.

✓ Blueberries and Caneberries (blackberries, boysenberries, loganberries, raspberries)

Apply 2.5-5 lbs. per acre in the spring or apply a split application of 2.5 lbs. per acre in the spring plus 2.5 lbs. per acre in the fall. Apply in a minimum of 40 gals. of water per acre.

On plantings less than 6 months old, use 1/2 the above rate.

To control quackgrass, apply 5 lbs. per acre in the fall or split the application applying 2.5 lbs. per acre in the fall plus 2.5 lbs. per acre in the spring, when quackgrass is actively growing.

Note: Do not apply when fruit is present.

✓ Cranberries (see *comest. handbook*)

MA: Apply up to 5 lbs. per acre either before spring growth begins or in the fall after harvest.

Other areas: Apply 2.5 lbs. per acre before spring growth begins.

✓ Filberts

Apply 2.5-5 lbs. per acre in the fall or apply a split application of 2.5 lbs. per acre in the fall plus 2.5 lbs. per acre in the spring.

Precautions: 1) If trees are planted on a hillside, excessive soil erosion may result from the elimination of weeds.
2) Do not apply on gravelly, sand, or loamy sand soil.

Note: Do not apply when nuts are on the ground during the harvest period.

✓ Grapes (not recommended)

Apply 2.5-6 lbs. per acre any time between harvest and early spring.

Precaution: Do not use in vineyards established less than three years. Do not apply on gravelly, sand, or loamy sand soil.

✓ ✓ ✓
Grapefruit, Lemons, Oranges

AZ (Lemons and Oranges only): Apply a split application of 2 lbs. per acre in the spring plus 2 lbs. per acre in the fall.

CA: In grapefruit, lemons, and oranges, apply 2.5-5 lbs. per acre in a single application; or apply 2.5 lbs. per acre in the fall and 2.5 lbs. per acre in the spring. Do not use in the Imperial, Coachella, or Palo Verde valleys.

FL (Grapefruit and Oranges only): Apply only once per year. For control of weeds listed under General Information, apply 8-12 lbs. per acre in the spring; to control milkweed vine, use 10-12 lbs. per acre either preemergence or before vines are 12 inches tall.

TX (Grapefruit and Oranges only): Apply 5-6 lbs. per acre.

Precautions (All areas): 1) Do not use in nurseries. 2) Do not apply to bedded grapefruit, lemons, or oranges. 3) To avoid possible injury, do not apply to trees under stress from freeze damage for one year after the freeze.

✓ Macadamia Nuts

Apply 2.5-5 lbs. in 50 gals. of water per acre before harvest and just prior to weed emergence. Repeat application as necessary. Do not apply when nuts are on the ground during the harvest period.

✓ Olives

Apply 2.5-5 lbs. per acre following grove preparation in the fall. ✓ Repeat annually in midwinter.

✓ Peaches, Plums, Sweet Cherries

Apply 2-5 lbs. per acre. Apply in late fall to early spring prior to weed emergence.

Precautions: 1) Do not apply on gravelly, sand, or loamy sand soil. 2) Peaches: use only in AR, LA, MO, OK, TX, and states east of the Miss. River. For California, see specific directions in the section "Almonds and Peaches (CA only)." 3) Plums and sweet cherries: use only in MO and states east of the Miss. River except TN.

✓ Pecans

Apply the appropriate rate in the following table before weeds emerge in the spring.

Soil texture	Rate per acre
Sand, loamy sand	Do not use
Sandy loam	2.5 lbs.
Loam or clay soil low in organic matter	2.5-3.5 lbs.
Clay soil high in organic matter	3.5-5 lbs.

Precautions: Do not use west of the Pecos River in TX or in NM, AZ, or CA, or injury will occur. Do not make applications to transplanted trees that have been established less than two years in the grove or injury may occur.

Note: Do not apply when nuts are on the ground. Do not allow animals to graze treated areas.

✓ Strawberries

Oregon and Washington: For control of chickweed, groundsel, mustard, and shepherdspurse, apply broadcast 1.25 lbs. per acre. In fields where overhead irrigation is used to activate this product, apply after harvest at time of bed renovation. In fields where overhead irrigation is not available, apply during early October through November.

Precautions: Make only one application per growing season. Do not apply within 4 months after transplanting. Limit overhead irrigation to 1/2 inch immediately following post-harvest application. Do not use on sand.

✓ Walnuts

Apply 2.5-5 lbs. per acre.

Precautions: 1) Do not apply on gravelly, sand, or loamy sand soil. 2) Leveling and furrowing operations after application will lessen effectiveness of weed control.

Note: Do not apply when nuts are on the ground.

Princep 80W Plus Evik® 80W for Grapefruit and Oranges

Use in grapefruit and oranges in FL and TX for control of weeds listed in the following directions and under General Information. Apply the rates given below in a minimum of 40 gals. of water per acre thoroughly covering existing vegetation, but avoiding contact with citrus fruit, foliage, or stems. Do not spray when windy. Apply when weeds have resumed active growth in the spring, but before they exceed 4 inches in height (except as noted for vines). Apply only once per year. To avoid possible injury, do not use in nurseries, or around trees that have been established in the grove less than two years, or where trees are under stress from freeze damage for one year after the freeze. Do not apply within 30 days before harvest. Do not graze treated areas. Observe all precautions and limitations on the labeling of Princep 80W and Evik 80W.

FL (Ridge Area): Apply 5 lbs. of Princep 80W plus 5 lbs. of Evik 80W per acre; controls milkweed vine when applied either preemergence or on vines up to 24 inches tall.

FL (Bedded Areas): Apply 4 lbs. of Princep 80W plus 6 lbs. of Evik 80W per acre; controls balsamapple when applied either preemergence or on vines up to 12 inches tall.

TX: Apply 4-5 lbs. of Princep 80W plus 2-3 lbs. of Evik 80W per acre plus 2 qts. nonionic surfactant, such as X-77®, per 100 gals. of spray mixture (add surfactant to spray tank last); controls buffelgrass, marinevine (ivy treevine), TX panicum, sandbur, and sunflower, and gives top kill of johnsongrass. Use 5 lbs. of Princep 80W per acre for longer residual control; use 3 lbs. of Evik 80W per acre for control of heavy weed growth and for top kill of johnsongrass; use the higher rate of both products for control of marinevine.

Princep 80W Plus Bromacil 80W for Grapefruit and Oranges
(Florida Only)

Use in grapefruit and oranges in FL for control of balsamapple, black nightshade, carpetweed, crabgrass, cudweed, dayweed, Florida pusley, horseweed, pepperweed, pigweed, poorjoe, ragweed, rattlebox, spanishneedles and sandbur, and for partial control of bermudagrass, bahiagrass, pangolagrass, paragrass, and torpedograss. Apply 4-8 lbs. of Princep 80W plus 3-4 lbs. of bromacil 80W per acre beneath trees in a minimum of 40 gals. of water per acre before or soon after weed growth begins. Avoid contact of foliage and fruit with spray or mist. Apply only once per year. Use the lower rates for light weed infestations or all applications in bedded citrus areas. Use the higher rates for heavy weed infestations only in ridge grown citrus areas. Temporary yellowing of citrus leaves may occur following treatment.

Precautions: To avoid possible injury, do not use in nurseries, or around trees that have been established in the grove less than four years, or where trees are under stress from freeze damage for one year after the freeze. Do not use on soil with less than 1% organic matter or on poorly drained soil. Do not treat trees planted in irrigation furrows. Do not treat diseased trees such as those with foot rot. Do not use in groves interplanted with other trees or desirable plants, nor in home grapefruit or orange plantings, or in areas where roots of other valuable plants or trees may extend. Treated areas may be planted to citrus trees one year after application. Do not rotate to other crops within two years after application.

Princep 80W Plus Paraquat

This tank mix is effective in the following fruit and nut crops for kill of existing vegetation and for residual control of the annual broadleaf and grass weeds claimed for Princep 80W applied alone. This combination is also effective for top kill and suppression of perennial weeds.

Use this tank mix on these crops:

almonds (CA only)	macadamia nuts
apples	olives
avocados (CA & FL ¹ only)	oranges
cherries (sour and sweet ²)	peaches ³
filberts	pears
grapefruit (CA, FL ¹ & TX only)	pecans
grapes	plums ⁴
lemons (AZ & CA only)	walnuts

Apply the Princep 80W rate given under the appropriate crop on this label plus 1-2 qts. paraquat* in 50-200 gals. (30-50 gals. for pecans) of water per acre to the orchard floor avoiding contact with fruit, foliage, or stems. Add a nonionic surfactant, such as X-77, at 0.5 pt. per 100 gals. of spray. Apply when weeds are succulent and new growth is 1-6 inches tall. For mature woody weeds or difficult to control perennial weeds, re-treat or spot treat with paraquat if regrowth occurs. Apply the tank mix only once per year. Do not spray under windy conditions. Use a shield for young trees or vines.

Add Princep 80W to the spray tank first (refer to Mixing Instructions section of this label) then add paraquat, and add the surfactant last. Provide constant agitation during mixing and application to keep the mixture in suspension.

Refer to the Princep 80W and paraquat labels for further directions, specific weeds controlled, and precautions and limitations on each crop.

¹In avocados in Florida, this tank mix also controls balsam-apple vine, rattail amaranth, and at the higher rate of each herbicide, it suppresses coral vine.

²In grapefruit and oranges in Florida, apply 4 lbs. Princep 80W per acre in bedded areas and 8-12 lbs. Princep 80W per acre in ridge areas.

³Limited to CA, AR, LA, MO, OK, TX, and states east of the Miss. River. As appropriate, refer to Princep 80W sections "Almonds and Peaches (CA only)" or "Peaches, Plums, Sweet Cherries" for rate of Princep 80W and other information.

⁴Limited to MO and states east of the Miss. River except TN.

*Based on a product containing 2 lbs. paraquat cation per gal.

Tank Mixture with Roundup*

This tank mixture is effective in grape vineyards and in the following bearing or nonbearing tree crops for control of existing vegetation and for residual control of the annual broadleaf and grass weeds claimed for this product applied alone. This combination is also effective for partial control of perennial weeds contacted by the spray mixture during application. Take extreme care in applying the tank mixture to keep it off fruit, foliage, of stems, or suckers or desirable plants. Observe precautions on both the Princep and Roundup labels for each crop involved. Use this tank mixture on these crops:

Almonds** (CA only)	Macadamia Nuts**
Apples*	Oranges* (AZ,CA,FL ¹ & TX)
Avocados* (CA and FL only)	Peaches* ²
Cherries (sour and sweet ³)	Pears*
Filberts**	Pecans**
Grapefruits* (CA,FL ¹ ,TX)	Plums* ³
Grapes*	Walnuts**
Lemons* (AZ & CA)	

Use the appropriate rate given elsewhere on this label for Princep applied alone in the crop being treated. Add to the spray tank 1-5 qts. of Roundup per acre depending on weeds present and their growth stage. Also add an agriculturally approved nonionic surfactant at 0.5% by volume of spray solution. Apply the mixture in 10-40 gals. of water per acre as a postemergence spray to the weeds at the appropriate weed growth stage given on the Roundup label.

Add Princep to the spray tank first, then add Roundup. Provide constant agitation during mixing and application to keep the mixture in suspension. Refer to Application Procedures section of this label for further directions.

Refer to the labels of both herbicides for further directions, specific weeds controlled, precautions and limitations on each crop.

*Allow a minimum of 14 days between last application and harvest.

**Allow a minimum of 21 days between last application and harvest of these crops.

¹In grapefruit and oranges in Florida, apply 3.5 lbs. of this product per acre in bedded areas, and 7.1-10.6 lbs. per acre in ridge areas.

²Limited to CA, AR, LA, MO, OK, TX, and states east of the Mississippi River. For CA, see specific directions in the section "Almonds and Peaches" (CA only).

³Plums and sweet cherries: limited to MO and states east of the Mississippi River except TN.

Princep 80W Plus Surflan*

Use this tank mix in the following crops for preemergence control of all weeds claimed on both labels:

almonds	lemons
apples	oranges
avocados	peaches
caneberries	pears
cherries	pecans
filberts	plums
grapes	English walnuts
grapefruit	

Apply the Princep 80W rate given under the appropriate crop on this label plus 2.67-5.33 lbs. of Surflan 75W or 2-4 qts. of Surflan A.S. in 20-40 gals. of water per acre. Refer to the Surflan 75W or Surflan A.S. label for complete tank mix directions. Observe all precautions and limitations on the Princep and Surflan labels.

Field and Forage Crops

Alfalfa

Pure alfalfa at least one year old, pure alfalfa less than one year old, and alfalfa-orchardgrass mixtures at least one year old

For ground application, apply Princep 80W in a minimum of 15 gals. of water per acre. Fluid fertilizer may replace all or part of the water used as a carrier, except fluid fertilizers should not be used when alfalfa is actively growing as injury may occur. Check the physical compatibility of Princep 80W with fertilizer before use. Refer to the General Information section of this label for a compatibility test procedure.

For aerial application, apply Princep 80W in a minimum of 2.5 gals. of water per acre.

✓ Pure Alfalfa At Least One Year Old
(All Areas Listed)

Apply once per year in the fall after the last cutting but prior to permanently frozen ground conditions. Alfalfa may be actively growing, semidormant, or dormant when the application is made.

Alfalfa must be growing for at least one year before treatment.

Alfalfa Regions	Weeds controlled	Soil texture	Rate of Princep 80W per acre
WA, OR, NV, UT, ID, MT, WY, CO, and CA counties of Del Norte, Siskiyou, Modoc, Shasta, Lassen, Plumas, Sierra, and Nevada	Downy brome (cheatgrass), tansymustard, prickly lettuce, pepperweed, seedling shepherds- purse, yellow flower pepperweed	Sand and loamy sand Sandy loam Loam soil Clay soil	DO NOT USE 1 lb. 1-1.5 lbs. 1.5-2 lbs.
ND, SD, MN, WI, NE, KS, IA, MI, and NY	Henbit, wild mustard, common chickweed, knawel (German moss), alyssum, downy brome (cheatgrass), wild oats, pepper- weed, tansymustard, pigweed, speedwell, and seedlings of yellow rocket, shepherdspurse, white cockle, fan- weed (pennycress), dandelion, Russian thistle, Kochia	Sand and loamy sand Sandy loam Loam soil Clay soil	DO NOT USE 1 lb. 1.5 lbs. 2 lbs.

(Continued ... 18

IL, IN, OH, KY, TN, NC*. VA, PA, NJ, DE, MD, WV, VT, MA, CT, ME, NH, RI, and MO	Pigweed, common chickweed, mouse- ear chickweed, speedwell, henbit, knawel (German moss), pepperweed, downy brome, alyssum, common lambsquarters, wild oats, and seedlings of shepherdspurse, fanweed (penny- cress), yellow rocket (winter- cress), Russian thistle, dandelion, kochia, white cockle	Sand and loamy sand Sandy loam Loam soil Clay soil	DO NOT USE 1 lb. 1.25 lbs. 1.5 lbs.
---	--	--	--

*use only on loam and clay soils.

Pure Alfalfa Less Than One Year Old (Northeastern U.S. Only)

Pure alfalfa which has been seeded in the spring before June 1 and has become well established may be treated in the fall after the last cutting but before permanently frozen ground conditions. Alfalfa may be actively growing, semidormant, or dormant when Princep is applied. Apply 1 lb. of Princep 80W per acre to control henbit, wild mustard, common lambsquarters, common chickweed, German moss, alyssum, downy brome (cheatgrass), wild oats, tansymustard, pigweed, speedwell, and pepperweed.

Established Alfalfa-Orchardgrass Mixtures (Northeastern U.S. Only)

Apply only to alfalfa-orchardgrass mixtures which have been established at least one year. Make a single application in the fall after the alfalfa-orchardgrass mixture has gone dormant. Applications should be uniformly broadcast at the rate of 1 lb. of Princep 80W per acre to control henbit, wild mustard, common lambsquarters, common chickweed, German moss, alyssum, downy brome (cheatgrass), wild oats, tansymustard, pigweed, speedwell, and pepperweed.

Precautions (For Each Use): 1) Do not use on gravelly, sand, loamy sand, or on soil low in organic matter, or crop injury may occur. 2) Applications made to alfalfa growing on highly calcareous or highly alkaline soil may result in crop injury. 3) Use only in areas where frost occurs and alfalfa goes into a semidormant or dormant state. 4) Do not apply on snow. 5) Princep will not control dodder and established perennial or biennial weeds such as dandelion, white cockle, quackgrass, cinquefoil, wild carrot, and dock. 6) Temporary crop chlorosis may appear on areas where overlapping occurs. The sprayer should be shut off when turning around at the end or corners of the field. 7) Do not rotate to any crops except corn for the growing season following application as injury may occur.

Note: Allow 30 days between application and grazing of dairy and beef cattle and sheep, and allow 60 days between application and cutting for hay.

Princep 80W Plus Paraquat for Alfalfa (WA, OR, NV, UT, ID, MT, WY, CO, and CA counties of Del Norte, Siskiyou, Modoc, Shasta, Lassen, Plumas, Sierra, and Nevada)

Controls filaree, Jim Hill mustard, and shepherdspurse, and gives improved control of bluegrass, cheatgrass, common chickweed, and tansymustard. Apply 1-2 lbs. of Princep 80W plus 1-3 pts. of paraquat* in 20-100 gals. of water per acre with ground equipment or in 5-10 gals. of water per acre aerially. Refer to the Princep 80W rate chart for rates on specific soil textures. Use 1 pt. of paraquat when weeds and grasses are less than 4 inches tall. Apply when weeds are succulent and growth is 1-6 inches tall. Apply to established alfalfa after the last fall cutting when the crop is dormant and before spring growth starts. Do not apply if regrowth following the last fall cutting is more than 2 inches tall. Alfalfa foliage present at application will be burned. Apply once per growing season. Do not pasture animals in treated fields before first cutting. Do not graze, cut, or harvest within 90 days after application. Observe all precautions and limitations on the labeling of Princep 80W and paraquat.

*Based on a product containing 2 lbs. paraquat cation per gal.

✓ Forage Bermudagrasses (Such as Coastal and Midland)

For preemergence control of most annual broadleaf and grassy weeds including barnyardgrass, crabgrass, Brachiaria spp., goosegrass, foxtail, carelessweed, carpetweed, annual morning-glory, pigweed, common purslane, Russian thistle, common chickweed, henbit, and burclover, apply Princep 80W shortly after sprigging. Apply it to established bermudagrass from fall to spring depending on the weed problem, but before weed emergence. Use the appropriate rate in the following table in a minimum of 15 gals. of water per acre for ground application and in a minimum of 1 gal. of water for each 1-1.5 lbs. of wettable powder per acre for aerial application.

Soil texture	Rate per acre
Fine sandy soil low in organic matter	1 lb.
Sandy loam soil	1.5-2.5 lbs.
Loam and clay soil low in organic matter	2.5-3 lbs.
Clay soil	3-4 lbs.

Precautions: 1) Do not use Princep 80W on bermudagrass in NM and the High Plains of TX. 2) Do not apply within 7 days after cutting.

Note: Do not graze dairy or beef cattle or sheep for 30 days after 1-3 lbs. per acre application or for 60 days after 4 lbs. per acre application. Do not cut for hay for 60 days after 1-3 lbs. per acre application or for 90 days after 4 lbs. per acre application.

✓ Corn

Nitrogen solutions or complete fluid fertilizers may replace all or part of the water in the spray. Determine the physical compatibility of Princep 80W with fertilizer before use. Refer to the General Information section of this label for a compatibility test procedure. Do not apply after corn has emerged as there is danger of liquid fertilizers causing crop injury.

Preemergence: Apply before weeds and corn emerge. Use the appropriate rate in the table below.

Preplant: Apply in the spring after plowing at the appropriate rate in the table below. Apply before, during, or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation of Princep 80W. Best results will be obtained when Princep 80W is applied within 2 weeks before planting.

Note: Under dry weather conditions, preplant applications may give better weed control.

Soil texture	Broadcast* rate per acre
Coarse-textured soil: Sand, silt, and loam that is low in organic matter	2.5 lbs.
Medium-textured soil: Soil containing a moderate amount of clay and organic matter	3 lbs.
Fine-textured soil: Loam that is high in organic matter and clay (including dark prairie soils of Corn Belt)	3.75 lbs.
Organic soil: Peat, muck, and high-organic clay	5 lbs.

*For calculation of band treatment rate, see the General Information section.

To control quackgrass: Apply 3.75-5 lbs. per acre in the fall. Plow two to three weeks later, or if erosion is a problem, delay plowing until spring. Do not plant any crop except corn in the spring following treatment. Do not graze treated area.

Note: If weeds develop, particularly under relatively dry conditions, a shallow cultivation will generally result in better weed control.

Precautions: 1) Do not apply more than 5 lbs. of Princep 80W per acre to corn per year. 2) Do not rotate to any crop except corn until the following year or injury may occur. 3) After harvest, plow and thoroughly till the soil in fall or spring to minimize possible injury to spring-seeded rotational crops, regardless of the rate used. 4) If more than 3.75 lbs. of Princep 80W is used per acre (or equivalent rate in a band), a crop of untreated corn should precede the next rotational crop. 5) Do not apply Princep 80W preplant incorporated in corn in the High Plains and Intermountain areas of the West (including central and western KS, western NE, western OK, and the Panhandle of TX) where rainfall is sparse and erratic or where irrigation is required. 6) In the High Plains and Intermountain areas of the West where rainfall is sparse and erratic or where irrigation is required, use Princep 80W to control weeds in corn only when corn is to follow corn or when a crop of untreated corn is to precede another rotational crop. 7) In western MN and eastern parts of the Dakotas, NE, and KS, do not plant soybeans following corn treated with Princep if more than 2.5 lbs. of Princep 80W per acre (or equivalent rate in a band) was applied or injury may occur. 8) Injury may occur to soybeans planted in north-central and northwest IA, south-central and southwest MN, northeast NE, southeast SD, and other areas the year following application on soils having a calcareous surface layer. 9) Do not plant sugar beets, tobacco, vegetables (including dry beans), spring-seeded small grains, or small-seeded legumes and grasses the year after Princep 80W application or injury may occur.

Tank Mixtures on Corn

AAtrex*: Use Princep 80W in a preplant or preemergence tank mixture with AAtrex Nine-O*, AAtrex 80W, or AAtrex 4L (4LC) for control of many annual weeds, including carpetweed, crabgrass, fall panicum, foxtail, lambsquarters, morningglory, pigweed, ragweed, and velvetleaf.

Apply at the rates given below. Use the 1:1 ratio for most weeds and the 2:1 ratio for expected heavy infestations of crabgrass and fall panicum.

Soil texture	Broadcast rate per acre			
	1:1 Ratio		2:1 Ratio	
	Princep 80W	AAtrex 80W*	Princep 80W	AAtrex 80W*
Sand, loamy sand, sandy loam	1.25 lbs.	1.25 lbs.	1.67 lbs.	0.83 lb.
Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay, or silty clay with low organic matter	1.5 lbs.	1.5 lbs.	2 lbs.	1 lb.
Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay, or silty clay with medium to high organic matter, and clay (including dark prairie soils of the Corn Belt)	1.8 lbs.	1.8 lbs.	2.4 lbs.	1.2 lbs.

*When using AAtrex 4L (4LC) or AAtrex Nine-O, use equivalent rates. One lb. of AAtrex 80W equals 0.8 qt. of AAtrex 4L (4LC) or 0.9 lb. of AAtrex Nine-O.

Refer to the AAtrex label for complete directions. Also refer to the AAtrex label and the Corn section of this label for precautions and limitations.

Eradicane®: Use in a preplant incorporated tank mixture for control of all weeds claimed on both the Princep and Eradicane labels including partial control (suppression) of shattercane (wild cane). Fluid fertilizer may replace all or part of the water in the spray. Check the physical compatibility of mixture with fertilizer before use. Refer to the General Information section of this label for a compatibility test procedure. Use a minimum of 20 gals. of spray volume per acre. Refer to the Eradicane label for incorporation directions. Use the higher rate of Princep on fine-textured soil and where heavy broadleaf infestations are expected. Use the higher rate of Eradicane for heavy nutsedge and bermudagrass infestations. For partial control (suppression) of shattercane, broadcast and incorporate immediately before planting 1.25-2.5 lbs. of Princep 80W plus 7.33 pts. of Eradicane 6.7E (or 8 pts. Eradicane Extra) per acre. For control of the other weeds claimed on both labels, broadcast and incorporate 1.25-3.75 lbs. of Princep 80W plus 4.75-7.33 pts. of Eradicane 6.7E (or 5.33-8 pts. of Eradicane Extra) per acre. Observe all precautions and limitations on the Princep 80W and Eradicane labels.

Paraquat: Use in a tank mixture where corn will be planted directly in a cover crop, established sod, or previous crop residues. This combination controls existing vegetation and provides residual control of the annual broadleaf and grass weeds listed under General Information.

Add Princep 80W to the spray tank, mix thoroughly with water, and then add paraquat and a nonionic surfactant. Provide constant agitation during mixing and application to keep the mixture in suspension.

Apply 2.5-3.75 lbs. of Princep 80W plus 1-2 pts. of paraquat* in 20-60 gals. of water per acre as a broadcast spray either before or after planting but before corn emerges. Add a non-ionic surfactant at the rate of 0.5 pt. per 100 gals. of spray volume.

For further information, see general information, caution and warning statements, precautions, and notes on the Princep 80W and the paraquat labels.

*Based on a product containing 2 lbs. paraquat cation per gal.