

AAtrex[®]

80W

Herbicide

For season-long
weed control in corn
and sorghum

For weed control in
certain other crops;
in noncrop areas;
and industrial sites

Five Pounds
Net Weight

Active Ingredients:
Atrazine: 2-chloro-
4-ethylamino-6-
isopropylamino-
s-triazine 76%
Related
compounds 4%
Inert Ingredients: 20%
Total: 100%

EPA Reg. No. 100-439

AAtrex 80W is a
wettable powder

Keep Out of Reach
of Children.

Caution

See additional
precautionary
statements on
side of bag.

See directions for use
on panel at right.

CIBA-GEIGY

EPA Est. 100-AL-1

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the Directions for Use and Conditions of Sale and Warranty before using this product.

Conditions of Sale and Warranty

The Directions for Use of this product are based on the best available information at the time of publication. The manufacturer does not warrant that the product will perform under all conditions of use. The manufacturer does not warrant that the product will perform under all conditions of use. The manufacturer does not warrant that the product will perform under all conditions of use.

CIBA-GEIGY warrants that this product conforms to the specifications in the Directions for Use and Conditions of Sale and Warranty. CIBA-GEIGY makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall CIBA-GEIGY or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product.

Directions for Use

This is a violation of Federal Law to use this product in a manner not as intended.

FAILURE TO FOLLOW ALL PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES

General Information

This herbicide controls many annual broadleaf and grass weeds in corn, sorghum, sugarcane, rice, and other crops. It is a contact herbicide. It is a contact herbicide. It is a contact herbicide.

For best results, use this product in a manner consistent with the Directions for Use. Do not use this product in a manner not as intended. Do not use this product in a manner not as intended.

Since this product acts through contact, its effectiveness depends on the ability of the herbicide to reach the weeds. Do not use this product in a manner not as intended.

Try to avoid injury to economic and desirable plants. Do not use this product in a manner not as intended. Do not use this product in a manner not as intended.

When the use directions cover a range of rates, use the higher rate on coarse textured soils and the lower rate on fine textured soils.

Use CIBA-GEIGY product recommendations and labels in conjunction with other herbicides only as expressly permitted on the label.

Application Procedures

Ground application: Use conventional ground sprayers equipped with nozzles that provide accurate and uniform application. Be certain that nozzles are clean, unobstructed, and are the same size. Do not use a spray nozzle that is larger than the nozzle specified on the label.

Aerial application: Use an aircraft with the proper spray application system. Do not use a spray nozzle that is larger than the nozzle specified on the label. Do not use a spray nozzle that is larger than the nozzle specified on the label.

Use of CIBA-GEIGY products in conjunction with other herbicides: Do not use this product in a manner not as intended. Do not use this product in a manner not as intended.

Use of CIBA-GEIGY products in conjunction with other herbicides: Do not use this product in a manner not as intended. Do not use this product in a manner not as intended.

Use of CIBA-GEIGY products in conjunction with other herbicides: Do not use this product in a manner not as intended. Do not use this product in a manner not as intended.

Use of CIBA-GEIGY products in conjunction with other herbicides: Do not use this product in a manner not as intended. Do not use this product in a manner not as intended.

Use of CIBA-GEIGY products in conjunction with other herbicides: Do not use this product in a manner not as intended. Do not use this product in a manner not as intended.

Use of CIBA-GEIGY products in conjunction with other herbicides: Do not use this product in a manner not as intended. Do not use this product in a manner not as intended.

Use of CIBA-GEIGY products in conjunction with other herbicides: Do not use this product in a manner not as intended. Do not use this product in a manner not as intended.

Use of CIBA-GEIGY products in conjunction with other herbicides: Do not use this product in a manner not as intended. Do not use this product in a manner not as intended.

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Corn

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Table 1. Broadleaf and Grass Weed Control on Corn*

Soil texture	Broadcast rate A
COARSE Sandy loam, sandy loam, sandy clay loam	2.5 b
MEDIUM Loam, silty loam, silty clay loam, sandy loam, silty clay loam, sandy clay loam, silty clay loam, silty clay loam	3.0
FINE Loam, silty loam, silty clay loam, sandy loam, silty clay loam, sandy clay loam, silty clay loam, silty clay loam	3.75 b
Very heavy, high organic clay, high clay content	3.75 b

*Broadcast rate: eastern CO, western AS, western NE, NW, OK, Panhandle, west TX, and eastern WI, OK, and sandy soil, sandy loam, mid to strong alkaline soil, and a recent heavy frost. Apply no more than 1.5 lb A per acre per application. On other soils in these areas, apply rate in Table 1 for broadcast and preplant.

Preplant (Broadleaf and grass control): Broadcast 2.5 to 3.75 lb A per acre before or after tillage. Do not use this product in a manner not as intended.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Postemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

Preplant (Broadleaf and grass control): Broadcast preplant after tillage in the field. Apply before planting or after the seedbed preparation. Apply after the weeds have emerged.

Preemergence (Broadleaf and grass control): Apply after the weeds have emerged. Do not use this product in a manner not as intended.

EST DOCUMENT AVAILABLE

Weeds are present in orchards and citrus groves. Use the use information on this label to determine the treatment of the orchard and citrus grove.

Use AATree-BOW in orchards and citrus groves to control weeds. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

Precautions: Do not use in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves.

Note: Do not graze treated areas within 6 months after application.

Aerial application: In orchards and citrus groves, apply AATree-BOW to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

Precaution:

For best results, apply AATree-BOW to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

Precaution: Use only one application per year.

Rotational crops: See Rotational crops in Con section.

Rangeland:

To control certain weeds, including cheatgrass, apply AATree-BOW to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

Aerial application: In orchards and citrus groves, apply AATree-BOW to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

Establishment of perennial range grasses in West (CA, ID, NV, OR, UT, WA): Broadcast 1.25 lb AATree-BOW per acre after the ground has been prepared. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

Renovation of existing stands of perennial range grasses: Central Great Plains (CO, KS, MT, NE, ND, SD, WY): Broadcast 0.6 lb AATree-BOW per acre before ground freezes. Southern Great Plains (OK, TX): Broadcast 1.25 lb AATree-BOW per acre in spring before April 30.

West (CA, ID, NV, OR, UT, WA): Concentrate AATree-BOW on the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

E. grass and intermediate wheatgrass may be included in this production of intermediate grasses.

Note: Do not graze treated areas within 6 months after application. Do not graze treated areas within 6 months after application. Do not graze treated areas within 6 months after application.

Warm season perennial range grasses — Midwest: Switchgrass and Big Bluestem: For control of weeds, apply AATree-BOW to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

Use of AATree-BOW in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves.

New seeding: Broadcast 2.5 lb AATree-BOW per acre after the ground has been prepared. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

Established orchard grass green for seed (Pacific Northwest Only): AATree-BOW is used to control weeds in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves.

Note: Do not graze treated areas within 6 months after application. Do not graze treated areas within 6 months after application. Do not graze treated areas within 6 months after application.

Rotational crops: See Rotational crops in Con section.

Perennial Ryegrass Green for Seed (Pacific Northwest Only):

For control of weeds, apply AATree-BOW to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

Precautions: Do not use in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves.

Note: Do not graze treated areas within 6 months after application. Do not graze treated areas within 6 months after application. Do not graze treated areas within 6 months after application.

Established Orchard Grass Green for Seed (Pacific Northwest Only):

AATree-BOW is used to control weeds in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves.

Precautions: Do not use in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves.

Note: Do not graze treated areas within 6 months after application. Do not graze treated areas within 6 months after application. Do not graze treated areas within 6 months after application.

Sugarcane:

For control of weeds, apply AATree-BOW to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

Additional applications may be made after the emergence of weeds. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

Note: AATree-BOW is used to control weeds in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves.

Aerial application: In orchards and citrus groves, apply AATree-BOW to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

Florida:

For control of weeds, apply AATree-BOW to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

For control of weeds, apply AATree-BOW to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

1. Apply 5 lb AATree-BOW per acre before weeds emerge. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

2. Apply 1.3 times as needed at 2.5 lb AATree-BOW per acre after weeds emerge. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

Texas:

Use AATree-BOW to control weeds in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves.

Apply 5 lb AATree-BOW per acre before weeds emerge. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

For best results when weeds are emerged, add a nonionic surfactant at a concentration of 2 to 100 gal to the spray and apply before weeds exceed 1.5 inches in height.

Precautions: Do not use in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves.

Turf Grasses for Sod:

St. Augustinegrass, Centipedegrass, and Zoysia Grass: Broadcast 2.5 lb AATree-BOW per acre to control weeds. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

Muck or peat	5 lb	0 to 2 days after mowing sod. New sods 3-4 days after spring or planting.
Sandy soil	2.5 lb	0 to 2 days after mowing sod. New sods 3-4 days after spring or planting.

Weeds should be applied additional 2.5 lb AATree-BOW per acre in muck or peat or 1.25 lb AATree-BOW per acre in sandy soil.

Precautions: Do not use in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves.

Macadamia Nuts:

For control of weeds, apply AATree-BOW to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

Precautions: Do not use in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves.

Pineapple:

For control of weeds, apply AATree-BOW to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove. Do not apply to the ground in the orchard or citrus grove.

Precautions: Do not use in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves.

Note: Do not graze treated areas within 6 months after application. Do not graze treated areas within 6 months after application. Do not graze treated areas within 6 months after application.

Guava:

Use of AATree-BOW in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves. Do not use in orchards and citrus groves.

Do not use in orchards and citrus groves.

Comments

For control of annual broadleaf and grass weeds or for pre-planting aftertransplanting or in established fields, this includes: Dog's head, groundnut, white, yellow and red top, pigweed, and other broadleaf weeds. In established fields, this includes: pigweed, morning glory, groundnut, and other broadleaf weeds. For control of annual broadleaf and grass weeds, broadcast 2.5 to 4 lb (1.1 to 1.8 kg) of water per acre (or 10 to 15 gal of water per acre) before weeds are 15 inches tall. Apply to established weeds before and early spring when weeds are dormant. For pre-planting, apply during or soon after transplanting. For applications prior to transplanting, a low surfactant rate is not to activate AAtres before transplanting. In areas where spring and summer rainfall is inadequate to activate AAtres, apply during fall prior to spring transplanting.

For the most weeds controlled, see AAtres 60W Applied Alone - Corn or Grain Sorghum - Preplant Incorporated or Preemergence.

Quackgrass control: Broadcast 5 lb (2.3 kg) of water per acre (or 10 to 15 gal of water per acre) before and early spring when weeds are dormant and before quackgrass is more than 15 inches tall.

Note: Temporary injury to trees may occur following the use of AAtres on coarse textured soil.

Precautions: (1) Do not graze treated areas. (2) Do not apply to seedbeds. (3) Apply only once per year.

Aerial application: In order to assure that spray will be complete within the target area when used according to label directions, make applications at a maximum height of 10 ft above vegetation, using no wind forces at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply AAtres 60W by aircraft at a minimum ground speed of 400 ft/min from sensitive plants.

Non-selective Weed Control on Noncrop Land

This product provides long-term weed control on industrial sites, grain elevators and yards, roads, rights-of-way, lumberyards, petroleum refineries, and other noncrop areas on farms such as around buildings, equipment and the storage areas along fences and areas. Do not use near adjacent desirable plants or in greenhouses or in any manner.

Apply before or soon after weeds begin to grow. Make post-emergence applications when weeds are 4 to 10 inches and actively growing. Use when temperatures are 50 to 80°F. Use 10 to 15 gal of water per acre to assure thorough coverage. Use 10 to 15 gal of water for each pound of product. Use in all practices.

Note: Do not graze livestock on treated areas.

Annual broadleaf and grass weeds: including: barnyardgrass, crabgrass, foxtail, grass, amaranth, pigweed, ragweed, sunflower, and yellow foxtail. Broadcast 2.5 to 4 lb (1.1 to 1.8 kg) of water per acre.

Hard-to-kill annual and many perennial broadleaf and grass weeds: including: quackgrass, ryegrass, Canada thistle, dogbane, foxtail, grass, plantain, quackgrass, purple top, redtop, and amaranth. Broadcast 2.5 to 4 lb (1.1 to 1.8 kg) of water per acre.

Hard-to-kill biennial and perennial weeds: including: thistle and sowthistle. Broadcast 2.5 to 4 lb (1.1 to 1.8 kg) of water per acre.

For long-term residual control in grain areas, including growing season, broadcast 2.5 to 4 lb (1.1 to 1.8 kg) of water per acre.

For final areas, 4.5 to 6 lb (2.0 to 2.7 kg) of water per acre.

Storage and Disposal

Store in a dry place. Do not contaminate water, food, or feed by storage or disposal. Cleaning of equipment. Open burning is prohibited. Pesticide sprays, mixtures, or residues that cannot be used according to label instructions must be disposed of according to applicable federal, state, or local procedures. Completely empty bags by shaking and rapping sides and bottom to remove all residues. Empty bags must be placed in a secure container. Do not dispose of bags in sanitary and other non-designated areas. Follow by state and local authorities.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. May be absorbed through skin. Do not breathe dust or spray. Avoid contact with eyes, skin, or clothing.

First Aid: If swallowed, contact your local poison control center, hospital, or physician. If the patient is unconscious, maintain breathing and heartbeat. Cardiac monitoring is essential. If the patient is conscious and alert, induce vomiting. Swallowing of contents may irritate the mouth and throat. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

In case of skin contact, remove contaminated clothing and wash with soap and water. In case of contact with eyes, flush with plenty of water. Get medical attention if irritation persists.

Note to Physician: There is no specific antidote for AAtres. This product is ingested in case of acute or repeated oral use. The acute oral LD50 is 1.5 g/kg (0.675 lb/kg) in rats. The acute oral LD50 is 1.5 g/kg (0.675 lb/kg) in rats.

Environmental Hazards

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

AAtres[®] trademark of CIBA GEIGY for AAtres.
Baxton[®] trademark of Dow Chemical Co. for propachlor.
Ca-ben[®] trademark of CIBA GEIGY.
Concept[®] trademark of CIBA GEIGY.
Dua[®] trademark of CIBA GEIGY for metolachlor.
U.S. Patent No. 3,937,730.
Igran[®] trademark of CIBA GEIGY for terbutryn.
U.S. Patent No. 3,634,062.
Lasso[®] trademark of Monsanto Company for alachlor.
Princep[®] trademark of CIBA GEIGY for simazine.
Ramrod[®] trademark of Monsanto Company for propachlor.
Roundup[®] trademark of Monsanto Company for glyphosate herbicide.
Sutan[®] trademark of Stauffer Chemical Co. for butyltin.
X-77[®] trademark of NALCO Laboratories, Inc.
©1982 CIBA GEIGY Corporation
Agricultural Division
CIBA GEIGY Corporation
Greensboro, North Carolina 27419
CGA 70290