

D·z·n[®]

Lawn & Garden Insect Control

This product must not be used on golf courses and sod farms.

Spray concentrate. Contains diazinon insecticide.

Active Ingredient:
Diazinon: O, O-Diethyl
O-(2-Isopropyl-6-
methyl-4-pyrimidinyl)
phosphor-
thioate 25.0%

Inert
Ingredients* : 75.0%

Total: 100.0%

* Contains xylene range aromatic solvent.

KEEP OUT OF REACH OF CHILDREN.

WARNING

See side panel for additional precautionary statements.

EPA Reg. No. 100-456

Protects
Lawns
Vegetables
Fruits and Nuts
Roses, Flowers
Trees, Shrubs

Controls
Ants, Aphids
Chinch Bugs
Cutworms
Fire Ants*
Flea Beetles ∴∴
Grubs, Mites . . .
Sod Webworms
...and many others

* Aids in control
See full label in booklet under cap for complete directions for mixing and applying this product, and a complete list of pests controlled.

One Pint
Net Contents

ACCEPTED
SEP 17 1990
Made for the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the products
registered under
EPA Reg. No. 100-456

CIBA-GEIGY

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

General Information

D·z·n Lawn & Garden Insect Control is an emulsifiable concentrate product which must be diluted with water before application. This product kills many outdoor pests which damage lawns, roses, flowers, trees, shrubs, vegetables and fruits and insects which annoy pets and people.

SEE THE BOOKLET UNDER CAP FOR COMPLETE DIRECTIONS FOR USE AND PESTS CONTROLLED.

Notes: (1) USE CONTENTS WITHIN 6 MONTHS AFTER OPENING. (2) Do not use in the home. (3) Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia, or orchids since plant injury may occur. (4) Do not use on commercially-grown potatoes which will be hand-harvested. (5) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties.

After spraying wait the following number of days before harvesting fruits, nuts, or vegetables:

| | |
|----------|---|
| Same day | Peas, Sweet Corn |
| 1 day | Tomatoes |
| 3 days | Melons, Winter Squash |
| 5 days | Broccoli, Cauliflower, Peppers, Strawberries |
| 7 days | Beans, Blackberries, Boysenberries, Cabbage, Cranberries, Cucumbers, Dewberries, Loganberries, Summer Squash |
| 10 days | Apricots, Carrots, Celery, Cherries, Chinese Broccoli, Chinese Cabbage, Collards, Endive, Kale, Lettuce, Nectannes, Onions, Parsnips, Plums, Prunes, Radishes, Spinach, Turnips |
| 14 days | Beets, Red (Table) |
| 18 days | Grapes |
| 20 days | Peaches |
| 35 days | Potatoes (Irish) |
| 45 days | Almonds, Walnuts |

**Precautions: (1) Wear a long-sleeved shirt and long-legged pants when spraying. (2) Spray with the wind to your back. (3) Do not spray on windy days. (4) If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water and launder clothing before wearing again.*


**This table is only a partial listing of the pests controlled by this product. See booklet under cap for complete directions for use and pests controlled.*

| Problem Area | Pests* |
|---|---|
| Nuisance Pests in Outside Areas | Ants, Brown Dog Ticks, Chiggers, Cockroaches, Crickets, Fleas, Fire Ants, Spiders and others |
| Lawns | Chinch Bugs, Ants, Sod Webworms, Brown Dog Ticks, Fleas, White Grubs, Lawn Billbugs and others* |
| Vegetables: Foliar Application | |
| Beans, Peas (except dried varieties) | Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Mites |
| Broccoli, Chinese Broccoli, Cabbage, Chinese Cabbage, Cauliflower, Collards, Kale | Aphids, Flea Beetles, Dipterous Leafminers, Diamond-back Moths, Imported Cabbage Worms |
| Celery, Endive (Escarole), Lettuce, Spinach | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites |

D-z-n® LGIC

395

| | |
|--|---|
| Peppers. Tomatoes | Aphids, Colorado Potato Beetles, Fall Armyworms (Tomatoes), Flea Beetles, Leafhoppers, Southern Armyworms (Tomatoes), Vinegar Flies - <i>Drosophila</i> spp. (Tomatoes) |
| Onions (Bulb and Green) | Onion Thrips |
| Radishes, Turnips, Carrots, Parsnips, Potatoes | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Colorado Potato Beetles (Potatoes), Southern Armyworms (Potatoes) |
| Sweet Corn | Aphids, Flea Beetles, Grasshoppers, Mites |
| Melons, Squash (Summer, Winter) | Aphids, Thrips, Mites |
| Vegetables: Preplant Soil Application | |
| Beans, Peas (except dried varieties) | Cutworms, Mole Crickets, Wireworms (snap beans) |
| Beets, Red (Table) | Mole Crickets |
| Cabbage | Cutworms, Root Maggots, Wireworms |
| Carrots, Sweet Corn | Cutworms, Mole Crickets, Wireworms |
| Lettuce, Tomatoes | Cutworms, Wireworms |
| Radishes, Turnips | Cutworms |
| Fruits & Nuts | |
| Almonds, Walnuts (California only - both crops) | Aphids, Scale Crawlers, Spider Mites and others |
| Apricots, Peaches, Nectarines | Aphids, Apricot Mealybugs (Apricots), Spider Mites, Scale Crawlers |
| Cherries | Black Cherry Aphids, Cherry Fruitflies, Spider Mites, Scale Crawlers and others* |
| Plums, Prunes | Aphids, Mites |
| Caneberries (blackberries, boysenberries, dewberries, loganberries, raspberries) (California, Oregon, Washington only - all crops) | Aphids, Leafhoppers, Two-Spotted Mites, Thrips, Raspberry Fruitworms |
| Strawberries | Aphids, Two-Spotted Mites, Strawberry Leafrollers |
| Grapes | Aphids, Grape Berry Moths, Grape Leafrollers, Leafhoppers |
| Roses & Flowers | |
| Azalea, Camellia, Chrysanthemum, Carnation, Gladioli, Lilac, Rhododendron, Roses | Aphids, Flea Beetles, Dipterous Leafminers, Leafhoppers, Mealybugs, Two-spotted Spider Mites, Thrips, Whiteflies and others |



Trees & Shrubs

Arborvitae,
Azalea, Birch,
Boxwood, Camellia,
Douglas Fir, Elm,
Hawthorn Holly,
Juniper, Lilac,
Locust, Maple,
Mimosa, Oak, Pine,
Ornamental Plum,
Poplar,
Rhododendron,
Spruce, Willow

Aphids, Mites, Scale Crawlers, Whiteflies, Tent Caterpillars and others*

Storage and Disposal**Storage**

Do not store near heat or open flame. Store in original container in an area inaccessible to children or pets. USE CONTENTS WITHIN 6 MONTHS AFTER OPENING.

Disposal

Do not reuse empty container. Thoroughly rinse container, securely wrap in several layers of newspaper, and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

Precautionary Statements**Hazards to Humans and Domestic Animals****WARNING**

May be fatal if swallowed or absorbed through skin. Avoid breathing spray mist. Causes substantial but temporary eye damage. Do not get in eyes, on skin, or clothing. Wear goggles or safety glasses. Avoid contact with eyes and skin. Wash thoroughly if on skin and after handling. Do not use in the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not permit children or pets to go onto sprayed grass until the spray has completely dried. Do not contaminate ornamental fish ponds. Food utensils such as spoons and measuring cups must not be used for food purposes after use in measuring pesticides.

Statement of Practical Treatment

If swallowed: Call a physician immediately. DO NOT INDUCE VOMITING UNLESS NO OTHER TREATMENT IS AVAILABLE. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If inhaled: Remove victim to fresh air and apply artificial respiration if indicated.

If on skin: Wash promptly with soap and water. Rinse thoroughly.

If in eyes: Rinse eyes with plenty of water and call a physician immediately.

Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains a xylene range aromatic solvent which may present an aspiration

hazard. Gastric lavage may be indicated if product was taken internally.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. Where irrigation (watering) is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply a minimum of 1/4 inch of water immediately after applying this product. Stop irrigation before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.

D-z-n® LGIC

595

D-z-n® trademark of CIBA-GEIGY for diazinon

©1990 CIBA-GEIGY Corporation


Agricultural Division

CIBA-GEIGY Corporation

Greensboro, North Carolina 27419

CGA A1L1R18U

050



BEST AVAILABLE COPY