

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions For Use and the Conditions Of Sale And Warranty before using this product.

Conditions Of Sale And Warranty

The Directions For Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Geigy or the Seller. All such risks shall be assumed by the Buyer.

Geigy warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions For Use, subject to the inherent risks referred to above. Geigy makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Geigy or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. Geigy and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions Of Sale And Warranty which may be varied only by agreement in writing signed by a duly authorized representative of Geigy.

Method of Application

Planter-Box Treatment:

Corn:

Seed Corn Beetles: Mix 3 oz. per bushel of seed as a dry mixture in the planter box as a seed treatment prior to planting.

Corn, Peas, ~~Pumpkins~~ ^{9/21}, Soybeans and Beans (Lima, Snap, White and Kidney)

Seed Corn Maggots: Mix 1-2 oz. per bushel of seed as a dry mixture in the planter box as a seed treatment prior to planting.

Planter-Box Mixing Instructions: Stir seed slightly using a mixing stick. Additional mixing will result from the operation of the planter. DO NOT MIX WITH BARE HANDS.

Note: If seed has not been treated with a fungicide, use Captan or Thiram in combination with Diazinon 50W at rates indicated on fungicide package. Failure to use a fungicide with Diazinon may result in injury to the seed. If the seed treatment increases the friction between seeds sufficiently to reduce the rate of delivery through the planter, 1-2 oz. of graphite per bushel of seed should be added as a drill lubricant. Adjust the planter to give the required rate of seed delivery with the treated seed.

Caution: Diazinon treated grain cannot be used for feed for livestock, or for the preparation of products for human consumption.

USDA Reg. No. 100-460

U. S. Patent No. 2,754,243

GEIGY
Geigy Agricultural Chemicals
Division of Geigy Chemical Corporation
Ardsley, New York

March 26, 1969