

Crop	Pest	Rate	Timing/Special Directions
Cranberries (Cont'd)	Cranberry Fruitworms	4 to 6 lbs./A	Cranberry growth stage and/or berry inspection for eggs may be used to optimize treatment timing; consult your local extension agent or pest management advisor. Repeat application every 14 days as needed. A maximum of 4 applications per season is permitted for the 6 lb. rate. A maximum of 6 applications per season is permitted for the 4 lb. rate.

Notes: (1) This product may be applied to cranberries via ground, aerial, or sprinkler irrigation. For ground application, apply the product in a minimum of 15 gals. of water/A, for aerial application, a minimum of 20 gals. of water/A, for chemigation up to 400 gals. of water/A. Refer to the "Chemigation" section of this label for specific chemigation directions. (2) Do not apply more than 24 lbs. of product per acre per season. (3) Allow a minimum of 7 days between the last application and harvest.

Grapes	Pacific Spider Mites, Leafhoppers, Grape Berry Moths, Grape Leaf-folders, Omnivorous Leafrollers, Aphids, Spider Mites	1-2 lbs. per acre in a minimum of 100 gals. of water (200 gals. of water maximum)	Apply as a thorough cover spray when pest appears. Repeat application after 7 days if necessary.
--------	--	---	--

Crop	Pest	Rate	Timing/Special Directions
Grapes (Cont'd)	<i>Drosophila</i> spp.	6 oz. per acre in a minimum of 100 gals. of water (300 gals. of water maximum)	Apply as a thorough cover spray at 7-day intervals.

Notes: (1) Do not apply more than 2 lbs. product per acre per application and no more than 10 lbs. product per acre per season. (2) To avoid illegal residues, allow a minimum of 28 days between the last application and harvest.

Nectarines	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, European Red Mite Eggs, Twig Borers, Apple Aphid Eggs, Black Cherry Aphid Eggs, Mealy Plum Aphid Eggs, Mealybugs	1 lb. + 2-3 gals. of dormant oil or 1-1 1/2 gals. superior type oil in 100 gals. of water	Apply as a dormant spray. Do not apply more than 4 lbs. of product or 6 gals. of oil per acre.
	Aphids, Brown Mites (Clover Mites), Two-Spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers	1 lb. in 100 gals. of water	Apply as infestations occur. Repeat application after 7 days if necessary.
	Peach Twig Borers, Parlatoria Scale	1 lb. + 1 gal. light medium summer horticultural oil in 100 gals. of water	Apply at petal fall. Do not apply more than 4 lbs. of product or 4 gals. of summer oil per acre.

Crop	Pest	Rate	Timing/Special Directions
Nectarines (Cont'd)	Olive Scale Crawlers	1/2 lb. + 1 1/2 gals. light medium horti- cultural oil in 100 gals. of water	Apply as insects occur. Do not apply more than 4 lbs. of product or 4 gals. of summer oil per acre per applica- tion. Allow 7 days between applications.
	Oriental Fruitmoths	1 lb. in 100 gals. of water	Apply at 100% petal fall. Make 2 additional applications at 10-15 day inter- vals. Time the last application to peak adult flight.

Notes: (1) Do not apply more than 4 lbs. of product per acre per application and no more than 12 lbs. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.

Peaches	San Jose Scale, Parlatoria Scale Black Scale, Brown Scale, Apricot Scale, European Red Mite Eggs, Brown Mite Eggs, Twig Borers, Apple Aphid Eggs, Black Cherry Aphid Eggs, Mealy Plum Aphid Eggs, Mealybugs	1 lb. + 2-3 gals. dormant oil or 1-1 1/2 gals. superior type oil in 100 gals. of water	Apply as a dormant spray. Do not apply more than 4 lbs. of product or 6 gals. of oil per acre.
---------	--	---	--

33/05

Crop	Pest	Rate	Timing/Special Directions
Peaches (Cont'd)	Leafhoppers	1 lb. in 100 gals. of water	Make 3-4 applications beginning at petal fall and continuing through the growing season. Allow at least 7 days between applications.
	White Peach Scale	1 1/2-2 lbs. in 100 gals. of water	Apply post harvest timed to coincide with peak crawler and immature scale activity. Under conditions of heavy infestation 2 applications (7 days apart) may be needed.

Notes: (1) Do not apply more than 4 lbs. of product per acre per application and no more than 12 lbs. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.

Pears	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite Eggs, Brown Mite Eggs, Pear Leaf Blister Mite Eggs, Twig Borers, Apple Aphid Eggs, Black Cherry Aphid Eggs, Mealy Plum Aphid Eggs, Mealybugs	1 lb. + 2-3 gals. dormant oil or 1-1 1/2 gals. superior type oil in 100 gals. of water	Apply as a dormant spray. Do not apply more than 4 lbs. of product or 6 gals. of oil per acre.
-------	---	---	--

Crop	Pest	Rate	Timing/Special Directions
Pears (Cont'd)	Pear Psylla	1/2-1 lb. in 100 gals. of water	Apply as a cover spray at pre-pink stage. Make 1-2 additional appli- cations during the first and second cover sprays, if neces- sary. Allow at least 14 days between applica- tions.
	Tentiform Leafminers, Mealybugs	1 lb. in 100 gals. of water	Make 3 cover sprays, beginning in April. Allow at least 14 days between applica- tions.
	Codling Moths, Rosy Apple Aphids, Green Apple Aphids, Wooly Apple Aphids, Spider Mites	1 lb. in 100 gals. of water	Make 3 cover sprays beginning as infestations occur. Allow at least 14 days between applica- tions. Note: Mites will be suppressed if diazinon is used in this regular cover spray pro- gram.
	Apple Maggots	1 lb. in 100 gals. of water	Make 3 cover sprays when flies are active and laying eggs. Allow at least 14 days between applications. Note: Diazinon will not control organophosphate- resistant apple maggot.

Crop	Pest	Rate	Timing/Special Directions
Peaches (Cont'd)	Aphids, Brown Mites (Clover Mites), Two-Spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers	1 lb. in 100 gals. of water	Apply as infestations occur. Repeat application after 14 days if necessary.
	Peach Twig Borers, Parlatoria Scale	1 lb. + 1 gal. light medium summer horti- cultural oil in 100 gals. of water	Apply at petal fall. Do not apply more than 4 lbs. of product or 4 gals. of oil per acre.
	Olive Scale Crawlers	1/2 lb. + 1 1/2 gals. light medium horti- cultural oil in 100 gals. of water	Apply as insects occur. Do not apply more than 4 lbs. of product or 4 gals. of summer oil per acre per applica- tion. Allow 7 days between applications.
	Oriental Fruit Moths	1 lb. in 100 gals. of water	Apply at 100% petal fall. Make 2 additional applications at 10-15 day inter- vals. Time the last application to peak adult flight.
	Peach Twig Borers	1 lb. in 100 gals. of water	Apply as insects occur in May and June. Allow 7 days between applications.

26/82

Crop	Pest	Rate	Timing/Special Directions
Pears (Cont'd)	San Jose Scale Crawlers, Forbes Scale Crawlers	1 lb. in 100 gals. of water	Apply early in the season when crawlers first appear. If infestation warrants, make a second applica- tion 14 days later.
<p>Notes: (1) Do not apply more than 4 lbs. of product per acre per application and no more than 12 lbs. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.</p>			
Pineapple	Pineapple Mealybugs	2 lbs. per acre in 200-500 gals. of water	Repeat applica- tion as necessary every 28 days up to a maximum of 8 applications. Do not apply within 7 days of harvest or illegal resi- dues may occur.
Plums	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite Eggs, Brown Mite Eggs, Twig Borers, Apple Aphid Eggs, Black Cherry Aphid Eggs, Mealy Plum Aphid Eggs, Mealybugs	1 lb. + 2-3 gals. dormant oil or 1-1 1/2 gals. superior type oil in 100 gals. of water	Apply as a dor- mant spray. Do not apply more than 4 lbs. of product or 5 gals. of oil per acre. If applied in late April or May, mix diazinon with 1 gal. of light medium summer horticul- tural oil per 100 gals. of water.

27/28

Crop	Pest	Rate	Timing/Special Directions
Plums (Cont'd)	Brown Mites, Clover Mites, European Red Mites	1 lb. in 100 gals. of water	Apply as needed every 7 days.
	Leafcurl Plum Aphids, Mealy Plum Aphids, Thistle Aphids	1/2-1 lb. in 100 gals. of water	

Notes: (1) Do not apply more than 4 lbs. of product per acre per application and no more than 12 lbs. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.

Prunes	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite Eggs, Brown Mite Eggs, Twig Borers, Apple Aphid Eggs, Black Cherry Aphid Eggs, Mealy Plum Aphid Eggs, Mealybugs	1 lb. + 2-3 gals. dormant oil or 1-1 1/2 gals. superior type oil in 100 gals. of water	Apply as a dormant spray. Do not apply more than 4 lbs. of product or 6 gals. of oil per acre. For late April or May applications, mix diaiznon with 1 gal. of light medium summer horticultural oil per 100 gals. of water.
	Brown Mites, Clover Mites, European Red Mites	1 lb. in 100 gals. of water	Apply as needed every 7 days.
	Leafcurl Plum Aphids, Mealy Plum Aphids, Thistle Aphids	1/2-1 lb. in 100 gals. of water	

Notes: (1) Do not apply more than 4 lbs. of product per acre per application and no more than 12 lbs. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.

Crop	Pest	Rate	Timing/Special Directions
Straw-berries	Aphids, Two-Spotted Spider Mites	1 lb. in 100 gals. of water per acre	Apply as pests occur. Repeat application, if necessary, after 7 days.
	Cyclamen Mites	2 lbs. in 100 gals. of water per acre	Direct spray to the plant crowns. Plants should be sufficiently agitated to assure thorough coverage of the foliage. Repeat application after 7 days if necessary.
	Mole Crickets	2 lbs. per acre	Apply 1-2 days before trans- planting and immediately incorporate into the top 1-2 inches of soil.
	Strawberry Leafrollers	3/4-1 lb. in 100 gals. of water	Apply 200 gals. of spray per acre when blossoms show color.

Notes: (1) Do not make more than 3 foliar applications per season. (2) Do not apply more than 2 lbs. of product per acre per application and no more than 8 lbs. of product per season. (3) Allow a minimum of 5 days between the last application and harvest.

Crop	Pest	Rate	Timing/Special Directions
Walnuts (CA only)	Aphids, Spider Mites	2 lbs. in 100 gals. of water per acre	Apply as a foliar spray as needed. Repeat applica- tion in 14 days if necessary. Do not make more than 3 applica- tions per season. Do not apply within 45 days of harvest or after husks open or illegal residues may occur.
	Codling Moths, Scale Crawlers, Walnut Caterpillars	6 lbs. in 100 gals. of water per acre	

Vegetable Crops

REI - 24 hours.

For foliar or soil applications, D·z·n diazinon 50W should always be applied as a spray in sufficient water to assure thorough coverage of the foliage or soil. D·z·n diazinon 50W may be applied aerially or by ground application as specified in the following table. The specific amount of product to be used appears under the separate directions for use for each crop.

Crop	Ground Application		Aerial Application Minimum Gals./Acre
	Minimum Gals./ Acre	OR Minimum Gals./ 1000 Sq. Ft.	
Beans (Lima, Pole, and Snap)	10	2	5
(succulent only)			
Beets, Red (Table)	10	2	5
Broccoli	10	2	5
Brussels Sprouts	10	2	5
Cabbage	10	2	5
Cantaloupes	5	1	5
Carrots	10	2	5
Casabas	5	1	5
Cauliflower	10	2	5
Chinese Broccoli	10	2	5
Chinese Cabbage	10	2	5
Chinese Mustard	10	2	5
Collards	10	2	5
Crenshaws	5	1	5
Cucumbers	5	1	5
Endive (Escarole)	10	2	5
Honeydew Melons	5	1	5
Kale	10	2	5
Lettuce (Head and Leaf)	10	2	5
Muskmelons	5	1	5
Mustard	10	2	5
Onions (Bulb and Green)	10	2	5
Parsley	10	2	5
Parsnips	10	2	5
Peas (succulent only)	10	2	5
Peppers	10	2	5
Persian Melons	5	1	5
Potatoes (Irish)*	10	2	5
Radishes	10	2	5
Spinach	10	2	5
Squash, Summer	5	1	5
Squash, Winter	5	1	5
Sweet Corn	20	4	5
Sweet Potatoes	10	2	5
Swiss Chard	10	2	5
Tomatoes	10	2	5
Turnips	10	2	5
Watermelons	5	1	5

*Do not use on commercially-grown potatoes which will be hand-harvested.

Observe stated time intervals between last application and harvest as well as intervals between and maximum numbers of applications per season.

Pre-Harvest Intervals (PHI*) for Foliar Applications

<u>Crop</u>	<u>PHI* (Days)</u>
Beans (Lima, Pole, and Snap)	7
Beets, Red (Table)	14
Broccoli	7
Brussels Sprouts	7
Cabbage	21
Cantaloupes	3
Carrots	14
Casabas	3
Cauliflower	7
Chinese Broccoli	10
Chinese Cabbage	10
Chinese Mustard	10
Collards	10
Crenshaws	3
Cucumbers	7
Endive (Escarole)	14
Honeydew Melons	3
Kale	10
Lettuce (Head and Leaf)	14
Muskmelons	3
Mustard	10
Onions (Bulb and Green)	14
Parsnips	14
Peas	7
Peppers	5
Persian Melons	3
Potatoes (Irish)	35
Radishes	14
Spinach	14
Squash, Summer	7
Squash, Winter	3
Sweet Corn	7
Swiss Chard	14
Tomatoes	1
Turnips (Roots and Tops)	14
Watermelons	3

*PHI=Pre-Harvest Interval or the number of days between the last application and harvest.

Notes: (1) **Soil Incorporation:** Following application of this product to control vegetable soil insects, immediately incorporate the product into the soil to the recommended depth (see **Timing/Special Directions**) using a rotary hoe, cultivator, disk, harrow, or other suitable means. For control of surface cutworms, incorporate 2-3 inches, for subterranean cutworms incorporate 3-6 inches. (2) **To Protect Bees:** Foliar applications of this product should not be made to peas or beans if these crops or weeds in the treatment area are in bloom, or to corn during the pollen shed period.

17/18

Crop	Pest	Rate	Timing/Special Directions
Beans (Lima, Pole, and Snap) (succulent only)	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorpora- tion Note.)
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Black Bean Aphids, Mexican Bean Beetles, Dipterous Leafminers	1-1 1/2 lbs. per acre or 2/5-3/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, every 5 days. Do not make more than 3 applications per season. Do not apply within 7 days of harvest or illegal resi- dues may occur.
	Cucumber Beetles	1/2-3/4 lb. per acre or 1/5-3/10 oz. per 1,000 sq. ft.	Bean forage or bean hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if for- age is to be cut for hay. Make cucumber beetle applica- tion in 100 gals. of water per acre. Note: Diazinon will not control organophosphate- resistant leaf- miners.

4/25

Crop	Pest	Rate	Timing/Special Directions
Beans (Lima, Pole and Snap) (succulent only) (Cont'd)	Leafhoppers, Spider Mites	1 lb. per acre or 2/5 oz. per 1,000 sq. ft.	Apply as pests occur. Repeat application as necessary every 5 days. Do not make more than 3 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. Bean forage or bean hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay.
Beets, Red (Table)	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	2 lbs. per acre or 3/4 oz. per 1,000 sq. ft.	Apply to plant beds or fields 1-2 days before planting, and immediately incorporate into the top 1-2 inches of soil
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before and immediately incorporate into the top 4-8 inches of soil.

Crop	Pest	Rate	Timing/Special Directions
Beets, Red (Table) (Cont'd)	Aphids, Dipterous Leafminers	1/2-1 lb. per acre or 1/5-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application as necessary up to 5 times with a minimum of 7 days between applica- tions. Do not apply within 14 days of harvest, or illegal resi- dues may occur. Note: Diazinon will not control organophosphate- resistant leaf- miners.
Broccoli	Root Maggots	4-6 lbs. per acre or 1 1/2-2 1/4 oz. per 1,000 sq. ft.	Broadcast just before planting or transplanting and immediately incorporate into the top 3-4 inches of soil.
		4-8 oz. in 50 gals. of water	In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply 1/2-1 cup (4 to 8 oz.) per plant by hand or by tractor mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.

Crop	Pest	Rate	Timing/Special Directions
Broccoli (Cont'd)	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	2 lbs. per acre or 3/4 oz. per 1,000 sq. ft.	Apply to plant beds or fields 1-2 days before planting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers	1/2-1 lb. per acre or 1/5-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, with up to 5 times, with a minimum of 7 days between applications. Do not apply within 7 days of harvest, or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Brussels Sprouts	Root Maggots	4-6 lbs. per acre or 1 1/2-2 1/4 oz. per 1,000 sq. ft.	Broadcast just before planting or transplanting and immediately incorporate into the top 3-4 inches of soil.

Crop	Pest	Rate	Timing/Special Directions
Brussels Sprouts (Cont'd)	Root Maggots (Cont'd)	4-8 oz. in 50 gals. of water	<p>In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply 1/2-1 cup (4 to 8 oz.) per plant by hand or by tractor mounted sprayer equipped with drop nozzles to direct spray to the base of the plant.</p> <p>Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.</p>
	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	2 lbs. per acre or 3/4 oz. per 1,000 sq. ft.	Apply to plant beds 1-2 days before planting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.

Crop	Pest	Rate	Timing/Special Directions
Brussels Sprouts (Cont'd)	Aphids, Diamondback Moths, Imported Cabbageworms	1/2-1 lb. per acre or 1/5-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, up to 5 times, with a minimum of 7 days between applications. Do not apply within 7 days of harvest or illegal residues may occur.
Cabbage	Root Maggots	4-6 lbs. per acre or 1 1/2-2 1/4 oz. per 1,000 sq. ft..	Broadcast just before planting or transplanting and immediately incorporate into the top 3-4 inches of soil.
		4-8 oz. in 50 gals. of water	In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply 1/2-1 cup (4 to 8 oz.) per plant by hand or by tractor mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.

40, 185
1

Crop	Pest	Rate	Timing/Special Directions
Cabbage (Cont'd)	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	2 lbs. per acre or 3/4 oz. per 1,000 sq. ft.	Apply to plant beds or fields 1-2 days before planting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers	1/2-1 lb. per acre or 1/5-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, up to 5 times, with a minimum of 7 days between applications. Do not apply within 21 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Carrots	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)

Crop	Pest	Rate	Timing/Special Directions
Carrots (Cont'd)	Mole Crickets	2 lbs. per acre or 3/4 oz. per 1,000 sq. ft.	Apply to plant beds or fields 1- 2 days before planting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Carrot Rust Fly	4 lbs. per acre or 1 1/2 oz. per 1,000 sq. ft.	Apply as a drench to the seed fur- row at planting.
	Aphids	1 lb. per acre or 2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, up to 5 times with a minimum of 7 days between applica- tions. Do not apply within 14 days of harvest, or ille- gal residues may occur.
Cauli- flower	Root Maggots	4-6 lbs. per acre or 1 1/2-2 1/4 oz. per 1,000 sq. ft.	Broadcast just before planting or transplanting and immediately incorporate into the top 3-4 inches of soil.

Crop	Pest	Rate	Timing/Special Directions
Cauliflower (Cont'd)	Root Maggots (Cont'd)	4-8 oz. in 50 gals. of water	<p>In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply 1/2-1 cup (4 to 8 oz.) per plant by hand or by tractor mounted sprayer equipped with drop nozzles to direct spray to the base of the plant.</p> <p>Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.</p>
	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	2 lbs. per acre or 3/4 oz. per 1,000 sq. ft.	Apply to plant beds 1-2 days before planting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate in the top 4-8 inches of soil.

Crop	Pest	Rate	Timing/Special Directions
Cauliflower (Cont'd)	Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers	1/2-1 lb. per acre or 1/5-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, up to 5 times, with a minimum of 7 days between applications. Do not apply within 7 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Chinese Broccoli, Chinese Cabbage (Bok Choy and Napa), Chinese Mustard (Gai Choy)	Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers	1/2-1 lb. per acre or 1/5-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, up to 5 times, with a minimum of 7 days between applications. Do not apply within 10 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Collards	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)

Crop	Pest	Rate	Timing/Special Directions
Cauliarids (Cont'd)	Mole Crickets	2 lbs. per acre or 3/4 oz. per 1,000 sq. ft.	Apply to plant beds 1-2 days before planting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Diamondback Moths, Imported Cabbageworms, Harlequin Cabbagebugs, Corn Earworms	1/2-1 lb. per acre or 1/5-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, up to 5 times, with a minimum of 7 days between applica- tions. Do not apply within 10 days of harvest or ille- gal residues may occur.
Cucumbers	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorpora- tion Note.)
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.

Crop	Pest	Rate	Timing/Special Directions
Cucumbers (Cont'd)	Aphids, Serpentine Leafminers, Thrips, Striped Cucumber Beetles, Melonworms	1 lb. per acre or 2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Endive (Escarole)	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	2 lbs. per acre or 3/4 oz. per 1,000 sq. ft.	Apply to plant beds or fields 1-2 days before planting or transplanting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate 4-8 inches into soil.

Crop	Pest	Rate	Timing/Special Directions
Endive (Escarole) (Cont'd)	Aphids, Dipterous Leafminers	1/2-1 lb. per acre or 1/5-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, up to 5 times with a minimum of 7 days between applications. Do not apply within 14 days of harvest. Note: Diazinon will not control organophosphate-resistant leafminers.
Kale	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	2 lbs. per acre or 3/4 oz. per 1,000 sq. ft.	Apply to plant beds 1-2 days before planting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.

Crop	Pest	Rate	Timing/Special Directions
Kale (Cont'd)	Aphids, Diamondback Moths, Imported Cabbageworms, Harlequin Cabbagebugs, Corn Earworms	1/2-1 lb. per acre or 1/5-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, up to 5 times, a mini- mum of 7 days be- tween applica- tions. Do not apply within 10 days of harvest or ille- gal residues may occur.
Lettuce (Head and Leaf)	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorpora- tion Note.)
	Mole Crickets	2 lbs. per acre or 3/4 oz. per 1,000 sq. ft.	Apply to plant beds 1-2 days before planting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers	1/2-1 lb. per acre or 1/5-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, up to 5 times with a minimum of 7 days between applica- tions. Do not apply within 14 days of harvest. Note: Diazinon will not control organophosphate- resistant leaf- miners.

5748
1

Crop	Pest	Rate	Timing/Special Directions
Melons (Cantaloupes, Casabas, Crenshaws, Honeydews, Musk- melons, Persians, and Hybrids of these) and Water- melons	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorpora- tion Note.)
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers, Thrips, Striped Cucumber Beetles, Melonworms, Leafhoppers, Spider Mites	1/2-1 1/2 lbs. per acre or 1/5-3/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 3 days of harvest or illegal resi- dues may occur. Note: Diazinon will not control organophosphate- resistant leaf- miners.
Mustard	Mole Crickets	2 lbs. per acre or 3/4 oz. per 1,000 sq. ft.	Apply to plant beds 1-2 days before planting and immediately incorporate into the top 1-2 inches of soil.

Crop	Pest	Rate	Timing/Special Directions
Mustard (Cont'd)	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers, Diamondback Moths, Imported Cabbageworms	1/2-1 lb. per acre or 1/5-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, up to 5 times, with a minimum of 7 days between applications. Do not apply within 10 days of harvest. Note: Diazinon will not control organophosphate-resistant leafminers.
Onions (bulb and green)	Onion Maggots	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and mix into the top 3-4 inches of soil. (See Soil Incorporation Note.) Note: Diazinon will not control organophosphate-resistant onion maggots.
	Onion Thrips	1 lb. per acre or 2/5 oz. per 1,000 sq. ft.	Apply as thrips appear. Repeat after 7 days, if necessary. Do not make more than 3 applications per season. Do not apply within 14 days of harvest or illegal residues may occur.

Crop	Pest	Rate	Timing/Special Directions
Onions (bulb and green) (Cont'd)	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Parsley	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Parsnips	Aphids, Flea Beetles, Dipterous Leafminers	1/2-1 lb. per acre or 1/5-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, up to 5 times, with a minimum of 7 days between applications. Do not apply within 14 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Peas (succulent only)	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)

6-7-85

Crop	Pest	Rate	Timing/Special Directions
Peas (succulent only) (Cont'd)	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorpora- tion Note.)
	Aphids, Dipterous Leafminers	3/4-1 lb. per acre or 3/10-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, every 5 days. Do not make more than 3 applications per season. To avoid illegal residues, allow a minimum of 7 days between the last applica- tion and harvest. Vines and hay may be fed to dairy and beef cattle and sheep. To avoid illegal residues, wait 7 days following treatment if forage is to be cut for hay. Note: Diazinon will not control organophosphate- resistant leaf- miners.
Peppers	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorpora- tion Note.)

Crop	Pest	Rate	Timing/Special Directions
Peppers (Cont'd)	Mole Crickets	2 lbs. per acre or 3/4 oz. per 1,000 sq. ft.	Apply to plant beds 1-2 days before planting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Serpentine Leafminers	1/2 lb. per acre or 1/5 oz. per 1,000 sq. ft.	Apply as insects appear. Repeat application after 7 days if necessary. Do not make more than 5 applications per season. Do not apply within 5 days of harvest, or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Potatoes	Cutworms (Surface and Subterranean)	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)

Crop	Pest	Rate	Timing/Special Directions
Potatoes (Cont'd)	Southern Potato Wireworms (Southeastern U.S.)	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	For potatoes grown on mineral soils in the Southeast, apply prior to planting and immediately incorporate into the top 3-6 inches of soil depending on location of wireworms.
	Aphids, Flea Beetles, Colorado Potato Beetles, Dipterous Leafminers	1/2-3/4 lb. per acre or 1/5-3/10 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, up to 5 times with a minimum of 7 days between applications. Do not apply within 35 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
	Banded Cucumber Beetles, Leafhoppers, Southern Armyworms	3/4-1 lb. per acre or 3/10-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, up to 5 times with a minimum of 7 days between applications. Do not apply within 35 days of harvest or illegal residues may occur.

Note: Do not use on commercially-grown potatoes which will be hand-harvested.

Crop	Pest	Rate	Timing/Special Directions
Radishes	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	2 lbs. per acre or 3/4 oz. per 1,000 sq. ft.	Apply to plant beds or fields 1-2 days before planting and immediately incorporate in the top 1-2 inches of soil.
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Flea Beetles, Dipterous Leafminers	1/2-1 lb. per acre or 1/5-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, up to 3 times, with a minimum of 7 days between applications. Do not apply within 14 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.

Crop	Pest	Rate	Timing/Special Directions
Rutabagas	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	2 lbs. per acre or 3/4 oz. per 1,000 sq. ft.	Apply to plant beds or fields 1-2 days before planting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Flea Beetles, Dipterous Leafminers	1/2-1 lb. per acre or 1/5-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, up to 5 times, with a minimum of 7 days between applications. Do not apply within 14 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.

Crop	Pest	Rate	Timing/Special Directions
Spinach	Aphids, Dipterous Leafminers	1/2-1 lb. per acre or 1/5-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, up to 5 times, with a minimum of 7 days between applications. Do not apply within 14 days of harvest. Note: Diazinon will not control organophosphate-resistant leafminers.
Squash, Summer	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Leafminers, Thrips, Striped Cucumber Beetles, Melonworms, Leafhoppers, Spider Mites	1/2-1 1/2 lbs. per acre or 1/5-3/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 7 days of harvest, or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.

66/85

Crop	Pest	Rate	Timing/Special Directions
Squash, Winter	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Leafminers, Thrips, Striped Cucumber Beetles, Melonworms, Leafhoppers, Spider Mites	1/2-1 1/2 lbs. per acre or 1/5-3/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 3 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Sweet Corn	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Seed Corn Maggots	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 2 inches of soil.

Crop	Pest	Rate	Timing/Special Directions
Sweet Corn (Cont'd)	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Corn Earworms	2-2 1/2 lbs. per acre or 3/4-1 oz. per 1,000 sq. ft.	Apply when silks first appear. Repeat application every 7 days up to a maximum of 5 applications, if necessary.
	Corn Leaf Aphids	1-2 lbs. per acre or 2/5-3/4 oz. per 1,000 sq. ft.	Apply in sufficient water to provide complete coverage to the upper portions of the plants. Repeat in 7 days if necessary. Do not make more than 5 applications per season.
	Grasshoppers, Spider Mites, Fleabeetles	1 lb. per acre or 2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat if necessary, but do not make more than 5 applications per season.
	Sap Beetles	2-2 1/2 lbs. per acre or 3/4-1 oz. per 1,000 sq. ft.	Consult local agricultural authorities for proper timing of application. Repeat application in 7 days, if necessary. Do not make more 5 applications per season.

Notes: (1) To protect bees, do not apply foliarly to corn during the pollen shed period. (2) Allow a minimum of 7 days between the last application and grazing or cutting for forage. (3) Corn may be picked 7 days after the last application.

Crop	Pest	Rate	Timing/Special Directions
Sweet Potatoes	Wireworms, Flea Beetles	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Swiss Chard	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Corn Earworms, Imported Cabbageworms, Diamondback Moths, Harlequin Cabbagebugs	1/2-1 lb. per acre or 1/5-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, up to 5 times, with a minimum of 7 days between applications. Do not apply within 14 days of harvest or illegal residues may occur.
Tomatoes	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	2 lbs. per acre or 3/4 oz. per 1,000 sq. ft.	Apply to plant beds 1-2 days before planting and immediately incorporate into the top 1-2 inches of soil.

6985

Crop	Pest	Rate	Timing/Special Directions
Tomatoes (Cont'd)	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers	1/2 lb. per acre or 1/5 oz. per 1,000 sq. ft.	Apply as insects appear. Repeat application after 7 days, if necessary. Do not make more than 5 applications per season.
	Banded Cucumber Beetles, Fall Armyworms, Southern Armyworms, Beet Armyworms	3/4-1 lb. per acre or 3/10-2/5 oz. per 1,000 sq. ft.	Do not apply within one day of harvest or illegal residue may occur.
	Vinegar Flies (<i>Drosophila</i> spp.)	1/2-1 1/2 lbs. per acre or 1/5-3/5 oz. per 1,000 sq. ft.	Note: Diazinon will not control organophosphate-resistant leafminers.
Turnips (Roots and Tops)	Cutworms	4-8 lbs. per acre or 1 1/2-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	2 lbs. per acre or 3/4 oz. per 1,000 sq. ft.	Apply to plant beds or fields 1-2 days before planting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	6-8 lbs. per acre or 2 1/4-3 oz. per 1,000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.

Crop	Pest	Rate	Timing/Special Directions
Turnips (Roots and Tops) (Cont'd)	Aphids, Flea Beetles, Dipterous Leafminers	1/2-1 lb. per acre or 1/5-2/5 oz. per 1,000 sq. ft.	Apply as insects occur. Repeat application, as necessary, up to 5 times, with a minimum of 7 days between applica- tions. Do not apply within 14 days of harvest or ille- gal residues may occur. Note: Diazinon will not control organophosphate- resistant leaf- miners.

Mushroom Houses

Mushroom flies (Phorids and Sciarids)

Mix 4-5 lbs. of D·z·n diazinon 50W in 50 gallons of water and spray on outside and inside walls, floors and sideboards of mushroom houses after compost has been pasteurized by heating. Also spray over the plastic covering the beds and trays after spawning. **Note: Do not spray the compost. To avoid illegal residues, do not apply more than 3 applications to mushroom houses per crop.**

Seed Treatment - Planter Box Treatment

In the planter box, mix with seed as a dry mixture. Stir slightly using a mixing stick. **DO NOT MIX WITH BARE HANDS.** Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seed beyond planting time. Exposed treated seeds may be hazardous to birds and other wildlife. Dispose of excess treated seed by burial away from bodies of water.

- Notes:** (1) If seed has not been treated with a fungicide, use an EPA-registered fungicide in combination with D·z·n diazinon 50W at rates indicated on the fungicide package. Failure to use a fungicide may result in injury to the seed.
(2) If the seed treatment increases the friction between seeds sufficiently to reduce the rate of delivery through the planter, 1/2 oz. of graphite per bushel seed should be added as a drill lubricant. Adjust the planter to give the required rate of seed delivery with the treated seed.
(3) Treated seed must not be used for or mixed with food or animal feed or processed for oil.

Corn

Seedcorn Beetle: Mix 3 oz. per bushel of seed.

Corn, Succulent Peas, and Succulent Beans

Seedcorn Maggot: Mix 1/2 oz. per bushel of seed.

Field Crops

REI - 24 hours.

Apply soil and foliar applications of D·z·n diazinon 50W in sufficient water to ensure thorough coverage for foliar applications. Observe stated time intervals between last application and harvest, as well as intervals between

applications and total number of applications that can be made per season:

Pre-Harvest Intervals (PHI*) For Foliar Applications

<u>Crop</u>	<u>PHI*</u>
Hops	14
Sugar Beets	14

*PHI = Pre-Harvest Interval or the number of days between the last application and harvest.

Crop	Pest	Rate	Timing/Special Directions
Hops	Aphids. Spider Mites	2 lbs. per acre	Apply as pests occur. Repeat applications, if necessary in 14 days. Do not make more than 4 applications per season or apply within 14 days of harvest or illegal residues may occur.
Sugar Beets	Wireworms	6-8 lbs. per acre	Broadcast just before planting and immediately incorporate into the top 4-6 inches of soil. Do not apply more than 8 lbs. of this product to the soil, per season.
	Beet Leafhoppers, Dipterous Leafminers, Aphids	3/4-1 lb. per acre	Apply as insects occur. Repeat application, as necessary, up to 5 times per season with a minimum of 7 days between applications. Do not apply within 14 days of harvest. Note: Diazinon will not control organophosphate-resistant leafminers.

11/25

Crop	Pest	Rate	Timing/Special Directions
Sugar Beets (Cont'd)	Grasshoppers	1 lb. per acre	Apply as insects occur. Repeat application, as necessary, up to 5 times per season with a minimum of 7 days between applications. Do not apply within 14 days of harvest.

Note: Tops may be fed to beef and dairy animals.

Grassland Insects

Ditch Banks, Roadsides, Wasteland, Noncrop Areas, Barrier Strips

Grasshoppers: Apply 3/4-1 lb. per acre in sufficient water or oil to obtain thorough coverage when insects first appear, preferably in the nymphal stage.

Do not graze or use treated grass for food or feed.

Livestock Insects

Sheep "Ticks" (Keds), Lice

) High gallonage, high pressure spray: Mix 1/2 lb. in 100 gals. of water, and apply at the rate of 1 gal. per animal.

Low gallonage, low pressure spray: Mix 1 lb. in 100 gals. of water. Apply at the rate of 1 qt. per animal.

Sprinkler Can: Mix 1 oz. in 6 gals. of water. Crowd sheep into a small pen and sprinkle 1 qt. of this mixture over the head, neck, top, and side of each sheep. Stir or agitate mixture frequently.

Note: Do not slaughter sheep within 14 days after treatment. Repeat applications as necessary.

Fly Control in Livestock Structures

) Fly control in barns and animal sleeping quarters (except dairy barns, milk rooms, and poultry houses).

Residual Sprays: Remove animals from buildings or corrals prior to treatment and keep animals out for at least 4 hours. Mix 2 lbs. of D·z·n diazinon 50W in 25 gals. of water and apply as an overall spray to ceilings and walls of livestock sheds, calf barns, hog barns, loafing sheds, and other farm buildings except poultry houses, dairy barns, and milk rooms. Areas where flies congregate, such as garbage dumps and corrals, should be sprayed thoroughly. Longer residual control may be obtained by increasing the rate to 4 lbs. in 25 gals. of water. Applications should be made to point of runoff. One gal. of the above spray should cover from 350-750 sq. ft. of area, depending on the surface treated. Repeat applications as fly populations warrant.

Bait Sprays: Mix the appropriate amount of D·z·n diazinon 50W with 1 lb. of sugar (or 2 cups of syrup or molasses) for the desired application.

For knapsack or similar type sprayer, mix 1/2 lb. with the sugar in 2 1/2 gals. of water and spot treat areas frequented by flies including cracks, crevices, doorways, around windows, or other areas where flies congregate. Repeat applications as fly populations warrant.

For sprinkling can application, mix 2 oz. with the sugar in 5 gals. of water and sprinkle lightly over the floor in areas frequented by flies. Applications every day or two will generally be required.

Do not contaminate milk, feed, drinking water, or apply to animals or poultry. Do not use in edible product areas of food processing plants or other areas where food is commercially prepared or processed. Do not use in dwellings.

Insect Control on Ornamentals

REI - 12 hours.

To control certain insects on Arborvitae, Azalea, Birch, Boxwood, Camellia, Carnation, Chrysanthemum, Crab Apple, Dogwood, Douglas Fir, Elm, Euonymus, Gladioli, Hawthorn, Holly, Honey Locust, Honeysuckle, Juniper, Lilac, Lilly, Locust, Maple, Marigold, Oak, Pansy, Petunia, Pine, Ornamental Plum, Poplar, Rhododendron, Rose, Scotch Pine, Spirea, Spruce, Sycamore, Wax Plant (Hoya), Willow, Yellowwood and Yew, apply the recommended rates indicated below. Try to spray underside of leaves and penetrate dense foliage.

Insects	Amount of D·z·n diazinon 50W to Use	
	Rate per 3 gals. water	Rate per 100 gals. water
Aphids, Bagworms, Carnation Bud Mites, Carnation Shoot Mites, Clover Mites, Cyclamen Mites, Dipterous Leafminers, European Pine Shoot Moths, European Red Mites, Flea Beetles, Holly Bud Moths, Leafhoppers, Obscure Root Weevils, Omnivorous Leaf-tiers, Privet Mites, Scale Crawlers (Cottonycushion, Lecanium, Pine Needle, San José, Soft), Thrips, Two-Spotted Spider Mites, Juniper Webworms, Whiteflies	1/2 oz.	1 lb.
Apple-and-Thorn Skeletonizers, Cotoneaster Webworms, European Pine Sawfly, Fall Webworms, Hemlock Chermes, Oak Loopers, Obliquebanded Leafrollers, Pear Slugs, Tent Caterpillars	1 1/2 oz.	3 lbs.

Note: Do not use on Ferns, Poinsettia, Hibiscus, Papaya, Pilea, and Gardenia, since injury to the plants may occur.

Mimosa Webworms: Mix 1/2 oz. in 3 gals. of water (or 1 lb. in 100 gals. of water) and apply as a thorough foliar spray when adults first appear. Make a second and third application at 4-5 week intervals.

Lawn Pest Control (Excluding Sod Farms and Golf Courses)

Precautions: Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds, especially waterfowl. Avoid overlapping sprays. If waterfowl, i.e., ducks or geese, can be expected to enter treated area after treatment (except spot treatments), water lawn with at least 1/4 inch of water immediately after application, however, stop irrigation before puddling occurs.

To control pests listed below, apply specified rates.

Pests	Rate/1,000 sq. ft.		Remarks
	D·z·n diazinon 50W	Water	
Lawn Chinch Bugs	2-3 oz.	3 gals.	Use higher rate for longer residual control and in turfgrass of dense growth, such as St. Augustine.
Ants, Armyworms, Bermudagrass Mites, Clover Mites, Springtails (Collembola), Crickets, Cutworms, Digger Wasps, Earwigs, Frit Flies, Lawn Billbugs, Leafhoppers, Sod Webworms (Lawn Moths), Sowbugs (Pillbugs), Brown Dog Ticks, Chiggers, Fleas	3 oz.	3 gals.	Spot spray ant hills and wasp ground nest openings. For digger wasps, apply product at dusk when wasps are not active. For frit flies, mow grass and water well before treatment, but delay application until grass is dry. Do not apply to animals. For brown dog ticks, spray grass and under shrubbery.
Hyperodes Weevils	3 oz.	3 gals.	Treat problem areas in mid-April and again in mid-May.
Millipedes	3 oz.	3 gals.	
Rhodesgrass Scales	3 oz.	3 gals.	Apply when crawlers first emerge.
White Grubs (Japanese Beetle Larvae, European Chafer, Southern Chafer)	3 oz.	3 gals.	Apply any time between late July and early October.

Nuisance Pests in Outside Areas

Pests	Amount of D·z·n diazinon 50W	Water	Remarks
Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Milli- pedes, Sowbugs (Pillbugs), Spiders, Springtails (Collembola)	3 oz.	3 gals.	<p>Spray window and door screens, sills, foundations, porch and patio walls, entranceways, walks, garbage cans, tree trunks, and into cracks and other places where insects can hide.</p> <p>To prevent these pests from entering the house, spray a 5-foot band of ground around the house next to the foundation of the house to a height of 2 to 3 ft. Repeat application as necessary.</p> <p>Note: Do not apply to animals. Do not apply to plants (except tree trunks as noted above) at this nuisance pest rate.</p>

Pests	Amount of D·z·n diazinon 50W	Water	Remarks
Fire Ants	2 teaspoons	None	Evenly apply this product in its dry, undiluted state over each fire ant mound.
	1 oz.	5 gals.	Apply 1 gal. of solution to each mound by sprinkling the mound and the immediate surrounding area until wet. Apply gently to avoid distributing ants.
			Note: For best results, apply in cool weather (60-80°F), or in early morning or late afternoon when the ants are most active. Treat new mounds as they appear.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If pesticide, spray mixture or rinsates cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. Stay out of smoke from burning container.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Do not breathe dust or spray mist. Avoid contact with skin, eyes, and clothing. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans or pets. Do not contaminate ornamental fish ponds. Be sure to read and follow "Work Safety Rules" appearing in this booklet.

Statement of Practical Treatment

This product is an organophosphate.

If swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If inhaled: Remove victim to fresh air. Apply artificial respiration if indicated. Get medical attention.

If on skin: Remove contaminated clothing and wash affected areas thoroughly with soap and water. Get medical attention.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation develops or persists.

Note to Physician: If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants

- Shoes plus socks
- Waterproof gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. On lawns, if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply a minimum of 1/4 inch of water immediately after applying this product. Stop irrigation before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

D.z.n® trademark of Ciba-Geigy Corporation

©1995 Ciba-Geigy Corporation

Ciba Crop Protection
Ciba-Geigy Corporation
Greensboro, North Carolina 27419

CGA 1L66

June 2, 1982
Revised Nov. 30, 1982 (Include
EPA comments of 11/22/82)

Revised Jan. 02, 1985 (EPA PR
Notices 83-2 & 83-3)

Revised April 1986 - reduced
rates, added precautions re:
avian toxicity

Revised April 5, 1988 - added
chemigation prohibition,
deleted dust formulation
instructions, deleted golf
course/sod farm dfu, deleted
golf course/sod farm
precautions in Env. Hazards,
added do not use on golf
courses/sod farms state-
ment to front panel.

Revised Sept. 21, 1988
Deleted non-supported crops
until May 1, 1987, EPA Data
Call-In; added spill
statement; added Chinese

broccoli, bok choy, and
Chinese mustard.

Revised Nov. 11, 1988
Deleted mushroom houses,
changed all references to Corn
to "Sweet Corn," deleted corn
from aerial application
statements, deleted
bermudagrass and rangeland per
EPA comments. Added "Do not
use on commercially grown
potatoes which will be hand-
harvested.

Accepted - 11/23/88 with
comments

Revised 6/7/89 - moved golf
course/sod farm prohibition
under product name, revised
Hazards to Humans and Domest.
Animals, revised precautionary
statements, revised Env.
Hazards section, added
"Recommended for
Agricultural or Commercial
Use Only, revised Note to
Physician, revised all
directions for use to reflect
Data Call-In Residue Program,
added Chinese radish, revised
reentry statements - 24 hrs.
vegs., fruits, field crops.

Revised 9/15/89 - Additional
minor revisions per EPA letter
of 6/23/89 and Reg.
Standard Changes.

Revised 11/21/89 per EPA
letter of 11/6/89 deleted in-
furrow treatment for root
vegs.

Revised 12/21/89 - added
cranberry chemigation

Revised 1/11/90 - referred to
chemig. directions in dfu's
for cranberries and revised
Env. Hazards per EPA letter of
12/4/89.

Revised 10/8/90 correct min.
gallons for grapes and
strawberries and other
corrections.

Revised 11/20/90 per EPA
letter of 10/9/90.

Revised 1/9/92 per EPA letter
of 8/30/91 - change in signal
word to Caution, change in

precautionary text.

Revised 5/11/92 - deleted celery and modified stone fruit, pome fruit, almond, leafy vegs, brassica dfus because of over tolerance residues.

Revised 1/18/93 - editorial changes per EPA letter of 11/16/92

Revised 8/20/93 - PR Notice WPS label changes, logo

Revised 2/2/94 - Added code 1L66M 024 per EPA acceptance of 1/25/94 change © date to 1994

Revised 7/6/94 - Added dfu's for fire ant control, changed to Ciba Crop Protection. EPA approved 9/2/94-CGA 1L66N 094

Revised 9/13/94 - Deleted pineapple refuse feeding restriction

Accepted 11/3/94 - Removal of pineapple refuse feeding restriction. CGA 1L66P 114

Revised 5/17/95 - added rutabagas, blueberries, changed min gals. of water to 5 gals. for cranberries, added cranberry tipworm, changed © date, added 15 ornamentals, deleted flammability statement and other changes per EPA letter of 5/20/95.

Revised 12/4/95 - revised cranberry dfu's per Cranberry Institute; change gallonage to 15 ground, 20 aerial

[GANNONC.LABELD\LBL-D-MS]DZN5BK-B - 12/5/95

PM 13 100-460

498426 P1725

DEC 12 1995

Ms. Carolyn Bussey
Ciba-Geigy Corporation
P.O. Box 18300
Greensboro, NC 27419-8300

Dear Ms. Bussey:

Subject: Amendments- Revise per Agency letter dated March 20, 1995,
Add Rutabagas, Add 25 Ornamentals and 1 pest,
Reinstate Blueberries including fire ants from 24C,
Revise Cranberry directions including aerial gallonage
D.z.n diazinon 50W
EPA Reg. No. 100-460
Your submission dated December 6, 1995

The amendments referred to above, submitted in connection with a registration under FIFRA section 3(c)(7)(a), are acceptable provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
 - a) For Apricots, add the missing horizontal line in the Pest, Rate, and Timing/Special Directions columns between the pests Olive Scale Crawlers and Apricot Mealybugs.
 - b) There are a number of crops for which the directions for control of Mole Crickets were not standardized on the revised label despite indication on your working copy that they should be. For the crops Brussels Sprouts, Cauliflower, Collards, Kale, Lettuce, Mustard, and Peppers revise the Timing/Special Directions to read as follows:

BEST COPY AVAILABLE

Apply to plant beds or fields 1-2 days before planting and immediately incorporate into the top 1-2 inches of soil.

- c) For Cabbage, delete the horizontal line in the Pest column under the name Root Maggots, since both dilutions/directions pertain to this same pest.
- d) Under the crop Sugar Beets, revise the Note to read:
Sugar beet tops may be fed to beef and dairy animals.
- e) In the second listing of pests for Ornamentals, revise "Sowfly" to "Sawfly".
- f) In the Remarks column for Ants, etc. in the Lawn Pest Control table, add "Note:" before "Do not apply to animals.".
- g) In the Nuisance Pests in Outside Areas directions, add the missing vertical lines between the Pests, Amount of D.z.n diazinon 50W, Water, and Remarks columns for both sets of pests and add the horizontal line across all columns at the end of each pest section, i.e., one under "nuisance pest rate" and one under "as they appear".
- h) For both the container label and the booklet, in the Hazards to Humans and Domestic Animals, revise "Do not breathe dust or spray mist." to "Avoid breathing dust or spray mist." Also, as noted in the Agency's March 20, 1995 letter, because a rate in teaspoons now appears in the Fire Ants directions, add "Food utensils such as teaspoons or measuring cups must not be used for food purposes after use in measuring pesticides." after "Avoid contamination of food and feed". On the container label, revise "Be sure to read and follow "Work Safety Rules" that appear in the enclosed booklet." with "Be sure to read and follow "Work Safety Rules" and Worker Protection precautions appearing on this label." In the booklet label, revise "Be sure to read and follow "Work Safety Rules" appearing in this booklet." with "Be sure to read and follow "Work Safety Rules" and Worker Protection precautions appearing on this label."
- i) For the container label, add the Personal Protective Equipment, Engineering Control Statements, and User Safety Recommendations headings and associated text to the Precautionary Statements, as they appear in the booklet label.

BEST COPY AVAILABLE

3/85

- j) In the Statement of Practical Treatment, revise the last sentence of the **If in Eyes** advice to read "Call a physician if irritation persists."

Sincerely,

George T. LaRocca
Product Manager (13)
Insecticide Rodenticide Branch
Registration Division (7505C)

BEST COPY AVAILABLE

RESTRICTED USE PESTICIDE
DUE TO AVIAN AND AQUATIC TOXICITY

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED
APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION,
AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED
APPLICATOR'S CERTIFICATION.

Container Label for Use With Booklet

D·z·n® diazinon 50W

INSECTICIDE

This product must not be used
on golf courses and sod farms.

For control of certain insects and mites on
fruits, nuts, vegetables, field crops, turfgrass,
and ornamentals.

Active Ingredient:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-
methyl-4-pyrimidinyl) phosphorothioate 50%

Inert Ingredients: 50%

Total: 100%

D·z·n diazinon 50W is a wettable powder.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements on left panel.

For directions for use and specific pests see booklet in bottom
flap of bag.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and
with the Worker Protection Standard, 40 CFR part 170.
Refer to supplemental labeling under "Agricultural Use
Requirements" in the Directions for Use section for
information about this standard.

ACCEPTED
with COMMENTS
in EPA Letter 100-460

DEC 12 1995

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for sale in the U.S.
registered under EPA Reg. No.

100-460

6/8

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

) The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Ciba-Geigy or the Seller. All such risks shall be assumed by the Buyer.

) Ciba-Geigy warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. Ciba-Geigy makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Ciba-Geigy or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. Ciba-Geigy and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of Ciba-Geigy.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

General Information

D·z·n diazinon 50W is a wettable powder which should be mixed with sufficient water to assure thorough coverage unless otherwise noted in the Directions for Use.

Best control is obtained when application is made at first sign of pests. Application should be repeated only as directed to maintain control.

Since D·z·n diazinon 50W is a wettable powder formulation, constant agitation is necessary during application.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, since crop or plant injury may result.

Do not use on food crops grown in greenhouses.

Work Safety Rules

REPEATED EXPOSURES TO CHOLINESTERASE INHIBITORS SUCH AS ARE CONTAINED IN THIS PRODUCT MAY, WITHOUT WARNING, CAUSE PROLONGED SUSCEPTIBILITY TO VERY SMALL DOSES OF ANY CHOLINESTERASE INHIBITOR.

When handling D·z·n Diazinon 50W, do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. See Practical Treatment section of this label.

8/10

Chemigation

Refer to supplemental labeling booklet in bottom flap of bag for chemigation use directions for cranberries. Do not apply this product to cranberries through any type of irrigation system unless the specific directions for chemigation are followed.

Do not apply this product to any other crop or site through any type of irrigation system.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If pesticide, spray mixture, or rinsates cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. Stay out of smoke from burning container.

For minor spills, leaks, etc. follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

FOR SPECIFIC DIRECTIONS FOR USE, SEE BOOKLET IN BOTTOM FLAP OF BAG.

D.z.n diazinon 50W is registered for use in or on:

9185

Almonds (CA only)	Kale	Sheep
Apples	Lawns (Excludes	Spinach
Apricots	Golf Courses and	Squash
Beans, (Lima, Pole,	Sod Farms)	Strawberries
and Snap)	Lettuce	Sugar Beets
(succulent only)	Loganberries (CA,	Sweet Potatoes
Beets	OR, WA only)	Sweet Corn
Blackberries (CA,	Melons	Swiss Chard
OR, WA only)	Mushroom Houses	Tomatoes
Blueberries	Mustard	Turnips
Boysenberries (CA,	Nectarines	Turnip Tops
OR, WA only)	Non-crop Areas	Walnuts (CA only)
Broccoli	Onions	
Brussels Sprouts	Ornamentals	
Cabbage	Parsley	
Caneberries (CA,	Parsnips	
OR, WA only)	Peaches	
Carrots	Pears	
Cauliflower	Peas (succulent	
Cherries	only)	
Chinese Broccoli	Peppers	
Chinese Cabbage	Pineapple	
(Bok Choy and	Plums	
Napa)	Potatoes**	
Chinese Mustard	Prunes	
Collards	Radishes	
Cranberries	Raspberries (CA, OR	
Cucumbers	WA only)	
Dewberries (CA,	Rutabagas	
OR, WA only)	Seed Treatment,	
Endive (Escarole)	Planter Box	
Fly control in	(Corn and	
barns and animal	Succulent Peas	
sleeping quarters*	and Beans)	
Grapes		
Hops		

*except dairy barns, milk rooms, and poultry houses

**except commercially-grown potatoes which will be hand-harvested

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Do not breathe dust or spray mist. Avoid contact with skin, eyes, and clothing. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans or pets. Do not contaminate ornamental fish ponds. Be sure to read and follow the "Work Safety Rules" that appear in the enclosed booklet.

Statement of Practical Treatment

This product is an organophosphate.

If swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If inhaled: Remove victim to fresh air. Apply artificial respiration if indicated. Get medical attention.

If on skin: Remove contaminated clothing and wash affected areas thoroughly with soap and water. Get medical attention.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation develops or persists.

Note to Physician: If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid

overlapping of sprays. On lawns, if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply a minimum of 1/4 inch of water immediately after applying this product. Stop irrigation before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

D·z·n® trademark of Ciba-Geigy Corporation

©1995 Ciba-Geigy Corporation

Ciba Crop Protection
 Ciba-Geigy Corporation
 Greensboro, North Carolina 27419

CGA 1L118 (5 Pounds)
 CGA 1L117 (4 Pounds)

[GANNONC.LABELD]DZN50CLA

Revised June 2, 1982
 Revised Nov. 30, 1982
 (Including EPA comments
 of 11/22/82)
 Revised Jan. 02, 1985
 (EPA PR Notices 83-2 &
 83-3)
 Revised March 3, 1987
 Reduced rates on turf,
 separate sections for golf
 courses and lawns, revised
Environmental Hazards
 Revised April 5, 1988
 Deleted golf courses, sod
 farms, added do not use on
 golf courses, sod farms,
 deleted ref. to golf courses

12/85

and sod farms from Env. Hazards, added chemigation prohibition.

Revised November 22, 1988 - Deleted non-supported crops per EPA's May 1, 1988 Data Call-In, added spill statement

Accepted 11/23/88 with comments

Revised 2/28/89 per EPA comments and per Data Call-In - Moved golf course/sod farm prohibition under product name, changed a.i. name back to original accepted version, revised Hazards to Humans and Domestic Animals, Revised precautionary statements, revised Env. Hazards, added "Recommended for Agricultural or Commercial Use only, revised note to physician, revised list of labeled crops, did not include apricots, plums or prune dfu's on container label because of lack of space.

Revised 6/7/89 - revised reentry statements - 24 hr. reentry for vegs., fruits, field crops, see booklet.

Revised 9/15/89 - minor revisions per EPA letter of 6/23/89 and Registration Standard.

Revised 11/21/89 - per EPA letter of 11/6/89

Revised 1/11/90 - referral statement for chemigation - cranberries and revised Env. Hazards per EPA letter of 12/4/89, deleted blueberries.

Revised 10/8/90 - amended "Failure to Follow" statement, revised Worker Safety lang. to match booklet.

Revised 11/20/90 per EPA letter of 10/6/90.

Revised 1/16/91 per EPA letter of 12/18/90

Revised 10/3/91 per EPA letters of 12/18/90 and

12/85

8/26/91 (signal word)
Revised 5/11/92 - deleted
celery (over tolerance) and
aerial application limita-
tions.
Revised 1/18/93 - editorial
changes per EPA letter of
11/16/92
Revised 8/25/93 - WPS & logo
changes
Revised 2/2/94 - WPS accep-
tance per EPA letter of
1/25/94 - CGA 1L118G 024 and
1L117F 024 codes assigned,
changed ● date
Revised 5/17/95 - deleted
flammability statements,
added rutabagas, blueberries,
changed ● date

[GANNONC.LABELD]DZN50CLA - 5/19/95

11/2/95

SECOND DRAFT
CROP/ORNAMENTALS, CRANBERRY DFU'S REVISED

RESTRICTED USE PESTICIDE
DUE TO AVIAN AND AQUATIC TOXICITY

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED
APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION
AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED
APPLICATOR'S CERTIFICATION.

Booklet for 4 and 5 Lbs.

D·z·n® diazinon 50W

INSECTICIDE

For control of certain insects and mites on
fruits, nuts, vegetables, field crops, lawns,
and ornamentals. For specific pests,
refer to use directions.

This product must not be used on
golf courses and sod farms.

Active Ingredient:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-	
methyl-4-pyrimidinyl) phosphorothioate	50%
Inert Ingredients:	50%
Total:	100%

D·z·n diazinon 50W is a wettable powder

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements
and directions for use inside booklet.

EPA Reg. No. 100-460

CGA 1L66

[GANNONC.LABELD\LBL-D-MS]DZN5BK-B - 12/5/95

ACCEPTED
with COMMENTS
in EPA Letter Dated

DEC 12 1995

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

100-460

.....
.....
.....
.....
.....

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire **Directions for Use** and the **Conditions of Sale and Warranty** before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

The **Directions for Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Ciba-Geigy or the Seller. All such risks shall be assumed by the Buyer.

Ciba-Geigy warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions for Use** subject to the inherent risks referred to above. **Ciba-Geigy makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Ciba-Geigy or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product.** Ciba-Geigy and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing **Conditions of Sale and Warranty**, which may be varied only by agreement in writing signed by a duly authorized representative of Ciba-Geigy.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval. Exception: If the product is soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

1785

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow entry into treated areas until sprays have dried. Do not permit children or pets to go onto sprayed grass until spray has completely dried.

Note: The California Department of Agriculture has set a 5-day reentry period for diazinon on grapes, peaches, and nectarines.

Five days should elapse between the time of application and the time a worker may enter the field to engage in any activity requiring substantial contact with treated foliage. When a mixture of two or more organophosphate pesticides are applied in combination, the interval should be prolonged by adding to the largest applicable interval an additional 50% of that interval.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

GENERAL INFORMATION

D·z·n® diazinon 50W is a wettable powder which should be mixed with sufficient water to assure thorough coverage unless otherwise noted in the **Directions for Use**.

Best control is obtained when application is made at first sign of pests. Application should be repeated only as directed to maintain control.

Since D·z·n diazinon 50W is a wettable powder formulation, constant agitation is necessary during application.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, since crop or plant injury may result.

Do not use on food crops grown in greenhouses.

Work Safety Rules

REPEATED EXPOSURES TO CHOLINESTERASE INHIBITORS SUCH AS ARE CONTAINED IN THIS PRODUCT MAY, WITHOUT WARNING, CAUSE PROLONGED SUSCEPTIBILITY TO VERY SMALL DOSES OF ANY CHOLINESTERASE INHIBITOR.

When handling D·z·n diazinon 50W, do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. See **Practical Treatment** section of this label.

CHEMIGATION

APPLICATION THROUGH AN IRRIGATION SYSTEM (CHEMIGATION) - CRANBERRIES ONLY

D·z·n diazinon 50W may be applied alone or in combination with other pesticides registered for application through sprinkler irrigation systems to cranberries. To insure compatibility, pour the products into a small container of water in the correct proportions. After thorough mixing, let stand for five minutes. If the combination remains mixed, or can be remixed readily, the mixture is compatible.

Do not apply this product through any type of irrigation system to other crops. For chemigation, apply this product to cranberries only through sprinkler irrigation systems. Do not apply this product to cranberries through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

Do not connect the irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments, if the need arises

Chemigation System Connected to Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction.

19/80

There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

Sprinkler Chemigation

1. The system must contain a functional check-valve, a vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Operating Instructions

1. Determine the acreage to be treated by the sprinkler system.
2. Measure the appropriate amount of D·z·n diazinon 50W needed to treat the acreage to be treated. Refer to the specific directions for use for cranberries for the application rate and the amount of water per acre.
3. Add the pre-measured D·z·n diazinon 50W to the chemigation system supply tank and dilute with water if needed. Constant agitation is required to keep the

product in solution when D-z-n diazinon 50W is used alone or tank mixed with other pesticides registered for sprinkler chemigation application to cranberries.

4. Start the irrigation system and bring the system up to full pressure, making certain that all sprinkler heads are functioning properly.
5. Engage the chemigation injection or venturi system to add the diazinon to the irrigation lines.
6. Continue to run irrigation system for a few minutes after supply tank is empty to allow material in the irrigation lines to reach the sprinkler heads furthest from the chemigation injection point.

Fruit and Nut Crops

REI - 24 hours.

D-z-n diazinon 50W may be applied using ground or aerial application equipment as specified in the table below. This table also indicates the **minimum** amount of water that can be used for the various methods of application. The rate of D-z-n diazinon 50W for aerial spray or concentrated spray applied by ground equipment is based on a dilute full cover spray applied to fruit or nut trees with conventional ground equipment at a rate of 300-400 gals. of water per acre. The specific amount of product to be used appears under the separate directions for use for each crop. (**Note:** "lbs." or "lbs. of product" in the directions means pounds of this product, not pounds of active ingredient). If the "Rate" calls for 1 lb. of D-z-n diazinon 50W per 100 gallons of water, this amounts to 3-4 lbs. of product per acre when applied to fruit trees at usual dilute ground spray volumes of 300-400 gals. of water per acre. In no case should the amount of product used per acre exceed the maximum amount per acre per application that is specified in the directions for use for individual crops.

<u>Crop</u>	<u>Aerial</u> <u>Application</u>	<u>Ground Application</u>	
	<u>Minimum</u> <u>Gals./</u> <u>Acre</u>	<u>Dilute</u>	<u>Minimum</u> <u>Gals./Acre</u> <u>Concentrate</u>
Almonds*	20	100	20
Apples	20	100	20
Apricots	20	100	20
Blackberries**	20	100	20
Blueberries	20	100	20
Boysenberries**	20	100	20
Cherries	20	100	20
Cranberries	20	15	--
Dewberries**	20	100	20
Grapes	20	100	20
Loganberries**	20	100	20
Nectarines	20	100	20
Peaches	20	100	20
Pears	20	100	20
Pineapple	20	200	--
Plums	20	100	20
Prunes	20	100	20
Raspberries**	20	100	20
Strawberries	20	100	20
Walnuts*	20	100	20

*CA only

**CA, OR, or WA only

Observe stated time intervals between last application and harvest, as well as intervals between applications and total number of applications that can be made per season.

Crop	PHI*
Almonds	**
Apples	21
Apricots	21
Blackberries	7
Blueberries	7
Boysenberries	7
Cherries	21
Cranberries	7
Dewberries	7
Grapes	28
Loganberries	7
Nectarines	21
Peaches	21
Pears	21
Pineapple	7
Plums, Prunes	21
Raspberries	7
Strawberries	5
Walnuts	45

*PHI = Pre-Harvest Interval or the number of days between the last application and harvest.

** = Dormant spray only.

Note: To protect bees, do not apply this product to fruit trees when the trees or substantial numbers of weed in the orchard are in bloom.

Crop	Pest	Rate	Timing/Special Directions
Almonds (CA only)	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite Eggs, Brown Mite Eggs, Twig Borers, Apple Aphid Eggs, Black Cherry Aphid Eggs, Mealy Plum Aphid Eggs, Mealybugs	1-1 1/2 lbs. + 2-3 gals. dormant oil or 1-1 1/2 gals. superior type oil in 100 gals. of water	Apply as a single dormant spray. Do not apply more than 6 lbs. of product or 9 gals. of oil per acre.
Apples	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite Eggs, Brown Mite Eggs, Twig Borers, Apple Aphid Eggs, Black Cherry Aphid Eggs, Mealy Plum Aphid Eggs, Mealybugs	1 lb. + 2 gals. dormant oil or 1-1 1/2 gals. superior type oil in 100 gals. of water	Apply as a dormant or delayed dormant spray. Do not apply more than 4 lbs. of product or 6 gals. of oil per acre.
	Leafhoppers	1 lb. in 100 gals. of water	Apply, beginning at petal fall, as infestations develop. Repeat application in 14 days if necessary.
	Eyespotted Bud Moths, Fruit Tree Leafrollers, Mealybugs	1 lb. in 100 gals. of water	Make 3 cover sprays, beginning at pink stage. Do not repeat application closer than 14 days.

2/1/25

Crop	Pest	Rate	Timing/Special Directions
Apples (Cont'd)	Codling Moths, Rosy Apple Aphids, Green Apple Aphids, Woolly Apple Aphids, Spider Mites	1 lb. in 100 gals. of water	Make 3 cover sprays beginning when infestations occur. Do not repeat applica- tions closer than 14 days. Note: Mites will be suppressed if diazinon is used in the regular cover spray program. Slight russeting may occur on some varieties of apples such as Golden Delicious.
	Apple Maggots	1 lb. in 100 gals. of water	Make 3 cover sprays when flies are active and laying eggs. Do not repeat applications closer than 14 days. Note: Diazinon may not control organophosphate- resistant apple maggots.
	San Jose Scale Crawlers, Forbes Scale Crawlers	1 lb. in 100 gals. of water	Apply early in the season when crawlers first appear. If infestation warrants, make a second applica- tion 14 days later.

Notes: (1) Do not apply more than 4 lbs. of product per acre per application and no more than 12 lbs. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.

Crop	Pest	Rate	Timing/Special Directions
Apricots	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite Eggs, Brown Mite Eggs, Twig Borers, Apple Aphid Eggs, Black Cherry Aphid Eggs, Mealy Plum Aphid Eggs, Mealybugs	1 lb. + 2 gals. dormant oil or 1-1 1/2 gals. superior type oil in 100 gals. of water	Apply as a dormant spray. Do not apply more than 4 lbs. of product or 6 gals. of oil per acre.
	Aphids, Brown Mites (Clover Mites), Two-Spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers	1 lb. in 100 gals. of water	Apply as infestations occur. Repeat application after 14 days if necessary.
	Olive Scale Crawlers Apricot Mealybugs	1/2 lb. + 1 1/2 gals. of light medium horticultural oil in 100 gals. of water 1 lb. in 100 gals. of water	Apply when scale crawlers are present. Do not apply more than 4 lbs. of product or 6 gals. of oil per acre per application. Apply as a cover spray from petal fall to June. Allow 14 days between applications.

Notes: (1) Do not apply more than 4 lbs. of product per acre per application and no more than 12 lbs. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.

Crop	Pest	Rate	Timing/Special Directions
Blue-berries	Cranberry fruitworms Cherry fruitworms, Blueberry maggots, Aphids, Thrips, Two-spotted spider mites	1 lb. per 100 gals. of water	Apply as infestations occur. Allow 14 days between applications.
	Fire ants	1 lb. in 100 gals. of water	Slowly apply 1 gal. of dilute mixture to each fire ant mound.

Notes: (1) Allow a minimum of 7 days between the last application and harvest. (2) Do not apply more than 2 pounds of product per acre per season.

Crop	Pest	Rate	Timing/Special Directions
Caneberries (CA, OR, and WA only) (Black- berries, Boysen- berries, Dew- berries, Logan- berries, Rasp- berries	Leafhoppers, Thrips, Aphids, Two-Spotted Spider Mites, Raspberry Fruitworms	2 lbs. per acre in a minimum of 100 gals. of water (200 gals. of water maximum)	Apply as insects occur. Repeat application after 14 days if neces- sary. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal resi- dues may occur. For raspberry fruitworm make one application when blossom buds separate. Make a second applica- tion immediately before blossoms open.
	Dryberry Mites	2-4 lbs. per acre in a mini- mum of 100 gals. of water	Apply as a dor- mant spray in the early Spring, when canes are still on the ground and buds are closed. Make a second applica- tion when buds are well devel- oped but still closed.
	Raspberry Crown Borers (Raspberry Root Borers)	4 lbs. per acre in a minimum of 100 gals. of water	Apply as a drench to the crown and lower canes in the Spring before buds break.

Crop	Pest	Rate	Timing/Special Directions
Cherries	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite Eggs, Brown Mite Eggs, Twig Borers, Apple Aphid Eggs, Black Cherry Aphid Eggs, Mealy Plum Aphid Eggs, Mealybugs	1 1/4 lbs. + 3 gals. dormant oil or 1-1 1/2 gals. superior type oil in 100 gals. of water	Apply as a dormant spray. Do not apply more than 4 lbs. of product or 6 gals. of oil per acre.
	Eyespotted Bud Moths, Fruit Tree Leafrollers	1 lb. in 100 gals. of water	Apply as infestations occur beginning at petal fall. Repeat application after 7 days if necessary.
	Leafhoppers	1 lb. in 100 gals. of water	Make 3-4 applications beginning at petal fall and continuing through the growing season as infestations occur. Allow 7 days between applications.
	Cherry Fruitflies	1/2-1 lb. in 100 gals. of water	Make 3-4 applications beginning when adult flies begin to emerge and repeating at 10 day intervals.
	Black Cherry Aphids	1 lb. in 100 gals. of water	Apply as a cover spray when aphids appear. Repeat application after 10 days if necessary.

Crop	Pest	Rate	Timing/Special Directions
Cherries (Cont'd)	San Jose Scale Crawlers	1 lb. in 100 gals. of water	Apply during the pre-pink stage or when crawlers first appear.
	Cherry Rust Mites	1/2-1 lb. in 100 gals. of water	Apply as a single supplemental spray to trees after harvest.

Notes: (1) Do not apply more than 4 lbs. of product per acre per application and no more than 12 lbs. of product per acre per year. (2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.

Cran- berries	Blackheaded Fireworms (<i>Rhopobota naevana</i>)	4 lbs.	Apply to larval stage. Pheromone trap captures and sweep monitoring may be used to optimize treat- ment timing. Repeat applica- tion after 14 days if neces- sary. A maximum of 6 applications per season is permitted.
	Cranberry Tipworms	4 lbs.	Presence of dam- age is not an in- dicator of active infestation. For best results, target early- stage larvae. Repeat applica- tion after 14 days if neces- sary. A maximum of 6 applications per season is permitted.