



**ACCEPTED**  
SEP - 2 1997  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 100-460

**SUPPLEMENTAL LABELING**

**D-Z-N® DIAZINON 50W  
INSECTICIDE**

**EPA Reg. No. 100-460**

This product must not be used on golf courses and sod farms.

**FOR CONTROL OF INSECTS AND MITES IN BLUEBERRIES**

Active Ingredient:

Diazinon: O, O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate . . . . . 50.0%

Inert Ingredients: . . . . . 50.0%

Total: . . . . . 100.0%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

**Follow all applicable directions, restrictions, and precautions, including statements pertaining to the Worker Protection Standard, on the EPA-registered D-z-n diazinon 50W label.**

**This label must be in the possession of the user at the time of application.**

**General Information**

D-z-n diazinon 50W is an wettable powder which should be mixed with sufficient water to assure thorough coverage. Constant agitation is necessary during

application. In the directions for use which follow, best control is obtained when application is made at the first sign of pests. Application should be repeated only as directed to maintain control.

**Blueberries**

**REI - 24 hours**

D-z-n diazinon 50W may be applied using ground or aerial application equipment as specified in the following table. This table indicates the **minimum** amount of water that can be used for the various methods of application.

Crop	Aerial Application Minimum Gals./Acre	Ground Application Minimum Gals./Acre	
		Dilute	Concentrate
Blueberries	20	100	20

Observe stated time interval between last application and harvest (PHI) , as well as intervals between applications and total amount of product that can be applied per season.

**Pre-Harvest Interval (PHI)\***

Crop	PHI
Blueberries	7 days

\*PHI = Pre-harvest Interval or the number of days to wait between the last application and harvest

**Note:** To protect bees, do not apply when crop or substantial numbers of weeds in the treatment area are in bloom.

Crop	Pest	Rate	Timing/Special Directions
Blueberries	Cranberry fruitworms, Cherry fruitworms, Blueberry maggots, Aphids, Thrips, Two-spotted spider mites	2 lbs./A	Apply as infestations occur. Allow 14 days between applications.

**Notes:** (1) Do not apply more than 2 lbs. of product per acre per application and no more than 10 lbs. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 7 days between the last application and harvest.

D-z-n® trademark of Novartis  
©1997 Novartis

Novartis Crop Protection, Inc.  
Greensboro, North Carolina 27419

July 25, 1997  
revised per EPA  
letter of 7/22/97

