

100-461

11-8-72

1043

REVISED LABEL CLAIMS FOR
DIAZINON® AG500 - EPA REG. NO. 100-461


Lawn Insects

White grubs, such as Japanese beetle larva, European chafer and Southern chafer and billbugs - Apply 4 fl. oz. per 1000 sq. ft. of lawn. Water grass thoroughly after application. For grub control make application any time between late July and early October. For billbug control apply when activity is first observed or when chewed or brown grass is present, indicating damage from this insect.

Hyperodes weevil (New York Only) - Apply 3 fl. oz. per 1000 sq. ft. of grass. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

Agricultural Division
CIBA-GEIGY Corporation
Ardsley, New York 10502

August 25, 1972



100-461

6853

No-Quibble Guarantee
 If for any reason you are not satisfied with results after applying this product within a "reasonable" period, you may be asked to get your money back. No ifs, ands or conditions of purchase and no bill will ever be mailed should you wish to return it.

Lawn Insect Control

Lawn Insect Control

FOR USE ON GRASS OR DICHOONDRA LAWNS

controls

white grubs, such as—Japanese beetle, European chafer, Southern chafer
 sod webworms (lawn moth larvae) chinchbugs,
 (and others listed on back panel)

A PROBLEM-SOLVING PRODUCT BY SCOTT'S, THE LAWN PEOPLE

Lawn Insect Control

NET WEIGHT 30 LBS

6853

6853

CAUTION: KEEP OUT OF REACH OF CHILDREN.
 Harmful if swallowed. Avoid breathing of dust. Avoid contact with skin, eyes and clothing. Avoid contamination of feed and feedstuffs. Wash thoroughly after handling. May be absorbed through skin. As directed water lawn after application and allow to dry before permitting children and pets on treated area.
 Fish in waters treated with this product may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.
 Apply this product only as specified on this label. Do not reuse container. Destroy when empty.
 Active ingredient: 0,0-Diethyl 0-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate 2.00%
 Inert ingredients 98.00%
Total 100.00%
 EPA Reg No 338-81
 Distributed by O M Scott & Sons
 Product of USA
 Marysville, Ohio 43040

6853

Lawn Insect Control

Best Available copy

3 of 3

Treats 5,000 sq ft / \$12.95

Lawn Insect Control

TO
OPEN
BOLD
TAP
▶

Treats 5,000 sq ft / \$12.95

Lawn Insect Control

A PROBLEM-SOLVING PRODUCT BY SCOTT'S, THE LAWN PEOPLE

Controls these insects in lawns:
white grubs, such as—Japanese beetle,
European chafer, Southern chafer

army worms	cutworms
Bermuda moths	crickets
billbugs	fleas
chiggers	hedgehoppers
chinchbugs	ticks
clover moths	soil webworms
crickets	(lawn moth larvae)

When to use

For control of insect damage apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat at signs of renewed insect activity.

For grub control make an application anytime between late July and early October.

How to apply

Apply to dry foliage. Distribute over the entire lawn area. Water after application to remove granules from the grass blades.

For chinchbug control water sufficiently to carry material into the thatch layer of the turf.

For grub control water thoroughly to carry the material into the soil.

To ensure proper distribution, use a Scotts Spreader (Standard or Automatic) at setting 7 1/2

Precautions

As directed, water lawn after application and allow to dry before permitting use of treated area.