



# D·z·n diazinon 4E

## DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

**IMPORTANT** Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

### Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of CIBA-GEIGY or the Seller. All such risks shall be assumed by the Buyer.

CIBA-GEIGY warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. CIBA-GEIGY makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall CIBA-GEIGY or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. CIBA-GEIGY and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of CIBA-GEIGY.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with this labeling.

### General Information

D·z·n diazinon 4E is an emulsifiable concentrate and is intended for use by pest control operators. Apply the product after dilution with water according to the directions for use for the control of those pests specified on this label. D·z·n diazinon 4E is not intended for use in preparing of base sprays.

This product may be used indoors in households, apartments, and other structures including schools (except classrooms in use), hospitals (except in patient rooms while occupied), office buildings, museums, libraries, sports facilities, stores, warehouses, and food and feed handling establishments. Read and follow the "Precautions" listed below when applying this product in any of these indoor areas. This product is also approved for use in USDA Meat and Poultry Plants in food and feed areas of food/feed handling establishments. D·z·n diazinon 4E may be applied only as a crack and crevice treatment or limited spot treatment. See the specific directions for use for treatment of these areas.

D·z·n diazinon 4E may also be used outdoors only on home lawns and ornamentals around the home. See the specific directions for use for treatment of these areas.

### INDOOR USE

Indoors D·z·n diazinon 4E controls:

ants	carpet beetles	scorpions
brown dog ticks	cockroaches (waterbugs)	spiders

In households and in food and feed areas of food/feed handling establishments this product also aids in the control of the following exposed stages of pantry pests which come into contact with treated surfaces:

drugs ore beetle	flour beetle	rice weevil
cigarette beetle	Indian meal moth	sawtooth grain beetle

**Precautions:** (1) Do not use this product indoors as an overall surface spray or as a space spray. Take care not to introduce this product into the air during application. (2) Use only in well-ventilated areas. (3) Do not apply where children are likely to frequently contact treated surfaces. (4) Remove or cover fish bowls and remove pets before application. (5) Do not treat pets with this product. (6) Do not allow children or pets in treated areas until spray has dried. (7) Do not apply this product in patient rooms or in any room while occupied by the elderly or infirm. (8) Do not apply to classrooms when in use. (9) In institutions (including, but not limited to, office buildings, museums, libraries, and sports facilities) do not apply this product when occupants are present in the immediate area. (10) The solvent used in this formulation is a petroleum distillate which may stain certain plastic, rubber, and enamel materials such as tiles and floor coverings. Do not treat such materials. (11) Avoid excessive wetting of carpets, floor coverings or unfinished materials because damage may occur. (12) Care should be taken to avoid spotting of walls, wall coverings, and fabrics. (13) Do not apply directly to conduits, motor housings or electrical switch boxes. (14) Do not apply to areas, surfaces or utensils which will contact food or feed. (15) Exposed food should be covered or removed during application. (16) All food processing surfaces and utensils should be covered during application or thoroughly washed with soap and water before use. (17) If anything is sprayed during application, remove clothing immediately after spraying, wash affected body areas with soap and water, and launder clothing before wearing again.

### Mixing Instructions

Dilute D·z·n diazinon 4E with water to make a 1/2% to 1% spray as indicated below.

Gals. of finished spray	Amount D·z·n diazinon 4E to use for	
	1/2% spray	1% spray
1	1 1/4 fl. oz.	2 1/2 fl. oz.
2	2 1/2 fl. oz.	5 fl. oz.
3	3 3/4 fl. oz.	7 1/2 fl. oz.
10	12 1/2 fl. oz.	25 fl. oz. (1 3/4 pts.)
25	32 fl. oz. (2 pts.)	64 fl. oz. (4 pts.)

\*2 1/2 tablespoons = 1 1/4 fl. oz.

**Indoor Use - Except Food and Feed Handling Establishments, Households, Apartments, and Non-Food Areas of Offices, Schools (except classrooms in use), Libraries, Hospitals (except patient rooms while occupied), Museums and Sports Facilities (See "Food and Feed Handling Establishments" for special directions when applying this product in food handling or serving areas in non-residential buildings).**

**Cockroaches (Waterbugs), Ants, Silverfish:** Make spot applications of 1/2% coarse spray or apply with a paintbrush to baseboards, areas around water pipes, cracks and crevices, surfaces behind and beneath cabinets, refrigerators, sinks, stoves, and similar areas where these insects hide.

For heavy cockroach infestations, make the first application at a strength of 1% followed by monthly applications of 1/2% spray.

**Brown Dog Ticks:** Make thorough application of a 1/2% spray in infested areas around baseboards, window and door frames, wall cracks, sleeping quarters of household pets, and localized areas of floors and floor coverings. Do not use as an overall surface spray. Fresh bedding should be placed in animal quarters following treatment. Do not treat animals with this product.

**Carpet Beetles:** Apply a 1/2% spray in spot applications along baseboards and edges of carpeting, under carpeting and rugs, and furniture in closets and on shelving, and wherever these insects are seen or suspected.

**Spiders and Scorpions:** Apply a 1/2% spray around windows, doors, and along baseboards. Make spot applications to other areas over which these pests may crawl.

**Pantry Pests - Exposed Stages (Sawtoothed grain beetle, Flour beetle, Rice weevil, Cigarette beetle, Drugstore beetle, Indian meal moth):** Apply a 1/2% spray to shelves and cupboards where food containers suspected of being infested are stored. Remove all food and shelving paper before spraying. Do not replace the food packages until spray has dried. Cover shelves with shelving paper. This treatment is supplemental to good housekeeping, since it will have no effect on insects breeding within food packages. Infested food packages should be destroyed.

### Food and Feed Handling Establishments

Places other than private residences in which food or feed is being processed, prepared or served (includes restaurants, canneries, bakeries, food processing plants, mills, dairies, and USDA Meat and Poultry Plants). D·z·n diazinon 4E may be used in the food/feed areas and the non food/non-feed areas of these establishments. Apply D·z·n diazinon 4E either as a limited spot or crack and crevice treatment as directed below.

#### Food/Feed Areas of Food/Feed Handling Establishments

Include areas for receiving, serving, storage (dry, cold, frozen, raw), packaging (canning, bottling, and grinding), enclosed processing systems (mills, dairies, and systems for processing edible oils and syrups).

**APPLICATION OF THIS PRODUCT IN THE FOOD/FEE AREAS OF FOOD HANDLING ESTABLISHMENTS, OTHER THAN AS A CRACK AND CREVICE TREATMENT OR LIMITED SPOT TREATMENT, IS NOT PERMITTED.**

#### Limited Spot Treatment

Apply this product diluted to 1/2-1% as a coarse, low pressure fan spray around water pipes and door frames, to surfaces beneath cabinets, refrigerators, sinks, stoves, storage areas, and similar floor areas. Limit spot treatments to floor surfaces where pests have been seen or where they are suspected of hiding. Limit the sum of individual spot treatments to an area no larger than 20% of the floor surface area. Any individual spot treatment must not exceed 2 square feet.

#### Crack and Crevice Treatment

Apply this product diluted to 1/2-1% with equipment capable of delivering a pin stream of insecticide. Apply in small amounts directly into cracks and crevices in such areas as expansion joints, spaces between equipment and the floor, openings leading to voids, and hollow space in walls and equipment legs and bases.

**Non-Food/Non-Feed Areas of Food/Feed Handling Establishments:** Include garbage rooms, lavatories, floor drains, sewer entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, mop closets and storage areas (after canning or bottling).

# D·z·n· diazinon 4E

## Spot Treatment

Apply to spot treated areas only. Spray. Apply at a low pressure. Use spray as a spot treatment along baseboards, in areas such as window sills, beneath cabinets, surfaces behind and under sinks, lockers and pallets, and to storage areas where cockroaches, ants, silverfish and spiders hide.

## OUTDOOR USE

### Control of pests on lawns and on ornamentals around the home only

**Precautions:** (1) Do not use this product on ornamentals or on turf areas other than around the home. (2) Do not apply this product through any type of irrigation system. (3) This product is highly toxic to bees exposed to direct treatment or to residues on blooming plants. Do not apply this product or allow it to drift to blooming plants if bees are visiting the treatment area. (4) Birds, especially waterfowl, feeding or drinking on treated area may be killed. (5) Because of the migratory habits of certain Atlantic coast waterfowl, do not apply this product to lawns in Nassau County, New York between November 1 and May 20. (6) Do not exceed maximum permitted label rates since rates above those recommended significantly increase potential hazards to birds and waterfowl. (7) Avoid overlapping of sprays. (8) If waterfowl, i.e., ducks, geese, can be expected in the treated lawn after treatment (except spot treatment) water lawn with at least 1/4 inch of water immediately after spraying this product. However, stop watering before puddling occurs. (9) Wear a long-sleeved shirt and long-legged pants when spraying. Spray with the wind to your back. Do not spray on windy days. If clothing becomes wet from spray during use, remove clothing after spraying. Wash affected body areas thoroughly with soap and water, and launder clothing before wearing again.

**Pests of Ornamental Plants:** To control certain pests on Arborvitae, Azalea, Birch, Boxwood, Camellia, Carnation, Chrysanthemum, Douglas Fir, Elm, Gladiolus, Hawthorn, Holly, Juniper, Lilac, Locust, Maple, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Rose, Spruce and Willow, apply the recommended rates indicated in the following chart.

Do not use on ferns, Impatiens, Mimosa, Papaya, Pilea and Gardenia since injury to the plants may occur. Do not spray underside of leaves and penetrate dense foliage.

Pests	Amount D·z·n diazinon 4E to use	
	per 3 gals water	per 100 gals water
Aphids, Bagworms, Carnation bud mites, Carnation shoot mites, Clover mites, Cyclamen mites, Dipterous leafminers, European pine shoot moths, European red mites, Flea beetles, Holly bud moths, Leafhoppers, Obscure root weevils, Omnivorous leafminers, Priver mites, Scale crawlers, Cottony cushion, Lecanium, Pine needle scale, San Jose scale, Thrips, Two-spotted spider mites, Juniper Webworms, Whiteflies	1/2 oz	1 qt
Apple and thorn skeletonizers, Cottoncaster webworms, Fall webworms, Hemlock chermes, Oak leafhoppers, Obliquebanded leafrollers, Pear sawfly, Tent caterpillars	1/2 oz	1 qt

1 qt = 32 fl oz; 1 fl oz = 1.28 fl oz

**Mimosa Webworm:** Mix 1/2 oz (1 tablespoon) in 3 gals of water or 1 qt in 100 gals of water and apply as a thorough foliar spray when adults first appear. Make a second and third application at 4-5 week intervals.

## Home Lawn Pests

Pests	Rate per 1000 sq. ft.		Remarks
	D·z·n diazinon 4E	Water	
Lawns (not Sugg.)	2.3 fl. oz.	3 gals.	Use higher rate for longer retention; dual control and in lawns of dense growth such as St. Augustine lawns.
Ants, Armyworms, Bermudagrass mites, Clover mites, Springtails (Collembola), Crickets, Cutworms, Digger wasps, Earwigs, Fire flies, Lawn billbugs, Leafhoppers, Sod webworms (Lawn moth), Sowbugs (billbugs)	2.3 fl. oz.	3 gals.	Spot spray anthills and wasp ground nest openings. For digger wasps, apply at dusk when the wasps are not active. For fire flies, mow grass and water well before treatment but delay application until grass is dry.
Brown dog ticks, Chiggers, Fleas	1 fl. oz.	3 gals.	Note: Do not apply to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house.
Two-spotted weevils	3 fl. oz.	3 gals.	Treat problem areas in mid April and again in mid May.
Milipedes	1 fl. oz.	3 gals.	
Rhodesgrass scales	3 fl. oz.	3 gals.	Apply when crawlers first emerge.
White grubs (such as Japanese beetle larvae), European chafers, Southern chafers	3 fl. oz.	3 gals.	Apply any time between late July and early October.

## Nuisance Pests Outside the Home

Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Fire Flies, Millipedes, Sowbugs (Billbugs), Spiders, and Springtails (Collembola).

Mix 3 1/2 oz in 3 gals of water.

Thoroughly spray window and door screens and sills, foundations, porch and patio walls, entranceways, walks, garbage cans, tree trunks and into cracks and other places where insects can hide. Repeat application as necessary.

To prevent entry of insects into the house, spray a five-foot band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 feet. Repeat application as necessary.

## Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

**Pesticide:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**Container:** Triple rinse or equivalent. Then offer for recycling, reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-888-8372 day or night.

Diazinon trademark of CIBA-GEIGY Corporation.

© 1992 CIBA-GEIGY Corporation

Agricultural Division  
CIBA-GEIGY Corporation  
Greensboro, North Carolina 27412  
CGA 11474A 062

BEST AVAILABLE COPY