

D·z·n®
Granular

Treats up to 2500 square feet.

lawn
insect
control

This product must not be used
on golf courses and sod farms.

Controls:

- Brown Dog Ticks
- Chinch bugs
- Sod webworms
- Lawn moths
- Earwigs
- Fleas
- Fire ants*
- White grubs of
 - . Japanese beetle
 - . European chafer
 - . Southern chafer
- Billbugs
- and many others

*Aids in control.

(See complete list on
back of bag)

ACCEPTED
OCT 12 1989
 Under the Federal Insecticide,
 Fungicide and Rodenticide Act
 as amended, for the Pesticide
 registered under
 EPA Reg. No. 100-468

Active Ingredient:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	2%
Inert Ingredients:	98%
Total:	100%


CAUTION

KEEP OUT OF REACH OF CHILDREN.

See additional precautionary statements
on back of bag.

12 1/2 Lbs.
Net Weight

Ready-to-use. Contains 2% diazinon
insecticide.



DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of CIBA-GEIGY or the Seller. All such risks shall be assumed by the Buyer.

CIBA-GEIGY warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. CIBA-GEIGY makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall CIBA-GEIGY or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. CIBA-GEIGY and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of CIBA-GEIGY.

D·z·n kills:

Ants	Crickets	Springtails
Armyworms	Cutworms	Sod webworms
Bermudagrass mites	Digger wasps	Sowbugs
Billbugs	Earwigs	White grubs of:
Brown dog ticks	Fire ants*	. Japanese beetle
Chiggers	Fleas	. European chafer
Chinch bugs	Lawn moths	. Southern chafer
Clover mites	Leafhoppers	
	Millipedes	

*Aids in control.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

OVERALL LAWN APPLICATION

D·z·n® Granules control lawn pests on both grass and dichondra lawns. Apply D·z·n Granules to your lawn with a fertilizer spreader at the rate shown on side panel chart. For application with other types of equipment, apply at the rate of 5 lbs. per 1000 sq. ft. (an area 20' by 50') or 8 oz. per 100 sq. ft. (an area 10' by 10').

Precautions: Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds. Avoid overlapping treated areas. If waterfowl, i.e., ducks, geese can be expected to enter the treated area after treatment (except spot treatments), water lawn with at least 0.25 inch water immediately after application, however, stop watering before puddling occurs.

Chinchbugs, Earwigs, Ants, Lawn Moths (Sod Webworms), Bermudagrass Mites, Crickets, Armyworms, Brown Dog Ticks, Chiggers, Clover Mites, Cutworms, Fleas, Leafhoppers, Sowbugs, Millipedes, Springtails (Collembola). Application should be made when grass is dry. Water lawn lightly after application. Repeat application in 7-10 days if necessary.

Billbugs; White grubs of Japanese beetle, European chafer, Southern chafer, Ataenius spretulus (dung beetle). Water grass thoroughly after application. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in Spring or early Summer when damage is first noticed. Water grass thoroughly with 1/4-1/2 in. of water after application. Treat again in late summer or early fall if a second generation occurs.


Hyperodes Weevils

Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

NOTE: The spreader settings listed below are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader setting if necessary in order to assure correct rate of application.

SPREADER SETTING NO.

Spreader Make and Model	For insects listed Grass and Dichondra Lawns 5 lbs./1,000 sq. ft. of lawn
Central Imperial Mark No. 1662-G	7 1/2
Central Suburbia Lawn Spreader - No. 1622-0	7 1/2
Cyclone Model B1 Rotary	6
Greenfield Model 105	8
J. C. Penney Model 6005	8 1/2
J. C. Penney Rotary Spreader	8
Scott Model 35	6 1/2
Sears Craftsman Model 671.19198	7 1/2



SPOT TREATMENT

Ants: Apply one tablespoon of D·z·n Granules over and around each individual ant hill.

Fire Ants: To aid in the control of fire ants in lawns and other recreational areas, apply 1 1/4 cups over, and 2 ft. around, each fire ant mound. Immediately after application, water in thoroughly. A minimum of 1 gallon of water should be applied to each mound area. Reapply as necessary. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness.

Note: For best results, apply in cool weather, 65-80°F, or in early morning or late evening hours. Treat new mounds as they appear.

Digger wasps: Apply one tablespoon of D·z·n Granules in and around each individual ground nest opening. Make the application about dusk when the wasps are not active.

BAND TREATMENT AROUND FOUNDATION

The following insects are commonly found around or near the foundation of houses as well as in the lawn.

Ants	Crickets	Millipedes
Brown dog ticks	Earwigs	Sowbugs
Clover mites	Fleas	Springtails (Collembola)

For control of the above insects, apply a five-foot band of D·z·n Granules next to the foundation of the house. Distribute the granules uniformly using a rate of 8 ozs. for each 5' by 20' area to be treated.

Storage and Disposal

Storage: Store in original container in a cool, dry, well-ventilated area inaccessible to children and domestic animals. Do not store near fertilizers, herbicides, seeds, food, or feed. Do not reuse bag. Discard bag in trash.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse empty bag. Discard bag in trash.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust and contact with eyes, skin, or clothing. Wash thoroughly after handling and before smoking or eating. May cause contact sensitization following repeated contact with skin of susceptible individuals. If sensitization reactions result, contact a physician. Do not store near food or feed products. Food utensils such as tablespoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets onto the treated area until granules have been watered in and the grass has dried.

Statement of Practical Treatment

If swallowed, call a physician immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If inhaled, remove victim to fresh air and apply artificial respiration if indicated.

If on skin, wash promptly with soap and water. Rinse thoroughly.

If in eyes, rinse eyes with plenty of water and call a physician immediately.

Note to Physician:

This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Environmental Hazards

This product is highly toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water or wetlands (swamps, bogs, marshes, and pot-holes). Runoff may be hazardous to aquatic organisms in neighboring areas. Collect or incorporate granules that are spilled during loading or are visible on soil surfaces in turn areas. Do not contaminate water by disposal of equipment washwaters.

Continued...

Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates. Rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of granules. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources.

D·z·n® trademark of CIBA-GEIGY for diazinon

EPA Reg. No. 100-468
EPA Est.

© 1989 CIBA-GEIGY Corporation

Agricultural Division
CIBA-GEIGY Corporation
Greensboro, North Carolina 27419

CGA 9L7T

April 6, 1988-
Changed name to
D z·n, revised
Storage & Dis-
posal, added do
not use on golf
courses/sod farms
statement.
Accepted December 28, 1988
Data Call-In
revisions -
reduced rates on
turf to a max.
of 4 lbs. a.i./A,
added avian pre-
cautions, revised
precautionary
statements,
storage and
disposal, Env.
Hazards.
Revised October 10, 1989
Reg. Standard
changes, added
warranty state-
ment, light
watering for
surface feeding
pests.