

Alfa-tox

Insecticide

AMPHLE 1000

For control of insects on alfalfa, clover, vegetables, and small fruits

Active Ingredients:	
Methoxychlor, Insecticide	20%
Diazinon, Insecticide	10%
Inert Ingredients	63%
Total	100%

Eq. 1000 (1000) 2.2 bis. p. 1000
Phy. 1000 (1000) 1000
1000 (1000) 1000
1000 (1000) 1000
1000 (1000) 1000

Alfa-tox contains Diazinon and Methoxychlor

Caution:

Keep out of reach of children.

Avoid contact with skin, eyes, and clothing. Avoid contamination of food and feed. Wash thoroughly after handling. Harmful if swallowed. May be absorbed through skin. Avoid breathing spray mist.

Fish and Wildlife Caution:
This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes and containers.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the Directions For Use, Conditions of Sale, and Warranty.

Directions For Use
Directions For Use
Directions For Use
Directions For Use
Directions For Use
Directions For Use
Directions For Use
Directions For Use
Directions For Use
Directions For Use

Directions For Use
Directions For Use
Directions For Use
Directions For Use
Directions For Use
Directions For Use
Directions For Use
Directions For Use
Directions For Use
Directions For Use

Directions For Use
Directions For Use
Directions For Use
Directions For Use
Directions For Use
Directions For Use
Directions For Use
Directions For Use
Directions For Use
Directions For Use

1000 (1000) 1000
1000 (1000) 1000

Geigy

Alfa-tox

General Information

Alfa-tox is an emulsifiable solution which can be diluted with water. When ground applications are to be made, dilute Alfa-tox with water according to the Directions for Use section. For application by aircraft, refer to the aerial application section of the label. Alfa-tox is intended for use against insect pests of agricultural crops and should not be used in barns or dwellings. Applications should be repeated as necessary to maintain control.

Directions for Use

Forage Insects

Alfalfa and Clover (Pure Stands and Stands Containing Grass)

Alfalfa Weevil Larvae, Egyptian Alfalfa Weevils, Pea Aphids, Yellow Clover Aphids, Spotted Alfalfa Aphids, Leafhoppers, Plant Bug Nymphs, Mites. Apply 2-3 qts. per acre in sufficient water to achieve thorough coverage. Apply when 50% of the growing plants are feeding damage. Repeat as infestations warrant. Do not treat during bloom. Do not apply within 14 days of harvest.

Alfalfa Leafhoppers, Armyworms, Fall Armyworms, Flea Beetles, Grasshoppers, Japanese Beetles, Lygus Bugs, Spittlebugs. Apply 2 1/2 - 3 qts. per acre in sufficient water to achieve thorough coverage when insects first appear. Repeat as infestations warrant. Do not treat during bloom. Do not apply to beneficial insects.

Note: Dairy and beef cattle and sheep may be grazed 7 days following treatment. Do not feed green chop to dairy cattle or sheep until 14 days after treatment.

Small Fruit Insects

For the control of the insects on the following crops, apply 2 1/2 qts. per 100 gals. of water per acre as a thorough cover spray when insects first appear. Repeat applications at 7-10 day intervals as infestations warrant.

Cranberries

Cranberry Fruitworms, Leafhoppers, Spittlebugs. Do not apply within 14 days of harvest.

Grapes

Flea Beetles, Grape Berry Moths, Grape Leaf Folders, Grape Leaf Skeletonizers, Japanese Beetles, Leafhoppers, Pacific Mites, Rose Chafers. For Grape Berry Moths, apply as a post-blossom cover spray and repeat 10 days later, make third and fourth applications at 10-day intervals if necessary. Do not apply within 14 days of harvest.

Strawberries

Aphids, Flea Beetles, Spittlebugs, Strawberry Leaf Beetles, Strawberry Leaf Rollers, Strawberry Aphids, Two-spotted Mites. Do not apply within 5 days of harvest.

Vegetable Insects

For the control of the insects on the following crops, use 2 1/2 qts. per acre in sufficient water for thorough coverage. Begin treatment at first sign of insect infestation and repeat at 7-10 day intervals as infestations warrant.

Broccoli

Aphids, Imported Cabbageworms, Diamondback Moths, Flea Beetles.

Cabbage

Aphids, Imported Cabbageworms, Diamondback Moths, Flea Beetles. Do not apply within 7 days of harvest.

Cauliflower

Aphids, Imported Cabbageworms, Diamondback Moths, Flea Beetles. Do not apply within 7 days of harvest.

Carrots

Aphids, Leafhoppers. Do not apply within 10 days of harvest or 14 days if tops are used for food or feed.

Collards

Aphids, Corn Earworms, Diamondback Moths, Flea Beetles, Harlequin Cabbage Bugs, Imported Cabbageworms. Do not apply within 14 days of harvest.

Corn

Fall Armyworms, Flea Beetles, Grasshoppers, Japanese Beetles, Rose Chafers. Do not apply within 7 days of harvest.

Cucumbers

Aphids, Cucumber Beetles, Spittlebugs, Leaf Miners, Lygus. Do not apply within 7 days of harvest.

Kale

Aphids, Corn Earworms, Diamondback Moths, Flea Beetles, Harlequin Cabbage Bugs. Do not apply within 14 days of harvest.

Lettuce

Aphids, Dipterous Leaf Miners, Leafhoppers. Do not apply within 14 days of harvest.

Melons (Cantaloupes, Muskmelons, Watermelons)

Leafhoppers, Cucumber Beetles, Spittlebugs, Mites, Flea Beetles, Two-spotted Mites. Do not apply within 14 days of harvest.