

100-479

03/17/1965

Beigy  
PRAMITOL 5G

ACCEPTED
MAR 17 1965
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER- ED UNDER NO. 100-479

1/2  
74

ACTIVE INGREDIENTS:

Sodium chlorate (NaClO <sub>3</sub> ) . . . . .	40.0%
Sodium metaborate (Na <sub>2</sub> B <sub>2</sub> O <sub>4</sub> 5H <sub>2</sub> O)* . . . . .	50.0%
2-methoxy-4,6-bis (isopropylamino)-s-triazine . . . . .	5.0%
INERT INGREDIENTS: . . . . .	5.0%
<hr/>	
TOTAL . . . . .	100.0%

\*Equivalent to 4.97% boron expressed as elemental boron.

USDA Reg. No. 100-

CAUTION: Keep out of reach of children. Harmful if swallowed. Avoid contamination of feed and foodstuffs. Avoid inhalation of dust. Avoid contact with eyes. Avoid contact with skin and clothing. Wash clothing and equipment after application. Do not use such equipment for other insecticides, fungicides or fertilizers as plant injury may result. Do not contaminate water intended for irrigation or domestic purposes.

U. S. Patent No. 2,909,420

FOR NONSELECTIVE OR INDUSTRIAL WEED CONTROL ON NONCROP LAND SUCH AS: Industrial Plant Sites, Rights-of-Way, Lumber Yards, Petroleum Tank Farms and Similar Areas

GENERAL INFORMATION

Pramitol 5G is a broad spectrum, nonselective, deep-rooted broadleaf and grass weed killer. It is effective when applied before or after weeds emerge. Since Pramitol 5G acts through the roots, its effect on weeds is dependent upon adequate rainfall to move the chemical into the root zone. Excessively dry soil conditions and absence of rain may result in poor weed control.

Care should be taken to avoid the use of Pramitol 5G where adjacent desirable trees, shrubs or plants may be injured. Use only in areas where complete control of all vegetation is desired.

75 2/2

DIRECTIONS FOR USE

Apply 1/2 - 1 lb. per 100 sq. ft. of soil surface to control annual broadleaf weeds and grasses.

Apply 1 - 2 lbs. per 100 sq. ft. of soil surface to control perennial broadleaf weeds and grasses. Where deep-rooted perennial grasses are predominant, use the higher rate.

Use the higher application rates in areas of high rainfall, long growing season and where extended residual control is desired.

Since weather and other conditions under which this product may be used will vary, GEIGY AGRICULTURAL CHEMICALS, Division of Geigy Chemical Corporation and the Seller make no warranty of any kind, express or implied, concerning the use of this product. The user assumes all risks of use or handling whether or not in accordance with directions or suggestions.

GEIGY AGRICULTURAL CHEMICALS  
Division of Geigy Chemical Corporation  
SAW MILL RIVER ROAD ARDSLEY, NEW YORK

015