



Ametryne[®] 2B

Herbicide

For weed control in corn (post-emergence directed spray to weeds)

Active Ingredients:.....	25.3%
2-ethylamino-4-	
isopropylamino-6-	
methylthio-s-triazine.....	24.0%
Related compounds.....	1.3%
Inert Ingredients:.....	74.7%

	Total 100%

Ametryne[®] 2B contains 2 lbs. technical Ametryne per gallon

CAUTION:

Keep out of reach of children.

Harmful if swallowed. Avoid contamination of food and feed, contact with eyes, contact with skin, and breathing spray mist. In case of eye contact, wash with plenty of water. In case of skin contact, wash with soap and water.

Do not contaminate domestic or irrigation water supplies or lakes, streams or ponds.

Do not use or store near heat or open flame.

Do not reuse container. Destroy when empty.

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions For Use and the Conditions Of Sale And Warranty before using this product.

Conditions Of Sale And Warranty

The Directions For Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Geigy or the Seller. All such risks shall be assumed by the Buyer.

Geigy warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions For Use, subject to the inherent risks referred to above. Geigy makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Geigy or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. Geigy and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions Of Sale And Warranty which may be varied only by agreement in writing signed by a duly authorized representative of Geigy.

GENERAL INFORMATION

Ametryne 2E herbicide controls most annual broadleaf and grass weeds, such as Texas panicum (Texas millet), fall panicum, signalgrass (*Brachiaria* spp.), goosegrass, crabgrass, barnyardgrass, giant foxtail, yellow and green foxtails, cocklebur, lambsquarters, Florida purslane, morningglory, pigweed, wild mustard, ragweed, and smartweed. Ametryne is effective as a post-emergence herbicide, having considerable activity through foliage contact.

Care should be taken to avoid its use where adjacent to desirable trees, shrubs, or other plants or injury may occur. Ametryne 2E is noncorrosive to equipment and metal surfaces and has low electrical conductivity.

DIRECTIONS FOR USE:

Corn (Southeast and South Central Region):

Apply Ametryne 2E as a post-emergence directed spray to weeds after the smallest corn is at least 12 inches high (measured to the highest leaf surface on free standing plants).

CAUTION:

Do not spray over top of corn or injury will occur.

Apply in a minimum of 20 gallons of water per acre to assure uniform coverage (non-pressure nitrogen solution may be substituted for all or part of the water). Best weed control is attained when weeds are 2 inches or less in height. The addition of a surfactant is recommended such as Colloidal Products X-77, Dupont EK, or Tronic, at the rate of 0.5% of the spray volume (2 qts./100 gals. of spray solution).

It is recommended that gauge wheels and/or leaf lifter equipment be used to prevent leaf contact with the spray. Apply at a spray pressure of 30 psi or less to prevent "bounce back" -- spray bouncing off soil or weeds and settling on the corn leaves.

Weeds	Weed Height	Rate per Acre	
		Broadleaf	20" Band
Brachiaria Broadleaves	Up to 2 in. Up to 2 in.	1 qt.	1/2 qt.
Brachiaria Broadleaves Texas panicum	Up to 4 in. Up to 4 in. Up to 2 in.	2 qts.	1 qt.
Brachiaria Broadleaves Crabgrass Texas panicum and other grasses	Up to 6 in. Up to 6 in. Up to 2 in. Up to 4 in.	3 1/2-4 qts.	1 3/4-2 qts.

*Based on 40-inch row spacing. For other band or row widths, adjust the rate in proportion to the area actually treated.

Precautions: 1) Do not apply to corn less than 12 inches high or at a spray pressure above 30 psi or injury may occur. 2) Allow 30 days after application before harvesting, grazing or feeding forage to livestock. 3) Ametryne treated land should not be planted to any rotational crop except corn until the following year or injury may occur.

USDA Reg. No. 100-480
U. S. Patent No. 2,909,420

GEIGY
Geigy Agricultural Chemicals
Division of Geigy Chemical Corporation
Ardsley, New York