

Acaralate 2E

Ingredients

Paracetamol	1.0
Spill	1.0
Isobutyl nitrate	1.0
Water	1.0
Active Ingredients:	
Chloropropylate	1.0
dichlorobenzilate	1.0
Aromatic Petroleum	1.0
Derivative Solvent	1.0
Inert Ingredients	1.0
Total:	10.0

Acaralate 2E contains 2 lbs.
chloropropylate per gallon.

Caution:

Keep out of reach of children.

Harmful if swallowed. Do not breathe
spray mist. Avoid contact with eyes.
Avoid prolonged or repeated contact
with skin. Do not use in dwellings.

Fish Caution:

This product is toxic to fish. Keep
out of lakes, streams, or ponds. Do
not apply where runoff is likely to
occur. Do not apply when weather
conditions favor drift from areas
treated. Do not contaminate water
by cleaning of equipment or disposal
of wastes. Apply this product only
as specified on this label.

Do not reuse empty container. De-
stroy by perforating or crushing and
burying in a safe place.

Flammable mixture:

Do not use, pour, spill or store near
heat or open flame. Store at tempera-
tures above 32°F.

EPA Reg. No. 100-485-AA

CONDITIONS OF USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions For Use and the Conditions Of Sale And Warranty before using this product.

Conditions Of Sale And Warranty

The Directions For Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application all of which are beyond the control of CIBA-GEIGY or the Seller. All such risks shall be assumed by the Buyer.

CIBA-GEIGY warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions For Use, subject to the inherent risks referred to above. CIBA-GEIGY makes no other express or implied warranty of fitness, merchantability or any other express or implied warranty. In no case shall Ciba-Geigy or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. CIBA-GEIGY and the seller offer this product and the Buyer and user accept it, subject to the foregoing Conditions Of Sale And Warranty which may be varied only by agreement in writing with a duly authorized representative of CIBA-GEIGY.

General Information

Acaralate 2E miticide is an emulsifiable solution intended for use in mixing sprays which, when applied as directed, will control European Red Mites and Two-Spotted Spider Mites on apples and pears. Thorough coverage of trees is essential for effective control. Either dilute or concentrate spray equipment can be used, providing the spray reaches all parts of the tree, especially the undersides of the leaves. The mixing rates listed below are offered as a guide for preparing dilute, high-gallonage sprays. The suggested rates should be adjusted accordingly for applications with concentrate sprayers.

Acaralate 2E should be applied in pre-bloom sprays or when mite infestations first appear. Severe mite infestations will require repeated applications, properly timed to coincide with mite build-up.

Compatibility

Acaralate is physically compatible with many of the commonly used insecticides and fungicides such as: Captan WP; Diazinon EC, WP; Guthion EC, WP; lead arsenate WP; Dieldrin WP; and zineb WP. Acaralate has been found to be incompatible with dodine (Cyprex) products, urea, Dyrene WP, sulfur, tribasic copper sulfate, and disulfene (Thysan XL). When Acaralate is to be combined with other spray materials the physical compatibility of the mixture should be established by premixing the materials in small quantities.

Directions for Use

Apples and Pears

Two-Spotted Spider Mites, European Red Mites: Apply 1 1/2 pts. per 100 gals. of dilute spray in prebloom or regular spray at whenever infestations first appear. Repeat applications as needed. In areas where new mite infestations occur, make two applications of 2 pts. per 100 gals. of dilute spray 7-10 days apart.

- 4 -

Do not apply to apples or pears within 14 days of harvest.
Do not apply in combination with spray oils as injury may
occur. In western states, to avoid injury to fruit, do
not apply with hand gun. Do not graze treated areas.

Acaralate: trademark of CIBA-GEIGY for chloropropylate

Agricultural Division
CIBA-GEIGY Corporation
Greensboro, North Carolina 27409

October 19, 1973