

## **PLEASE NOTE**


**This image contains more than one label approved for this product on this date.**

100-501

04/11/1975

1/1

SUPRACIDE® 2E  
EPA Reg. No. 100-501  
PROPOSED LABEL REVISIONS



---

Directions for Use

---

Sunflowers

Apply 2 pts. per acre. Make ground application in a minimum of 10 gals. of water per acre; aerial application in a minimum of 5 gals. of water per acre. Thorough coverage of the heads is necessary. One to three applications may be made on a 7 day schedule. The number of applications will be determined by the severity of the infestations.

Sunflower Moth, Banded Sunflower Moth, and Sunflower Maggot Complex: Begin applications at the onset of bloom as the infestations develop.

Sunflower Seed Midge: Begin applications during early bud development.

Note: Do not apply within 50 days of harvest.

Do not graze treated areas or feed treated forage to livestock.

CIBA-GEIGY Corporation  
Agricultural Division  
Greensboro, North Carolina 27409


**NEXT**

**LABEL**

100-501

04/11/1975

Y/1



PROPOSED LABEL CLAIMS

SUPRACIDE® 2E - SORGHUM

EPA REG. NO. 100-501



DIRECTIONS FOR USE

Grain Sorghum Grown for Forage and Grain (Texas Only)

Make application in a minimum of 10 gals. of water per acre by ground application or 2 gals. of water per acre by aerial application. Up to three applications may be made on a 10-14 day schedule. The number of applications will be determined by the severity of the infestation.

Banks Grass Mite: Apply 2 pts. per acre.

Note: Application may result in phytotoxicity to certain sorghum varieties. This phytotoxicity is a temporary spotting, reddening or chlorosis of the leaves.

Do not apply within 30 days of harvesting or feeding to livestock.

Supracide® trademark of CIBA-GEIGY for Methidathion  
U.S. Patent Number 3,230,230 and 3,240,668

Agricultural Division  
CIBA-GEIGY Corporation  
Greensboro, North Carolina 27409

January 22, 1975