

100-501



06/27/1975

1/2

Best Available Copy

Supracide®

2E



Insecticide- Miticide

For control of certain insects of alfalfa, cotton, tobacco, grapefruit, and oranges

Five Gallons

U.S. Standard Measure

Active Ingredients:	91.7%
Methidathion: O,O-dimethyl phosphorodithioate, S-ester with 4-(mercaptomethyl)-2-methoxy-2,1,3,4-thiazolidin-5-one	24.4%
Aromatic petroleum derivative solvent	67.3%
Inert Ingredients:	8.3%
Total:	100.0%

Supracide 2E contains 2 lbs. active ingredients per gal.

Danger:

Keep out of reach of children. See additional danger statements on back of container.

EPA Reg. No. 100-501-AA

EPA Est. No. 100-AL-1

CIBA-GEIGY

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire **Directions for Use** and the **Conditions of Sale and Warranty** before using this product.

Conditions of Sale and Warranty

The **Directions for Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of CIBA-GEIGY or the Seller. All such risks shall be assumed by the Buyer.

CIBA-GEIGY warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions for Use**, subject to the inherent risks referred to above. **CIBA-GEIGY makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty.** In no case shall CIBA-GEIGY or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. CIBA-GEIGY and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of CIBA-GEIGY.

General Information

Supracide 2E is an emulsifiable concentrate which, when diluted with water according to the **Directions for Use** given below, will control certain insect pests and mites on agricultural crops.

Do not enter treated fields the same day of application. Wear adequate protective clothing during the preparation of spray mixture. During application stop when entering treated fields. Wash thoroughly before eating and at the end of each day's operation. Change clothing daily.

Directions For Use

Alfalfa: pure stands or stands containing clover or grass). Make applications at the rates listed below in a minimum of 10 gals. of water per acre for ground application or apply in a minimum of 2 gals. of water for aerial applications. Foliage applications for alfalfa weevil and Egyptian alfalfa weevil control should be made to standing alfalfa when 20-30% of the growing tips show feeding damage. Stubble applications should be made immediately following a cutting. For the other insects listed on alfalfa, apply when insects first appear. Make no more than one stubble application and no more than one foliage application per alfalfa cutting.

Alfalfa weevil, Egyptian alfalfa weevil*, aphids, leathoppers*, lygus bugs* and spittle bugs**. Apply 2-4 pts. per acre (Southern States: apply 3-4 pts. per acre).

*Pacific and Intermountain States only
**Northeastern U.S. only

Note: 1) To avoid injury to pollinating insects, do not apply during bloom. 2) Do not apply within 10 days of harvesting or feeding to livestock.

Cotton

Make applications at the rates listed below in a minimum of 10 gals. of water per acre for ground application or apply in a minimum of 3 gals. of water per acre for aerial application. Primarily for spider mite control, if bollworm and boll weevil are present at the time of application for spider mites, they will be controlled. Repeat applications can be made if necessary at 5-7 day intervals.

Spider mites, boll weevil, bollworm: Apply 4 pts. per acre when mite foliar feeding symptoms are present (feeding symptoms are red and yellow speckling of the leaves).

Note: 1) Do not make more than 3 applications per year. 2) Do not apply after the bolls begin to open, or within 60 days of harvest. 3) Do not graze treated cotton plants or feed gin waste to livestock.

Oranges and Grapefruit

California — California Red Scale
Texas — California Red Scale, Chaff Scale
Florida — Black Scale, Chaff Scale, Glovers Scale, Purple Scale, Snow Scale, Yellow Scale, White Flies

Apply 1-2 pts. per 100 gals. of dilute spray. A second application may be made if necessary. Allow at least 45 days between applications. For best results, apply in postbloom and/or summer sprays. Severe infestations may require treatment at both times. Use the low rate for light to moderate infestations, and the high rate for heavy infestations. Thorough coverage is essential for good control.

Do not apply more than 40 pts. per acre. The minimum amount of Supracide 2E applied per acre is dependent on the number of trees per acre and on tree height. See example below.

EXAMPLE:

Mature trees of normal foliage require at least the height (in feet) of the tree plus 5 gals. of water for adequate coverage. Eighty trees per acre 20 feet tall require $80 \times 25 = 2,000$ gals. of dilute spray per acre.

NOTE: Do not graze treated areas. Make no more than 2 applications per growing season and do not apply within 14 days of harvest.

Tobacco

Make applications at the rates listed below in a minimum of 25 gals. of water per acre when insects appear. Apply 1-3 repeat applications at a 7-14 day interval as needed.

Flea beetles, hornworms: Apply 2-3 pts. per acre
Budworms: Apply 4 pts. per acre.

Note: Do not apply in mixture with other pesticides or within 3 days of tobacco harvest.

Danger

Keep out of reach of children.

Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when mixing. Avoid contamination of food and feed.

In case of skin contact, wash with plenty of water. In case of eye contact, flush with water for 15 minutes.

Harmful or fatal if swallowed. Do not breathe spray mist (prolonged or repeated inhalation of mist may be harmful). Wear an approved respirator during prolonged use.

Keep out of reach of domestic animals.

Not for use or storage in or around the home.

Note to Physician:

Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

If swallowed and gastric lavage is not immediately available, induce vomiting taking care to avoid inhalation of stomach contents into the lungs.

Fish and Wildlife Caution

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when the weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy it by perforating or crushing and burying in a safe place.

Do not use, pour, spill or store near heat or open flame.

Store at temperatures above 32° F.

Supracide® trademark of CIBA-GEIGY for methidathion
U.S. Patent No. 3,230,230 and 3,240,668

Agricultural Division
CIBA-GEIGY Corporation
Greensboro, North Carolina 27409

CGA 30LIC

084

2/2