

Compatibility test: Nitrogen solutions or complete liquid fertilizers may replace all or part of the water in the spray in corn. Since liquid fertilizers can vary, even within the same analysis, check compatibility each time before use. Be especially careful when using complete suspension or fluid fertilizers as serious compatibility problems are more apt to occur. Commercial application equipment may improve compatibility in some instances. The following test assumes a spray volume of 25 gals. per acre. For other spray volumes, make appropriate changes in the ingredients. Check compatibility using this procedure:

1. Add 1 pint of fertilizer to each of 2 one-quart jars with tight lids.
2. To one of the jars add 1/4 tsp. or 1.2 milliliters of a compatibility agent approved for this use (1/4 tsp. is equivalent to 2 pts. per 100 gals. spray). Shake or stir gently to mix. Examples of compatibility agents include Compax and Unde.
3. To both jars add the appropriate amount of herbicide(s). If more than one herbicide is used, add them separately with dry herbicides first, flowables next, and emulsifiable concentrates last. After each addition, shake or stir gently to thoroughly mix. The appropriate amount of herbicides for this test follows:

Dry herbicides: For each pound per acre add 1.4 teaspoons to each jar.
Liquid herbicides: For each pint per acre add 0.5 teaspoon or 2.5 milliliters to each jar.

4. After adding all ingredients, put lids on and tighten, and invert each jar ten times to mix. Let the mixtures stand 15 minutes and then look for separation, large flakes, precipitates, gels, heavy oil film on the jar, or other signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the two jars. If either mixture separates, but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility: (A) slurry the dry herbicide(s) in water before addition, or (B) add 1/2 of the compatibility agent to the fertilizer and the other 1/2 to the emulsifiable concentrate or flowable herbicide before addition to the mixture. If still incompatible, do not use the liquid fertilizer and herbicide(s) in the same spray tank.

Rotational Crops and Perennial Crop Replanting

To avoid crop injury, observe the following precautions: 1) If rotating treated land the year following application, plant only corn, unless otherwise stated in this label. 2) If replanting perennial crops or if rotating land to crops other than corn, do not apply Princep in the year preceding planting these crops.

Fruit and Nut Crops

Apply the spray to the orchard or vineyard floor avoiding contact with fruit, foliage, or stems. Apply only to orchards or groves where trees have been established one year or more unless specified differently. Make only one application per year, except as noted otherwise. Recommended rates are based on broadcast treatment. For band applications or spot applications around trees in fruit or nut plantings, reduce the broadcast rate of Princep 4L and water per acre in proportion to the area actually sprayed.

Almonds and Peaches (CA only)

Apply 1-2 qts. of Princep 4L per treated acre in a 2-4 ft. band on each side of the tree row. Apply before weeds emerge in late fall or early winter. Weeds controlled by 1 qt. of Princep 4L include burdock, common chickweed, wild mustard, and shepherdspurse. Apply only once per year.

Precautions: 1) Do not treat trees established in the grove less than 3 years. 2) Do not treat the Mission (Texas) variety of almonds. 3) Do not apply to almond trees propagated on plum rootstocks. 4) Do not replant almonds or peaches in treated soil for 12 months after treatment. 5) Do not apply on gravelly, sandy, or foamy sand soil or on soil with less than 1% organic matter. 6) Do not treat areas where water will accumulate.

Apples, Peats, Sour Cherries

Apply 2-4 qts. per acre

Avocados

CA and FL only. Apply 2-4 qts. per acre after final preparation of grove.

Precaution: Do not apply on gravelly, sandy, or foamy sand soil.

Blueberries and Caneberries (blackberries, boysenberries, loganberries, raspberries)

Apply 2-4 qts. per acre in the spring or apply a split application of 2 qts. per acre in the spring plus 2 qts. per acre in the fall. Apply in a minimum of 40 gals. of water per acre.

On plantings less than 6 months old, use 1/2 the above rate.

To control quackgrass, apply 4 qts. per acre in the fall, or split the application applying 2 qts. per acre in the fall plus 2 qts. per acre in the spring when quackgrass is actively growing.

Note: Do not apply when fruit is present.

Cranberries

MA. Apply up to 4 qts. per acre after before spring growth begins or in the fall after harvest.

Other areas. Apply 2 qts. per acre before spring growth begins.

Filberts

Apply 2-4 qts. per acre in the fall or apply a split application of 2 qts. per acre in the fall plus 2 qts. per acre in the spring.

Precautions: 1) If trees are planted on a hillside, excessive soil erosion may result from the elimination of weeds. 2) Do not apply on gravelly, sandy, or foamy sand soil.

Note: Do not apply when nuts are on the ground during the harvest period.

Grapes

Apply 2-4 qts. per acre any time between harvest and early spring.

Precaution: Do not use in vineyards established less than three years. Do not apply on gravelly, sandy, or foamy sand soil.

Grapefruit, Lemons, Oranges

AZ (Lemons and Oranges only): Apply a split application of 1.6 qts. per acre in the spring plus 1.6 qts. per acre in the fall.

CA: Apply 2-4 qts. per acre in a single application, or apply 2 qts. per acre in the fall and 2 qts. per acre in the spring. Do not use in the Imperial, Coachella, or Palo Verde Valleys.

FL (Grapefruit and Oranges only): Apply only once per year. For control of weeds listed under General Information, apply 6.4-9.6 qts. per acre in the spring; to control milkweed vine, use 8-9.6 qts. per acre either preemergence or before vines are 12 inches tall.

TX (Grapefruit and Oranges only): Apply 4-8 qts. per acre.

Precautions (All areas): 1) Do not use in nurseries. 2) Do not apply to bedded grapefruit, lemons, or oranges. 3) To avoid possible injury, do not apply to trees under stress from freeze damage for one year after the freeze.

Macadamia Nuts

Apply 2-4 qts. in 60 gals. of water per acre before harvest and just prior to weed emergence. Repeat application as necessary. Do not apply when nuts are on the ground during the harvest period.

Olives

Apply 2-4 qts. per acre following grove preparation in the fall. Repeat annually in midwinter.

Peaches, Plums, Sweet Cherries

Apply 1.6-4 qts. per acre. Apply in late fall to early spring prior to weed emergence.

Precautions: 1) Do not apply on gravelly, sandy, or foamy sand soil. 2) Peaches: use only in AR, LA, MO, OK, TX, and states east of the Mississippi River. For California, see specific directions in the section Almonds and Peaches (CA only). 3) Plums and sweet cherries: use only in MO and states east of the Mississippi River except TX.

Pecans

Apply the appropriate rate in the following table before weeds emerge in the spring.

Soil texture	Rate per acre
Sand, foamy sand	Do not use
Sandy loam	2 qts.
Loam or clay soil low in organic matter	2-2.8 qts.
Clay soil high in organic matter	2.8-4 qts.

Precautions: Do not use west of the Pecos River in TX or in NM, AZ, or CA, as injury will occur. Do not make application to transplanted trees that have been established less than two years in the grove or injury may occur.

Note: Do not apply when nuts are on the ground. Do not allow animals to graze treated areas.

Strawberries

Oregon and Washington: For control of chickweed, groundnut mustard, and shepherdspurse, apply broadcast 1 qt. per acre in fields where overhead irrigation is used to activate this product. Apply after harvest at time of bud renovation. In fields where overhead irrigation is not available, apply during early October through November.

Precautions: Make only one application per growing season. Do not apply within 4 months after transplanting. Limit overhead irrigation to 1 inch immediately following postharvest application. Do not use on sand.

Walnuts

Apply 2-4 qts. per acre

Precautions: 1) Do not apply on gravelly, sandy, or foamy sand soil. 2) Leveling and furrowing operations after application will lessen effectiveness of weed control.

Note: Do not apply when nuts are on the ground.

BEST DOCUMENT AVAILABLE

added

Soil texture	Broadcast rate per acre			
	1:1 Ratio		2:1 Ratio	
	Princep 4L	AAtrex 80W*	Princep 4L	AAtrex 80W*
Sand, loamy sand, sandy loam	1 qt	1.25 lbs	1.33 qts	0.83 lb
Loam, silt loam, silty clay loam, sandy clay loam, silty clay, or silty clay with low organic matter	1.2 qts	1.5 lbs	1.6 qts	1 lb
Loam, silt loam, silty clay loam, sandy clay loam, silty clay, or silty clay with medium to high organic matter, and clay (including dark prairie soils of the Corn Belt)	1.44 qts	1.8 lbs	1.92 qts	1.2 lbs

*When using AAtrex 4L (4LC) or AAtrex Nine-O, use equivalent rates. One lb. of AAtrex 80W equals 0.8 qt. of AAtrex 4L (4LC) or 0.9 lb. of AAtrex Nine-O.

Refer to the AAtrex label for complete directions. Also refer to the AAtrex label and the Corn section of this label for precautions and limitations.

Eradicane: Use in a preplant incorporated tank mixture for control of all weeds claimed on both the Princep and Eradicane labels including partial control (suppression) of shattercane (wild cane). Fluid fertilizer may replace all or part of the water in the spray. Check the physical compatibility of mixture with fertilizer before use. Refer to the General Information section of this label for a compatibility test procedure. Use a minimum of 20 gals. of spray volume per acre. Refer to the Eradicane label for incorporation directions. Use the higher rate of Princep on fine textured soil and where heavy broadleaf infestations are expected. Use the higher rate of Eradicane for heavy nutsedge and bermudagrass infestations. For partial control (suppression) of shattercane, broadcast and incorporate immediately before planting 2-4 pts. of Princep 4L plus 7.33 pts. of Eradicane 7E (or 8 pts. Eradicane Extra) per acre. For control of the other weeds claimed on both labels, broadcast and incorporate 2-6 pts. of Princep 4L plus 4.75-7.33 pts. of Eradicane 7E (or 5.33-8 pts. of Eradicane Extra) per acre. Observe all precautions and limitations on the Princep 4L and Eradicane labels.

Paraquat: Use in a tank mixture where corn will be planted directly in a cover crop, established sod, or previous crop residue. This combination controls existing vegetation and provides residual control of the annual broadleaf and grass weeds listed under General Information.

Add Princep 4L to the spray tank, mix thoroughly with water, and then add paraquat and a nonionic surfactant. Provide constant agitation during mixing and application to keep the mixture in suspension.

Apply 4-6 pts. of Princep 4L plus 1-2 pts. of paraquat* in 20-60 gals. of water per acre as a broadcast spray either before or after planting but before corn emerges. Add a nonionic surfactant at the rate of 0.5 pt. per 100 gals. of spray volume.

For further information, see general information, caution and warning statements, precautions, and notes on the Princep 4L and the paraquat labels.

rev. seed

*Based on a product containing 2 lbs. paraquat cation per gal.

Sutan +: Use in a preplant incorporated tank mixture for control of all weeds claimed on both the Princep and Sutan + labels including partial control (suppression) of shattercane (wild cane). Fluid fertilizer may replace all or part of the water in the spray. Check the physical compatibility of mixture with fertilizer before use. Refer to the General Information section of this label for a compatibility test procedure. Use a minimum of 20 gals. of spray volume per acre. Refer to the Sutan + label for incorporation directions. Use the higher rate of Princep on fine-textured soil and where heavy broadleaf infestations are expected. For partial control (suppression) of shattercane, broadcast and incorporate immediately before planting 2-4 pts. of Princep 4L plus 7.33 pts. of Sutan + 6.7E per acre. For control of the other weeds claimed on both labels, broadcast and incorporate 2-6 pts. of Princep 4L plus 4.75 pts. of Sutan + 6.7E per acre. Observe all precautions and limitations on the Princep 4L and Sutan + labels.

Sugarcane

Apply 2-4 qts. per acre at planting or after harvest if ratoon crops to be grown, but before weeds or sugarcane emerge. Princep 4L may also be applied interline directed or over the sugarcane to provide continuing control of germinating weeds.

Precautions: 1) Do not apply more than one 2.4 qt. application over the plants or more than two interline directed applications. 2) Do not apply after close in 31. Total application should not exceed 10 qts. per acre crop cycle.

Note: Do not feed or graze treated forage crops.

Grasses Grown for Seed (Pacific Northwest Only)

Perennial ryegrass, Highland or Astoria bentgrass, orchardgrass, timothy, and fine fescues such as named Ramer Penn and Chewings and related species. For the control of broadleaf weeds and annual grasses including annual ryegrass, wild mustard, silver chaff, grass, and downy brome, apply 2 qts. Princep 4L (4LC) in a minimum of 20 gals. of water per acre as soon as weeds start. Early application is important.

Precaution: Apply only to grasses from which at least one seed crop has been cut.

Note: Allow 30 days between last application and grazing of dairy and beef cattle and sheep, and allow 60 days between the last application and cutting for hay.

Nurseries, Christmas Tree Plantings, Shelterbelts

Nurseries (see list below)

Apply 2-3 qts. in at least 25 gals. of water per acre in fall or spring at least one year after transplanting.

Christmas Tree Plantings and Shelterbelts (see list below)

Remove weed growth before application. Apply 2-4 qts. in at least 25 gals. of water per acre after transplanting. Use the same rate for annual maintenance applications.

For quackgrass control apply 4 qts. per acre in the fall or apply a split application of 2 qts. per acre in the fall plus 2 qts. per acre in early spring, after quackgrass begins growth.

Precautions: 1) Do not use Princep 4L on seedbeds or cutting beds. 2) In CA, OR, and WA, do not apply to Christmas trees or shelterbelts sooner than one year after transplanting. In other areas, do not apply to Christmas trees or shelterbelt transplants less than three years of age. 3) Do not apply more than once a year, except as directed for quackgrass control.

Apply Princep 4L to these species of trees and shrubs, as recommended above:

Conifers

arbutus
Austrian pine
Bishop pine
Knobcone pine
Lodgepole pine (shore pine)
Monterey pine
Mugho pine
red pine (Norway pine)
Scotch pine
white pine
blue spruce
Norway spruce
red spruce
white spruce
balsam fir
Douglas fir
Fraser fir
white fir
hemlock
juniper
red cedar
white cedar
yellow fir

Deciduous Trees and Woody Ornamentals

American elm
Siberian elm
barberry
boxelder
bush honeysuckle
caragana
cottonaster
dogwood
eucalyptus
holly (Ilex spp.)
(max 3.2 qts. A)
honey locust
Oregon grape (Mahonia spp.)
Pieris spp.
(max 3.2 qts. A)
red oak
Russian olive
oleander
palm (2 qts. A)
nettle brush
carob

*For California only

Surflan Tank Mix: On Christmas tree plantings, use this tank mix for preemergence control of weeds listed on the Surflan 75W (or Surflan A 5) and Princep 4L labels. Use on field grown conifer species listed on the labels for each herbicide plus grand fir, alpine fir, Engelmann spruce, black spruce, Colorado blue spruce, Coulter pine, giant redwood, and Veitch fir. Broadcast the mixture as a directed spray to the soil surface or as an overtop spray using 2-4 qts. of Princep 4L and 2-6.75-33 lbs. of Surflan 75W (or 2-4 qts. of Surflan A 5). Apply in sufficient water per acre to uniformly treat the area. Follow overtop sprays with sprinkler irrigation to move the herbicide from leaf surface to the soil. Remove weed growth before application. If a weed requires prunings or flash in the soil, or remove them before treatment. Soil should be in good tilth and free of clods at time of application. Shallow cultivation (1-2 inches) after treatment will not reduce weed control. Observe all precautions and limitations on the Princep 4L and Surflan labels.

added

Tree Plantations for Timber

Apply in the spring before weeds emerge or following removal of weed growth. Apply before or after seeding/transplanting. If applying after transplanting, application may be made over the top of tree seedlings before new growth appears. If new growth is present, direct the spray to avoid tree foliage contact. Do not use on seed nurseries or injury may occur. Apply only once per year.

The following rates are based on broadcast treatment in 20-100 gals. of water per acre. For band or spot applications, reduce the broadcast rate in proportion to the area actually sprayed.

Black Walnuts

Apply 2-3 qts. per acre.

Slash Pine and White Pine

Use for establishment in the South or, Apply 4 qts. per acre.

Loblolly Pine

Use for establishment in the South only. Apply either 4-6 qts. of Princep 4L per acre or a tank mix of 3-6 qts. of Princep 4L plus 1-2 lbs. of AAtrex 80W (or 0.9-1.75 lbs. AAtrex Nine-O or 0.8-1.6 qts. AAtrex 4L or 4LC) per acre. The tank mix provides broad spectrum annual weed control (see list in General Information section of this label) and better weed control than either Princep 4L alone.

deleted
bird foot
trefoil
or
clover

BEST DOCUMENT AVAILABLE

100-526

Tree Seedlings in Illinois

Remove weed growth before application. Apply 2-4 qts. of Princep 4L per acre in a minimum of 25 gals. of water after transplanting but before weed emergence. Use on the following species of trees:

American beech	honey locust	*sweet gum
balsam fir	maple	*white ash
*black locust	pecan	*white oak
black walnut	*red oak	*yellow poplar
eastern red cedar	Russian olive	

*Instead of Princep 4L alone, on these species a tank mix of 2 qts. of Princep 4L + 2.5 lbs. of AAtrex 80W (or 2.2 lbs. AAtrex Nine-O or 2 qts. AAtrex 4L or 4LC) per acre may be applied in a minimum of 25 gals. of water.

Deleted AAtrex 5L

Precautions: Apply only once per year. On sandy soil, do not use more than 2 qts. Princep 4L per acre and do not use the tank mix. Do not apply the tank mix directly to seedlings after the start of bud break.

Turfgrasses for Sod

St. Augustinegrass, Centipedegrass, and Zoysia Grass

Apply 2-4 qts. per acre, according to soil texture as indicated below:

Muck or peat	4 qts.	Old beds	within 2 days after lifting of sod
		New beds	3-4 days after springing or plugging
Sandy soil	2 qts.	Old beds	within 2 days after lifting of sod
		New beds	7-10 days after springing or plugging

If weeds regrow, apply an additional 2 qts. on muck or peat, or 1 qt. on sandy soil.

Precautions: 1) Do not apply within 30 days before cutting or lifting. 2) Do not use north of NC.

Turfgrass for Fairways, Lawns, and Similar Areas

Bermudagrass, Centipedegrass, St. Augustinegrass, and Zoysia Grass

Apply Princep 4L after October 1 before emergence of winter annual weeds for control of annual bluegrass, burclover, carpet burweed, chickweed, corn speedwell, henbit, hop clover, and spurneed. Princep 4L will control annual bluegrass even if it is emerged at time of treatment. For control of summer annual weeds listed in the General Information section of this label, also apply Princep 4L in late winter before the weeds emerge. Apply in a minimum of 15 gals. of water per acre or 1 gal. per 1,000 sq. ft.

When annual bluegrass is the major weed, use 1 qt. of Princep 4L per acre (22 fl. oz. or 75 fl. oz. per 1,000 sq. ft.). Use 2 qts. per acre (44 fl. oz. or 15 fl. oz. per 1,000 sq. ft.) for control of the other weeds named above. Do not exceed 1 qt. per acre per treatment on newly sprigged turfgrass or on hybrid bermudagrass such as Tiflawn, Tifway, and Diamond.

For continued summer annual weed control, apply another 1 qt. per acre at least 30 days after the previous application, but not after April 15. However, do not make more than two applications of this product per year.

Precautions: To avoid injury, use only on turfgrass reasonably free of infestations of insects, nematodes, and diseases. Do not use on golf greens. Do not use north of NC or west of the high rainfall areas of eastern OK and eastern TX. Do not use on muck or alkaline soils. Do not apply over the rooting area of trees or ornamentals not listed on this label. On newly sprigged turfgrass, temporary slowing of growth may follow application. On hybrid bermudagrass temporary yellowing may occur. Do not overseed with desirable turfgrass within 4 months before or 6 months after treatment.

Vegetable Crops

Asparagus (Established)

Apply 2-4 qts. per acre to beds established one year or more. Apply once per year after disking in the spring before weeds emerge or following harvest after removal of existing weed growth. Spring application must be made at least 3 days before first cutting.

Precautions: (WA and OR only) 1) Do not apply on gravelly sand, or heavy sand soil for crop injury may result. 2) Do not allow irrigation water to stand or to form pools in the field or injury may occur.

Artichokes

Apply 2-4 qts. per acre as a directed spray after the last foliage stage.

Precautions: 1) Do not apply on gravelly sand or heavy sand soil. 2) Apply only once per year.

Nonselective Weed Control on Noncrop Land

Princep 4L provides long-term weed control on industrial sites, highway medians and shoulders, railroad rights-of-way, lumberyards, petroleum tank farms, and in noncrop areas on farms, such as around buildings, equipment and fuel storage areas, along fences, roadsides, and ditches. Do not use this product for nonselective weed control on land to be cropped, near adjacent desirable trees, shrubs, or plants, or in greenhouses, as injury may occur. Do not use in areas accessible to livestock or allow livestock to graze treated weed forage. *Added*

Best results are obtained when Princep 4L is applied before weeds emerge. If weed growth has begun and removal is not feasible, include a contact or translocated herbicide in the spray. Aerial applications may be made where it is feasible. Use at least 1 gal. of water for each 1 qt. of Princep 4L for both ground and aerial applications. Use more water if practical.

Annual broadleaf and grass weeds (including barnyardgrass, cheat, crabgrass, lambsquarters, foxtail, ragweed, sunflower, and turkey mullein). Apply 4-8-10 qts. per acre.

Annual and perennial broadleaf and grass weeds (including quackgrass, bluegrass, redtop, burdock, Canada thistle, orchardgrass, purpletop, smooth brome, dogfennel, and plantain). Apply 2.5-5 qts. per acre.

Hard-to-kill perennial weeds (including bullthistle and sowthistle). Apply 5-10 gals. per acre.

For longer residual control in regions of high rainfall and along growing season, apply 5-10 gals. per acre.

Storage and Disposal

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited. Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable federal, state, or local procedures. Triple rinse container (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill or by incineration if allowed by state and local authorities.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Causes eye irritation. Harmful if inhaled or absorbed through the skin. Avoid breathing vapors or spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

First Aid: If in eyes, flush with plenty of water. Get medical attention on irritation persists. If on skin, wash with plenty of soap and water. If inhaled, remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

Environmental Hazards

Do not contaminate domestic livestock or irrigation water supplies or lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

AAtrex[®] trademark of CIBA-GEIGY for atrazine.

Complex[®] trademark of KALO Laboratories, Inc.

Fractane[®] trademark of Stauffer Chemical Co.

Lyik[®] trademark of CIBA-GEIGY for ametryn.

Princep[®] trademark of CIBA-GEIGY for simazine.

Nine-O[®] trademark of CIBA-GEIGY.

Surflan[®] trademark of E.I. du Pont de Nemours & Co. for oryzalin.

Sutan[®] trademark of Stauffer Chemical Co.

Unite[®] trademark of Hopkins Agricultural Chemical Co.

X-77[®] trademark of KALO Laboratories, Inc.

© 1982 CIBA-GEIGY Corporation

Agricultural Division
CIBA-GEIGY Corporation
Greensboro, North Carolina 27419
CGA 6132D

BEST DOCUMENT AVAILABLE

100-526

Princep

4L

Herbicide

For weed control in certain crops, ornamental plantings, on industrial sites, and around-the-farm.

Active ingredient:
 Simazine: 2-chloro-4,6-bis(ethylamino)-s-triazine 41.9%
Inert ingredients: 58.1%
Total: 100.0%

Princep 4L contains 4 lbs. active ingredient per gal.

Use entire contents at one time.
 © 1982 CIBA-GEIGY Corporation

2½ Gallons
 U.S. Standard Measure

Keep Out of Reach of Children. Caution

Precautionary Statements
Hazards to Humans and Domestic Animals
 Causes eye irritation. Harmful if inhaled or absorbed through the skin. Avoid breathing vapors or spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

First Aid: If in eyes, flush with plenty of water. Get medical attention if irritation persists. If on skin, wash with plenty of soap and water. If inhaled, remove victim to fresh air. If not breathing, give artificial respiration, preferably by mouth-to-mouth respiration.

Environmental Hazards
 Do not contaminate domestic livestock, or surface water supplies, streams, or rivers. Do not contaminate water by cleaning of equipment or disposal of wastes.

Princep® trademark of CIBA-GEIGY for simazine.

EPA Est. 100-LA-1
 EPA Reg. No. 100-526

See label booklet CGA 6L32D for complete directions for use.

Agricultural Division
 CIBA-GEIGY Corporation
 Greensboro, North Carolina 27419
 CGA 6L32D

CIBA-GEIGY

100 1982

Mr. Richard Conn
Ciba-Geigy Corp.
Agricultural Division
P. O. Box 11622
Greensboro, NC 27409

Dear Mr. Conn:

Subject: Various Amendments
Princep 4L
EPA Registration Number 100-526
Resubmission Dated October 15, 1982

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

In the biotype statement, add the word sequential so as to read xxx sequential combinations xxx.

Sincerely yours,

Richard P. Mountfort *RFM*
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (TS-767)

BEST DOCUMENT AVAILABLE

TS-767:RFMountfort:DCR-04133:WANG-0875C:lwj:Raven:479-2013:10/26/82

BEST DOCUMENT AVAILABLE