



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON D C 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Mr Montague Dixon
Syngenta Crop Protection LLC
PO Box 18300
Greensboro NC 27419 8300

DEC 17 2012

Subject Princep 4L Herbicide
EPA Registration Number 100 526
Application dated November 14 2012
Revisions to Environmental Hazards addition of New York restriction

Dear Mr Dixon

The labeling referred to above submitted in connection with registration under the Federal Insecticide Fungicide and Rodenticide Act (FIFRA) as amended is acceptable A stamped copy of your label is enclosed for your records This label supersedes all previously accepted labels

You must submit one (1) copy of the final printed label before you release the product for shipment Products released for shipment after eighteen (18) months from the date of this letter must bear the new revised label If these conditions are not complied with the registration will be subject to cancellation in accordance with FIFRA §6(e) Your release for shipment of the product constitutes acceptance of these conditions

If you have any questions please contact Hope Johnson at (703)305 5410 or at johnsonhope@epa.gov

Sincerely

Kable Bo Davis
Product Manager 25
Herbicide Branch
Registration Division (7505P)

[Booklet]

Sale use and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited

GROUP 5 HERBICIDE

Princep® 4L

Herbicide

For weed control in certain crops

Active Ingredient	
Simazine 2 chloro 4 6 bis(ethylamino) s triazine	41.9%
Other Ingredients	58.1%
Total	100.0%

Princep 4L contains 4 lbs active ingredient per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION


See additional precautionary statements and directions for use inside booklet

EPA Reg No 100 526

EPA Est

2.5 gallons
____ gallons
Net Contents

SCP 526A



FIRST AID	
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15 20 minutes • Call a poison control center or doctor for treatment advice
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 20 minutes • Remove contact lenses if present after the first 5 minutes then continue rinsing eye • Call a poison control center or doctor for treatment advice
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by a poison control center or doctor • Do not give anything by mouth to an unconscious person
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing call 911 or an ambulance then give artificial respiration preferably mouth to mouth if possible • Call a poison control center or doctor for further treatment advice
Have the product container or label with you when calling a poison control center or doctor or going for treatment	
HOT LINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill Leak Fire or Accident) Call 1 800 888 8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through the skin Causes moderate eye irritation Harmful if swallowed or inhaled Avoid breathing vapors or spray mist Avoid contact with skin eyes or clothing

Personal Protective Equipment (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

All mixers, loaders, applicators and other handlers must wear

- Long sleeved shirt and long pants
- Chemical resistant gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-5)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

Simazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Simazine has been found in ground water. Users are advised not to apply simazine to sand and loamy sand soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.

Product must not be mixed or loaded within 50 feet of intermittent streams and rivers, natural or impounded lakes and reservoirs. Product must not be applied within 66 feet of points where agricultural field (nurseries, Christmas tree plantings, shelterbelts, and turf grasses for sod farms) surface water runoff enters perennial or intermittent streams and rivers, or within 200 feet of natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66 foot buffer or setback from runoff entry points must be planted to crop or seeded with grass or other suitable crop.

Product must not be mixed or loaded or used within 50 feet of all wells, including abandoned wells, drainage wells, and sink holes. Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 ft. of any well are prohibited, unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rain water that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specified minimum containment capacities do not apply to vehicles when delivering pesticide to the mixing/loading sites.

Additional State imposed requirements regarding well head setbacks and operational area containment must be observed.

One of the following restrictions must be used in applying simazine to tile outletted terraced fields containing standpipes:

- Do not apply within 66 feet of standpipes in tile outletted terraced fields.
- Apply this product to the entire tile outletted terraced field and immediately incorporate it to a depth of 2-3 inches in the entire field.
- Apply this product to the entire tile outletted terraced field under a no-till practice only when a high crop residue management practice is practiced. High crop residue management is described as a crop management practice where little or no crop residue is removed from the field during and after crop harvest.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once unopened and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law, (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

ANY USE OF THIS PRODUCT IN AN AREA WHERE USE IS PROHIBITED IS A VIOLATION OF FEDERAL LAW. Before using this product, you must consult the Simazine Watershed Information Center (SWIC) to determine whether the use of this product is prohibited in your watershed. SWIC can be accessed through www.simazine-watershed.info or 1 888 365 2874. If the SWIC indicates that use of this product is prohibited in your watershed, you may return this product to your point of purchase or contact Syngenta Crop Protection, LLC for a refund.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

For Christmas trees, do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours. For all other crops and use patterns, do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves made of any waterproof material
- Shoes plus socks

NON AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170) The WPS applies when this product is used to produce agricultural plants on farms forests nurseries or greenhouses

Do not enter or allow others (including children or pets) to enter treated areas until sprays have dried

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR WEED CONTROL AND/OR ILLEGAL RESIDUES

Sale use and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited

Do not apply this product through any type of irrigation system

This product is not for use in formulating end use products for aquatic use

PRODUCT INFORMATION

Aerial application is prohibited

When tank mixing or sequentially applying simazine or products containing simazine the total pounds of simazine applied must not exceed the specific maximum rate per calendar year as noted in the use directions

Apply Princep 4L herbicide before weeds emerge or after removal of weed growth Princep 4L controls a wide variety of annual broadleaf and grass weeds when used at selective rates in agricultural crops

Where a range of application rates is given use the low rate on coarse textured soil and soil lower in organic matter use the high rate on fine textured soil and soil higher in organic matter

Since Princep 4L enters weeds mainly through their roots moisture is needed to move it into the root zone Very dry soil conditions following application may necessitate shallow cultivation or rotary hoeing

Princep 4L is nonflammable

ANNUAL WEEDS CONTROLLED

- | | |
|------------------------------------|---------------------------------------|
| alyssum | common lambsquarters |
| annual bluegrass | Nightshade |
| annual morningglory | pepperweed |
| annual ryegrass | pigweed |
| barnyardgrass (watergrass) | pineappleweed |
| burclover | prickly lettuce |
| carelessweed | common purslane |
| carpetweed | ragweed |
| common chickweed | rattail fescue |
| crabgrass (<i>Digitaria</i> spp) | redmaids |
| downy brome (cheatgrass) | Russian thistle |
| fall panicum | shepherdspurse |
| fiddleneck | signalgrass (<i>Brachiaria</i> spp) |
| filaree | silver hairgrass |
| fireweed | smartweed |
| fivehook bassia | spanishneedles |
| Florida s paintbrush | speedwell |
| Florida pusley | tansymustard |
| foxtails | wild mustard |
| goosegrass | wild oats |
| groundsel | witchgrass |
| henbit | yellow flower pepperweed |
| jungerice | |
| knawel (German moss) | |

RESISTANCE MANAGEMENT

Princep 4L is a **Group 5 Herbicide** (contains the active ingredient simazine)

Following many years of continuous use of this product and chemically related products biotypes of some of the weeds listed on this label have been reported which cannot be effectively controlled by this and related herbicides. Where this is known or suspected and weeds controlled by this product are expected to be present along with resistant biotypes we recommend the use of this product in registered combinations or in sequence with other registered herbicides which are not triazines i.e. which are not solely a Group 5 Herbicide. If only resistant biotypes are expected to be present use a registered herbicide which is not solely a Group 5 Herbicide. Consult with your State Agricultural Extension Service for specific recommendations.

PERENNIAL CROP REPLANTING

If replanting perennial crops do not apply Princep 4L in the year preceding planting these crops unless otherwise stated on this label or crop injury may occur

APPLICATION PROCEDURES

Ground application Use conventional ground sprayers equipped with nozzles that provide accurate and uniform application. Be certain that nozzles are uniformly spaced and the same size. Calibrate sprayer before use and recalibrate at the start of each season and when changing carriers. Unless otherwise specified use a minimum of 20 gals of spray mixture per acre

Use a pump with capacity to (1) maintain 35-40 psi at nozzles (2) provide sufficient agitation in tank to keep mixture in suspension and (3) to provide a minimum of 20% bypass at all times. Use centrifugal pumps which provide propeller shear action for dispersing and mixing this product. The pump should provide a minimum of 10 gals /minute/100 gal tank size circulated through a correctly positioned sparger tube or jets

Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on the suction side of the pump should be 16 mesh or coarser. Do not place a screen in the recirculation line. Use 50 mesh or coarser screens between the pump and boom and where required at the nozzles. Check nozzle manufacturer's recommendations

For band applications calculate amount to be applied per acre as follows

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

Spray Drift Management

A variety of factors including weather conditions (e.g. wind direction, wind speed, temperature, relative humidity) and method of application (e.g. ground chemigation) can influence pesticide drift. The applicator and grower must evaluate all factors and make appropriate adjustments when applying this product

Wind Speed

Do not apply at wind speeds greater than 10 mph

Droplet Size

Apply as a coarse or coarser spray (ASAE standard 572)

Release Height

For groundboom applications apply with a nozzle height no more than 4 feet above the ground or crop canopy

MIXING PROCEDURES

All Uses (1) Be sure sprayer is clean and not contaminated with any other materials or crop injury or sprayer clogging may result (2) Fill tank $\frac{1}{4}$ full with clean water nitrogen solution or complete fluid fertilizer (3) Start agitation (4) Be certain that the agitation system is working properly and creates a rippling or rolling action on the liquid surface (5) If any dry herbicides are being tank mixed add them to the tank and allow them to become thoroughly suspended (6) Then pour Princep 4L directly from jug into tank (7) Continue filling tank until 90% full Increase agitation if necessary to maintain surface action (8) Add other liquid tank mix herbicide(s) after this product is thoroughly suspended (9) Finish filling tank Maintain agitation to avoid separation of materials (10) Empty tank as completely as possible before refilling to prevent buildup of emulsifiable concentrate residue from possible tank mix herbicides (11) If an emulsifiable concentrate film starts to build up in tank drain it and clean with strong detergent solution or solvent (12) Clean sprayer thoroughly immediately after use by flushing system with water containing a detergent

Compatibility test To determine the tank mix compatibility of Princep 4L with liquid fertilizer crop oil spreaders or other recommended pesticides use this test method Nitrogen solutions or complete fluid fertilizers may replace all or part of the water in the spray in corn Since liquid fertilizers can vary even within the same analysis check compatibility each time before use Be especially careful when using **complete** suspension or fluid fertilizers as serious compatibility problems are more apt to occur Commercial application equipment may improve compatibility in some instances Check compatibility using this procedure

- 1 Add 1 pt of fertilizer to each of 2 one qt glass jars with tight lids
- 2 To **one** of the jars add $\frac{1}{4}$ tsp of a compatibility agent approved for this use ($\frac{1}{4}$ tsp is equivalent to 2 pts /100 gals spray) Cap and shake until mixed Examples of compatibility agents include Compex® and Unite®
- 3 To **both** jars add the appropriate amount of herbicide(s) If more than one herbicide is used add them separately with dry herbicides first flowables next and emulsifiable concentrates last After each addition cap and shake until

thoroughly mixed The appropriate amount of herbicides for this test follows (assuming a spray volume of 25 gals /A)

Dry herbicides For each pound per acre add 1.5 teaspoons to each jar Fluff up wettable powder products before measuring

Liquid herbicides For each pint per acre add 0.5 teaspoon to each jar

For a spray volume other than 25 gals /A change the teaspoons added to each jar as follows

$$\frac{25 \text{ gals } \times \text{ no. teaspoons given above}}{\text{desired gals spray volume per acre}} = \text{teaspoons to add to each jar}$$

- 4 After adding all ingredients put lids on and tighten Shake jars vigorously one minute Let the mixtures stand 15 minutes and then look for separation large flakes precipitates grease gels medium to heavy oily film on the jar or other signs of incompatibility Determine if the compatibility agent is needed in the spray mixture by comparing the 2 jars If either mixture separates but can be remixed readily the mixture can probably be sprayed as long as good agitation is used If the mixtures are incompatible test the following methods of improving compatibility (A) slurry the dry herbicide(s) in water before addition or (B) add 1/2 of the compatibility agent to the fertilizer and the other 1/2 to the emulsifiable concentrate or flowable herbicide before addition to the mixture If still incompatible do not use the materials mixed in the same spray tank

CROP USE DIRECTIONS

FRUIT AND NUT CROPS

Apply the spray to the orchard or vineyard floor avoiding contact with fruit foliage or stems Specified rates are based on broadcast treatment For band applications or spot applications around trees in fruit or nut plantings reduce the broadcast rate of Princep 4L and water per acre in proportion to the area actually sprayed

Precautions (All fruit and nut crops) To avoid crop injury (1) Apply only to orchards or groves where trees have been established 12 months or more unless specified differently (2) Make only one application per year except as noted otherwise (3) Do not use on gravelly sand or loamy sand soil (4) Immediately following application limit overhead sprinkler irrigation to 1/2 inch

Almonds Peaches and Nectarines (CA Only)

Princep 4L must be applied in a 2 to 4 ft band on each side of the tree row at a rate of 1 2 qts of Princep 4L per treated acre Apply before weeds emerge in late fall or early winter Weeds controlled by 1 qt of Princep 4L include burclover common chickweed wild mustard and shepherdspurse Do not apply more than 2 qts Princep 4L/A (2 lbs a 1 /per treated acre) per calendar year Do not apply more than once per calendar year

Precautions To avoid crop injury (1) Do not treat trees established in the grove less than 3 years (2) Do not treat the Mission (TX) variety of almonds (3) Do not apply to almond trees propagated on plum rootstocks (4) Do not replant almonds peaches or nectarines in treated soil for 12 months after treatment (5) Do not apply on soil with less than 1% organic matter (6) Do not treat areas where water will accumulate

Apples Pears and Tart Cherries

Apply 2 4 qts /A Princep 4L Do not apply more than 4 qts Princep 4L/A (4 lbs a 1 /A) per calendar year Do not apply more than once per calendar year For apples the pre harvest interval is 150 days

Avocados

CA and FL Only Apply 2 4 qts /A Princep 4L after final preparation of grove Do not apply more than 4 qts Princep 4L/A (4 lbs a 1 /A) per calendar year Do not apply more than once per calendar year

Blueberries and Caneberries (Blackberries, Boysenberries, Loganberries, and Raspberries)

Apply 2 4 qts /A Princep 4L in the spring or apply a split application of 2 qts /A Princep 4L in the spring plus 2 qts /A Princep 4L in the fall Apply in a minimum of 40 gals of water per acre

On plantings less than 6 months old use ½ the above rate

To control quackgrass apply 4 qts /A Princep 4L in the fall or split the application applying 2 qts /A Princep 4L in the fall plus 2 qts /A Princep 4L in the spring when quackgrass is actively growing

Do not apply more than 4 qts Princep 4L/A (4 lbs a 1 /A) per calendar year Do not apply more than twice per calendar year

Note Do not apply when fruit is present or illegal residues may result

Filberts

Apply 2.4 qts /A Princep 4L in the fall or apply a split application of 2 qts /A Princep 4L in the fall plus 2 qts /A Princep 4L in the spring. Do not apply more than 4 qts Princep 4L/A (4 lbs a.i./A) per calendar year. Do not apply more than twice per calendar year.

Precaution If trees are planted on a hillside excessive soil erosion may result from the elimination of weeds.

Note Do not apply when nuts are on the ground during the harvest period or illegal residues may result.

Grapes

Apply 2.4 qts /A Princep 4L any time between harvest and early spring. Do not apply more than 4 qts Princep 4L/A (4 lbs a.i./A) per calendar year. Do not apply more than once per calendar year.

Precaution Do not use in vineyards established less than 3 years or crop injury may occur.

Grapefruit Lemons and Oranges

AZ (Lemons and Oranges Only) Apply a split application of 1.6 qts /A Princep 4L in the spring plus 1.6 qts /A Princep 4L in the fall. Do not apply more than 3.2 qts Princep 4L/A (3.2 lbs a.i./A) per calendar year. Do not apply more than 1.6 qts Princep 4L per acre (1.6 lbs a.i./A) per application.

CA Apply 2.4 qts /A Princep 4L in a single application or apply 2 qts /A Princep 4L in the fall and 2 qts /A Princep 4L in the spring. Do not apply more than 4 qts Princep 4L/A (4 lbs a.i./A) per calendar year. Do not apply more than twice per calendar year.

Precaution Do not use in the Imperial Coachella or Palo Verde Valleys or crop injury may occur.

FL (Grapefruit and Oranges Only)

Broadcast Applications Apply up to 4 qts of Princep 4L/A per application to weed free soil once during the spring and/or once during the fall to control weeds expected to emerge during these periods. Apply prior to emergence of weeds or if weeds have emerged apply in tank mixture with a contact herbicide. Use caution to keep the treatment off the foliage fruit or trunk of citrus trees. Do not apply more than 4 qts Princep 4L/A (4 lbs a.i./A) per application. Do not apply more than 8 qts Princep 4L per acre (8 lbs a.i./A) per calendar year.

Banded Applications For control of difficult species such as balsam apple vine and spanish needles and partial control of honeyvine milkweed apply no more than 8 qts of Princep 4L (8 lbs a i/A) per treated acre as a single application in the spring using a 50% band application. The amount of Princep 4L per acre of field must not exceed 4 qts (4 lbs a i/A). Apply once in the spring growing season between January and April. Do not make a fall Princep 4L application if this treatment was used in the spring. When emerged weeds are present apply Princep 4L in tank mixture with a recommended contact herbicide. Follow all directions precautions limitations etc on the tank mix products.

TX (Grapefruit and Oranges Only) Apply 4 qts /A Princep 4L. Do not apply more than 4 qts of Princep 4L/A (4 lbs a i/A) per calendar year. Do not apply more than twice per calendar year.

Precautions (All areas) To avoid crop injury (1) Do not use in nurseries (2) Do not apply to bedded grapefruit lemons or oranges (except for FL grapefruit and oranges) (3) Do not apply to trees under stress from freeze damage for one year after the freeze (4) Do not exceed 4 qts /acre/year of Princep 4L on grove planted trees 1 year old or less or crop injury may occur **Note** Dark red grapefruit hybrids e.g. Star Ruby, have a higher risk of crop injury than non dark red types

Macadamia Nuts

Apply 2-4 qts Princep 4L in 50 gals of water per acre before harvest and just prior to weed emergence. Do not apply more than 4 qts Princep 4L/A (4 lbs a i/A) per calendar year. Do not apply more than once per calendar year.

Note Do not apply when nuts are on the ground during the harvest period as illegal residues may result.

Olives

Apply 2-4 qts /A Princep 4L following grove preparation in the fall. Repeat annually in midwinter. Do not apply more than 4 qts Princep 4L/A (4 lbs a i/A) per calendar year. Do not apply more than once per calendar year.

Peaches, Plums, and Sweet Cherries

Apply 1-6 qts /A Princep 4L. Apply in late fall to early spring prior to weed emergence. Do not apply more than 4 qts Princep 4L/A (4 lbs a i/A) per calendar year. Do not apply more than once per calendar year.

Precautions To avoid crop injury (1) Peaches use only in AR LA MO OK TX and states east of the Mississippi River. For CA see specific directions in the Almonds Peaches and Nectarines (CA only) section (2) Plums and sweet cherries use only in MO and states east of the Mississippi River except TN

Pecans

Apply 2.4 qts /A Princep 4L before weeds emerge in the spring. Do not apply more than 4 qts Princep 4L per acre (4 lbs a.i./A) per calendar year. Do not apply more than once per calendar year.

Precautions To avoid crop injury (1) Do not use west of the Pecos River in TX or in AZ, CA or NM. (2) Do not make applications to transplanted trees that have been established less than 2 years in the grove.

Note Do not apply when nuts are on the ground or illegal residues may result. Do not allow animals to graze treated areas.

Strawberries

OR and WA Only For control of chickweed, groundsel, mustard, and shepherdspurse, apply broadcast 1 qt/A Princep 4L. In fields where overhead irrigation is used to activate this product, apply after harvest at time of bed renovation. In fields where overhead irrigation is not available, apply during early October through November. Do not apply more than 1 qt Princep 4L/A (1 lb a.i./A) per calendar year. Do not apply more than once per calendar year.

Precautions To avoid crop injury (1) Make only one application per growing season. (2) Do not apply within 4 months after transplanting.

Walnuts

Apply 2.4 qts /A Princep 4L. Leveling and furrowing operations after application will lessen effectiveness of weed control. Do not apply more than 4 qts Princep 4L/A (4 lbs a.i./A) per calendar year. Do not apply more than once per calendar year.

Note Do not apply when nuts are on the ground or illegal residues may result.

Princep 4L Plus Bromacil 80W for Grapefruit and Oranges (FL Only)

Use in grapefruit and oranges in FL for control of balsam apple, black nightshade, carpetweed, crabgrass, cudweed, dayweed, Florida pusley, horseweed, pepperweed, pigweed, poorjoe, ragweed, rattlebox, spanish needles, and sandbur, and for partial control of bermudagrass, bahiagrass, pangolagrass, paragrass, and torpedograss. Apply 4 qts of Princep 4L plus 3.4 lbs of Bromacil 80W per acre beneath trees in a minimum of 40 gals of water per acre before or soon after weed growth begins. Use caution to keep the spray off the foliage, fruit, or trunk of citrus trees. When mixing, add Bromacil 80W slurry to water in spray tank, agitate thoroughly, then add Princep 4L and agitate thoroughly again. Use the lower rates for light weed infestations or all applications in bedded citrus areas. Use the higher rates for heavy weed infestations.

only in ridge grown citrus areas Temporary yellowing of citrus leaves may occur following treatment

Precautions To avoid crop injury (1) Do not use in nurseries or where trees are under stress from freeze damage for one year after the freeze (2) Do not use on soil with less than 1% organic matter or on poorly drained soil (3) Do not treat trees planted in irrigation furrows (4) Do not treat diseased trees such as those with foot rot (5) Do not use in groves interplanted with other trees or desirable plants nor in home grapefruit or orange plantings or in areas where roots of other valuable plants or trees may extend (6) Treated areas may be planted to citrus trees one year after application (7) Do not rotate to other crops within 2 years after application

Note Apply only once per year and avoid contact with foliage and fruit with spray or mist or illegal residues may result

Princep 4L Plus Gramoxone Inteon®

This tank mix is effective in the following fruit and nut crops for kill of existing vegetation and for residual control of the annual broadleaf and grass weeds claimed for Princep 4L applied alone This combination is also effective for top kill and suppression of perennial weeds In FL this mixture may be applied in spring or fall to emerged weeds

Use this tank mix on these crops

- | | |
|---|----------------------|
| almonds (CA only) | olives |
| apples | oranges ² |
| avocados (CA and FL ¹ only) | peaches ³ |
| cherries (tart and sweet ⁴) | pears |
| filberts | pecans |
| grapefruit (CA and TX only) | plums ⁴ |
| grapes | walnuts |
| lemons (AZ and CA only) | |
| macadamia nuts | |

Apply the Princep 4L rate given under the appropriate crop on this label plus the specified labeled rate of Gramoxone Inteon in 50 200 gals (30 50 gals for pecans) of water per acre to the orchard floor avoiding contact with fruit foliage or stems Add a nonionic surfactant such as X 77® at 0.5 pt/100 gals of spray Apply when weeds are succulent and new growth is 1.6 inches tall For mature woody weeds or difficult to control perennial weeds re treat or spot treat with Gramoxone Inteon if regrowth occurs

Add Princep 4L to the spray tank first (refer to **Mixing Procedures** section of this label) then add Gramoxone Inteon and add the surfactant last Provide constant agitation during mixing and application to keep the mixture in suspension

Precautions To avoid crop injury (1) Apply the tank mix only once per year (2) Use a shield for young trees or vines

Refer to the Princep 4L and Gramoxone Inteon labels for further directions specific weeds controlled and precautions and limitations on each crop

¹In avocados in FL this tank mix also controls balsamapple rattail amaranth and at the higher specified rate of each herbicide it suppresses coral vine

²In oranges in FL apply 4 qts of Princep 4L per acre per application Do not exceed 8 qts of Princep 4L per calendar year

³Limited to AR CA LA MO OK TX and states east of the Mississippi River As specified refer to Princep 4L sections **Almonds Peaches and Nectarines (CA only) or Peaches, Plums and Sweet Cherries** for rate of Princep 4L and other information

⁴Limited to MO and states east of the Mississippi River except TN

Tank Mixture with Roundup® or Touchdown® Brand Herbicides

This tank mixture is effective in grape vineyards and in the following bearing or nonbearing tree crops for control of existing vegetation and for residual control of the annual broadleaf and grass weeds claimed for this product applied alone This combination is also effective for partial control of perennial weeds contacted by the spray mixture during application

- | | |
|---|------------------------------|
| almonds** (CA only) | macadamia nuts** |
| apples | oranges*** (AZ CA FL and TX) |
| avocados* (CA and FL only) | peaches**** ¹ |
| cherries (tart and sweet**** ²) | pears* |
| filberts** | pecans** |
| grapefruits*** (CA FL and TX) | plums**** ² |
| grapes* | walnuts** |
| lemons* (AZ and CA) | |

Use the specified rate given elsewhere on this label for Princep 4L applied alone to the crop being treated Add to the spray tank the specified labeled rate of a Roundup or Touchdown brand product per acre depending on weeds present and their growth stage Also add an agriculturally approved nonionic surfactant at 0.5% by volume of spray solution Apply the mixture in 10-40 gals of water per acre as a postemergence spray to the weeds at the appropriate weed growth stage given on the Roundup label

Add Princep 4L to the spray tank first then add the Roundup or Touchdown brand product Provide constant agitation during mixing and application to keep the mixture in suspension Refer to **Application Procedures** section of this label for further directions

Precaution To avoid crop injury take extreme care to avoid contact of herbicide solution spray drift or mist with foliage or green bark of trunk branches suckers fruit or other parts of trees or vines Observe precautions on both the Princep 4L and Roundup or Touchdown labels for each crop involved

Always refer to the labels of both herbicides for further directions specific weeds controlled and precautions and limitations on each crop

*Allow a minimum of 14 days between last application and harvest

**Allow a minimum of 21 days between last application and harvest of these crops

***Allow a minimum of 1 day between last application and harvest of these crops

****Allow a minimum of 17 days between last application and harvest of these crops

¹Limited to AR CA LA MO OK TX and states east of the Mississippi River For CA see specific directions in the **Almonds Peaches and Nectarines (CA only)** section

²Plums and sweet cherries limited to MO and states east of the Mississippi River except TN

Tank Mixture with Solicam®

For improved control of such weeds as clover cutleaf eveningprimrose, dandelion henbit horseweed or marestail lambsquarters and puncturevine apply Princep 4L in tank mixture with Solicam DF on these crops

oranges
grapefruit
lemons

Apply 4 qts /A of Princep 4L plus 4.5 lbs of Solicam DF Herbicide (78.6% active ingredient) in 20-100 gals of water per acre Princep 4L + Solicam may be applied in tank mixture with Gramoxone Inteon or Roundup

Follow the labeled directions and restrictions of use on labels of the herbicides used in mixtures with Princep 4L

Precaution Keep Princep 4L + Solicam mixtures from contacting foliage fruits and stems of citrus trees during spraying

Princep 4L Plus Surflan®

Use this tank mix in the following crops for preemergence control of all weeds claimed on both labels

- | | |
|-------------|-----------------|
| almonds | lemons |
| apples | oranges |
| avocados | peaches |
| caneberries | pears |
| cherries | pecans |
| filberts | plums |
| grapes | English walnuts |
| grapefruit | |

Apply the Princep 4L rate given under the appropriate crop on this label plus 2 67 5 33 lbs of Surflan 75W or 2 4 qts of Surflan A S in 20 40 gals of water per acre Refer to the Surflan 75W or Surflan A S label for complete tank mix directions and use 0 8 qt of Princep 4L for each pound of Princep 80W Observe all precautions and limitations on the Princep 4L and Surflan labels

FIELD CROPS

Corn (Field and Sweet)

Apply a maximum of 2 qts Princep 4L per acre (2 0 lbs a /A) as a single preemergence application on soils that are not highly erodible or on highly erodible soils as defined by the Natural Resources Conservation Service if at least 30% of the soil is covered with plant residues If a second treatment is required following an earlier herbicide application the total simazine applied may not exceed 2 5 lbs a /A per calendar year

If the soil is highly erodible and covered with <30% plant residues apply a maximum of 1 6 lb a /acre as a single preemergence application For highly erodible soils and is covered with <30% plant residue do not apply more than one application of Princep 4L and not more than 1 6 lbs a /A (1 6 qt/A)

Pre grazing/pre harvest interval for field corn 60 days
Pre grazing/pre harvest interval for sweet corn 45 days

Nitrogen solution or complete liquid fertilizer may replace all or part of the water in the spray Determine the physical compatibility of Princep 4L with fertilizer before use Refer to the **Product Information** section of this label for a compatibility test procedure Do not apply after corn has emerged as there is danger of liquid fertilizer causing crop injury Use a minimum of 10 gals of spray mixture per acre

Preemergence Apply before weeds and corn emerge Use the specified rate as stated above

Preplant Broadcast in the spring after plowing at the specified rate as stated above Apply before during or after final seedbed preparation If soil is tilled or worked after application avoid deep incorporation of Princep 4L Best results will be obtained when Princep 4L is applied within 2 weeks before planting Under dry weather conditions preplant applications may give better weed control than preemergence If weeds develop particularly under relatively dry conditions a shallow cultivation will generally result in better weed control

Winter Annual Broadleaf Control – Preemergence Fall Application

For preemergence control of winter annual weeds such as common chickweed henbit shepherdspurse tansymustard wild mustard annual bluegrass downy brome and others broadcast 1 qt /A of Princep 4L after harvest of the preceding crop and prior to weed emergence on land to be planted to corn the following year A tillage operation may precede the application Do not apply to frozen ground If Princep 4L is used in the fall corn weed control program do not exceed 1.5 qts of Princep 4L preemergence in the spring

Precautions (1) Do not apply more than 2.5 qts of Princep 4L per acre of corn per calendar year (2) Do not plant any crop except corn until the following year or injury may occur (3) Following harvest plow and thoroughly till the soil in fall or spring to minimize possible injury to spring seeded rotational crops regardless of the rate used (4) Do not apply Princep 4L preplant incorporated for weed control in corn in the High Plains and Intermountain areas of the West (including central and western KS western NE western OK and the Panhandle of TX) where rainfall is sparse and erratic or where irrigation is required (5) In the High Plains and Intermountain areas of the West where rainfall is sparse and erratic or where irrigation is required use Princep 4L only when corn is to follow corn or when a crop of untreated corn is to precede other rotational crops (6) Injury may occur to soybeans planted in north central and northwest IA south central and southwest MN northeast NE southeast SD and other areas the year following application on soils having a calcareous surface layer (7) Do not plant sugar beets tobacco vegetables (including dry beans) spring seeded small grains or small seeded legumes and grasses the year following Princep 4L application or injury may occur

Note Do not graze treated areas or illegal residues may result

Tank Mixtures on Corn

AAtrex® Use Princep 4L in a preplant or preemergence tank mixture with AAtrex Nine O® or AAtrex 4L for control of many annual weeds including carpetweed crabgrass fall panicum foxtail lambsquarters morningglory pigweed ragweed and velvetleaf

When tank mixing or sequentially applying simazine and/or atrazine or products containing either active ingredient to corn the total pounds of simazine and/or atrazine applied must not exceed 2.5 lbs a.i./acre per calendar year

Refer to the AAtrex label for complete directions and use 1 qt of this product for each 1.1 lb of Princep Caliber 90. Also refer to the AAtrex label and the **Corn** section of this label for precautions and limitations

Gramoxone Inteon Use in a tank mixture where corn will be planted directly in a cover crop established sod or previous crop residues. This combination controls existing vegetation and provides residual control of the annual broadleaf and grass weeds listed under the **Product Information** section

Add Princep 4L to the spray tank mix thoroughly with water and then add Gramoxone Inteon and a nonionic surfactant. Provide constant agitation during mixing and application to keep the mixture in suspension

Apply up to 2 qts of Princep 4L plus the specified labeled rate of Gramoxone Inteon in 20-60 gals of water per acre as a broadcast spray either before or after planting but before corn emerges. Add a nonionic surfactant at the rate of 0.5 pt/100 gals of spray volume

For further information see product information, caution and warning statements, precautions and notes on the Princep 4L and the Gramoxone Inteon labels

NURSERIES, CHRISTMAS TREE PLANTINGS, AND SHELTERBELTS

Nurseries (see list below)

Apply 2-3 qts Princep 4L in at least 25 gals of water per acre in fall or spring. Do not apply more than 3 qts Princep 4L/A (3 lbs a.i./A) per calendar year. Do not apply more than once per calendar year

Precautions: To avoid plant injury do not apply for at least one year after transplanting

Christmas Tree Plantings and Shelterbelts (see list below)

Remove weed growth before application. Apply 2-4 qts Princep 4L in at least 25 gals of water per acre after transplanting. Use at the same rate for annual maintenance applications

For quackgrass control apply 4 qts/A Princep 4L in the fall or apply a split application of 2 qts/A Princep 4L in the fall plus 2 qts/A Princep 4L in early spring after quackgrass begins growth

Do not apply more than 4 qts Princep 4L/A (4 lbs a i /A) per calendar year Do not apply more than twice per calendar year

Precautions To avoid tree injury (1) Do not use Princep 4L on seedbeds or cuttingbeds (2) In CA OR and WA do not apply to Christmas trees or shelterbelts sooner than one year after transplanting In other areas do not apply to Christmas trees or shelterbelt transplants less than 2 years of age (3) Do not use until soil is firmly settled around roots (4) Do not apply more than once a year except as directed for quackgrass control

Apply Princep 4L to these species of trees and shrubs as directed above

Conifers

Deciduous Trees and Woody Ornamentals

- arborvitae
- Austrian pine
- Bishop pine*
- knobcone pine*
- lodgepole pine (shore pine)
- Monterey pine*
- Mugho pine
- red pine (Norway pine)
- Scotch pine
- white pine
- blue spruce
- Norway spruce
- red spruce
- white spruce
- balsam fir
- Douglas fir
- Fraser fir
- white fir
- hemlock
- juniper
- red cedar
- white cedar
- yew (*Taxus spp*)

- American elm
- Siberian elm
- barberry
- boxelder
- bush honeysuckle
- caragana
- cotoneaster
- dogwood
- eucalyptus
- holly (*Ilex spp*) (max 3 qts /A)
- honey locust
- Oregon grape (*Mahonia spp*)
- Pieris spp* (max 3 qts /A)
- red oak
- Russian olive
- oleander*
- palm (max 2 qts /A)*
- bottle brush*
- carob*

*For CA only

Pennant Magnum® Tank Mix Princep 4L may be tank mixed with Pennant Magnum Herbicide to control weeds in containerized field and liner* grown ornamentals and commercial landscapes See the Pennant Magnum label for ornamental species weeds controlled with this tank mixture and for other information

*Plants transplanted normally in rows in a nursery or similar area for further growth prior to transplanting to final growing place

Surflan Tank Mix On Christmas tree plantings use this tank mix for preemergence control of weeds listed on the Surflan 75W (or Surflan A S) and Princep 4L labels Use on field grown conifer species listed on the labels for each herbicide plus grand fir alpine fir Engelmann spruce black spruce Colorado blue spruce Coulter pine giant redwood and Veitchi fir

Broadcast the mixture as a directed spray to the soil surface or as an ovetop spray using 2.4 qts of Princep 4L and 2.67533 lbs of Surflan 75W (or 2.4 qts of Surflan A S) Apply in sufficient water per acre to uniformly treat the area Follow ovetop sprays with sprinkler irrigation to move the herbicide from leaf surfaces to the soil Remove weed growth before application Mix weed residues prunings or trash into the soil or remove them before treatment Soil should be in good tilth and free of clods at time of application Shallow cultivation (1.2 inches) after treatment will not reduce weed control Observe all precautions and limitations on the Princep 4L and Surflan labels

Note Length of weed control may be reduced when continuous wet soil conditions follow herbicide application

Precautions To avoid plant injury (1) Do not use on seedbeds or on unrooted cuttings (2) Do not use in greenhouses or other enclosed areas

TURFGRASSES FOR SOD (FL ONLY)

St Augustinegrass, Bermudagrass Centipedegrass and Zoysiagrass

Apply 2.4 qts /A of Princep 4L according to soil texture as indicated below This product must be watered in immediately after application if applied at a rate greater than 2 lbs a /A (2 qts /A of Princep 4L)

Muck or peat	4 qts	Old beds	within 2 days after lifting of sod
		New beds	3-4 days after sprigging or plugging
Sandy soil	2 qts	Old beds	within 2 days after lifting of sod
		New beds	7-10 days after sprigging or plugging

If weeds regrow apply an additional 2 qts /A or Princep 4L on muck or peat or 1 qt /A of Princep 4L on sandy soil

Precautions To avoid crop injury (1) Do not apply within 30 days prior to cutting or lifting (2) Do not apply in combination with surfactants or other spray additives (3) Use only on turfgrass reasonably free of infestations of insects nematodes and diseases (4) On newly sprigged turfgrass temporary slowing of growth may follow

application (5) Do not exceed 4 qts of Princep 4L per acre (4 lbs a /A) per application (6) Do not apply more than 2 applications per year (7) Do not exceed 6 qts Princep 4L per acre (6 lbs a /A) per calendar year

Note Do not graze or feed turf clippings to animals or illegal residues may result

TURFGRASSES FOR SOD (IN STATES EXCEPT FL)

St Augustinegrass Bermudagrass Centipedegrass and Zoysiagrass

Apply 1 2 qts /A of Princep 4L

Do not apply to established beds within 2 days after lifting of sod Do not apply to new beds before 7 10 days after sprigging or plugging

Do not apply more than 2 lbs a /A (2 qts of Princep 4L) per application and not more than two applications per calendar year Do not exceed 3 qts /A Princep 4L (3 lbs a /A) per calendar year

Precautions On newly sprigged turfgrass hybrid bermudagrass nondormant bermudagrass or nondormant zoysiagrass temporary slowing of growth and yellowing may occur following application To avoid turf injury (1) Use only on turfgrass reasonably free of infestations of insects nematodes and diseases (2) Do not use north of NC (except in the VA Coastal Plains) or on soils with a pH above 7.8 (3) Do not use on muck or alkaline soils (4) Do not apply over the rooting area of trees of ornamentals not listed on this label (5) Do not seed or overseed with desirable turfgrass within 4 months before or 6 months after treatment (6) Do not apply this product to newly seeded grasses until they have overwintered and have a well developed rhizome system (7) Do not exceed 2 qts /A of Princep 4L within 12 months of seeding grasses

Note Do not graze or feed turf clippings to animals or illegal residues may result

TURFGRASS FOR FAIRWAYS, LAWNS AND SIMILAR AREAS

Bermudagrass, Centipedegrass, St Augustinegrass, and Zoysiagrass

Apply Princep 4L after September 1 (after October 1 for annual bluegrass) before emergence of winter annual weeds Fall applications of Princep 4L will control annual bluegrass burclover lawn burweed common and mouseear chickweed corn speedwell henbit hop clover spurweed and parsley piert Princep 4L will also control these weeds soon after emergence For control of summer annual weeds listed in the **Product Information** section of this label apply Princep in late winter before the weeds emerge Apply in a minimum of 15 gals of water per acre

Irrigate with ½ inch of water if rainfall does not occur within 10 days after preemergence treatment

Where annual bluegrass is the major weed use 1 qt of Princep 4L per acre (22 ml or 0.75 fl oz per 1 000 sq ft) Use 1.2 qts /A of Princep 4L (22.44 ml or 0.7515 fl oz per 1 000 sq ft) for control of other weeds. However do not exceed 1 qt /A of Princep 4L per treatment on newly sprigged turfgrass or on hybrid bermudagrass

For control of summer annuals which emerge after the initial application apply an additional 1 qt /A of Princep 4L at least 30 days after the initial application but not after June 1. However do not make more than 2 applications per calendar year

Precautions On newly sprigged turfgrass hybrid bermudagrass nondormant bermudagrass or nondormant zoysiagrass temporary slowing of growth and yellowing may occur following application. To avoid turf injury (1) Use only on turfgrass reasonably free of infestations of insects nematodes and diseases (2) Do not use on golf greens (3) Do not use north of NC (except in the VA Coastal Plains) or on soils with a pH above 7.8 (4) Do not use on muck or alkaline soils (5) Do not apply over the rooting area of trees or ornamentals not listed on this label (6) Do not seed or overseed with desirable turfgrass within 4 months before or 6 months after treatment (7) Do not apply this product to newly seeded grasses until they have overwintered and have a well developed rhizome system (8) Do not exceed 2 qts /A of Princep 4L within 12 months of seeding grasses (9) Do not exceed 2 lbs a.i. /A (2 qts /A of Princep 4L) per application (10) Do not exceed 3 qts /A Princep 4L (3 lbs a.i. /A) per calendar year (11) Maximum of 2 applications per year

Note Do not graze or feed turf clippings to animals or illegal residues may result

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage or disposal

Pesticide Storage

Ground water contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material

Pesticide Disposal

Pesticide wastes may be toxic. Improper disposal of unused pesticide spray mixture or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods

Container Disposal [less than 5 gallons]

Non refillable container Do not reuse or refill this container Offer for recycling if available

Residue Removal [less than 5 gallons]

Triple rinse container (or equivalent) promptly after emptying Triple rinse as follows empty the remaining contents into application equipment of a mix tank and drain for 10 seconds after the flow begins to drip Fill the container ¼ full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begin to drip Repeat this procedure two more times

Container Disposal [Bulk/Mini Bulk]

Refillable container Refill this container with Princep 4L Herbicide only Do not reuse this container for any other purpose Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller

Residue Removal [Bulk/Mini Bulk]

To clean container before final disposal empty the remaining contents from this container into application equipment or mix tank Fill the container about 10 percent full with water Agitate vigorously or recirculate water with the pump for 2 minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat the rinsing procedure two more times

For minor spills leaks etc follow all precautions indicated on this label and clean up immediately Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes In the event of a major spill fire or other emergency call 1 800 888 8372 day or night

AAtrex® AAtrex® Nine O® Gramoxone Inteon® Princep® Solicam® Touchdown® and the Syngenta logo are trademarks of a Syngenta Group Company

Compex® trademark of KALO Agricultural Chemicals Inc

Roundup® trademark of Monsanto Company

Surflan® trademark of Dow AgroSciences

Unite® trademark of HACO Inc

X 77® trademark of Loveland Industries Inc

©201_ Syngenta

For non emergency (e g current product information) call
Syngenta Crop Protection at 1 800 334 9481

Manufactured for
Syngenta Crop Protection LLC
P O Box 18300
Greensboro North Carolina 27419 8300

SCP 526 MAS

[Container Label]

Sale use and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited

GROUP 5 HERBICIDE

Princep® 4L

Herbicide

For weed control in certain crops

Active Ingredient

Simazine 2 chloro 4 6 bis(ethylamino) s triazine 41 9%

Other Ingredients 58 1%

Total 100 0%

Princep 4L contains 4 lbs active ingredient per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Directions for Use in attached booklet

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170 Refer to supplemental labeling under Agricultural Use Requirements in the Directions for Use section for information about this standard

EPA Reg No 100 526

EPA Est

2 5 gallons

___ gallons

Net Contents

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful if absorbed through the skin Causes moderate eye irritation Harmful if swallowed or inhaled Avoid breathing vapors or spray mist Avoid contact with skin eyes or clothing

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15 20 minutes • Call a poison control center or doctor for treatment advice
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 20 minutes • Remove contact lenses if present after the first 5 minutes then continue rinsing eye • Call a poison control center or doctor for treatment advice
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by a poison control center or doctor • Do not give anything by mouth to an unconscious person
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing call 911 or an ambulance then give artificial respiration preferably mouth to mouth if possible • Call a poison control center or doctor for further treatment advice
Have the product container or label with you when calling a poison control center or doctor or going for treatment	
HOT LINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill Leak Fire or Accident) Call 1 800 888 8372	

Environmental Hazards

Simazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water Simazine has been found in ground water Users are advised not to apply simazine to sand and loamy sand soils where the water table (ground water) is close to the surface and where these soils are very permeable i.e well drained Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.

Product must not be mixed or loaded within 50 feet of intermittent streams and rivers, natural or impounded lakes and reservoirs. Product must not be applied within 66 feet of points where agricultural field (nurseries, Christmas tree plantings, shelterbelts, and turf grasses for sod farms) surface water runoff enters perennial or intermittent streams and rivers, or within 200 feet of natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66 foot buffer or setback from runoff entry points must be planted to crop or seeded with grass or other suitable crop.

Product must not be mixed or loaded or used within 50 feet of all wells, including abandoned wells, drainage wells, and sink holes.*

*For exceptions to this restriction, see the **Environmental Hazards** section of the **Precautionary Statements** in attached booklet.

Do not apply this product through any type of irrigation system.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

ANY USE OF THIS PRODUCT IN AN AREA WHERE USE IS PROHIBITED IS A VIOLATION OF FEDERAL LAW. Before using this product, you must consult the Simazine Watershed Information Center (SWIC) to determine whether the use of this product is prohibited in your watershed. SWIC can be accessed through www.simazine.watershed.info or 1 888 365 2874. If the SWIC indicates that use of this product is prohibited in your watershed, you may return this product to your point of purchase or contact Syngenta Crop Protection, Inc. for a refund.

Container Disposal [less than 5 gallons]

Non refillable container Do not reuse or refill this container Offer for recycling if available

Residue Removal [less than 5 gallons]

Triple rinse container (or equivalent) promptly after emptying Triple rinse as follows empty the remaining contents into application equipment of a mix tank and drain for 10 seconds after the flow begins to drip Fill the container 1/4 full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begin to drip Repeat this procedure two more times

Container Disposal [Bulk/Mini Bulk]

Refillable container Refill this container with Princep 4L Herbicide only Do not reuse this container for any other purpose Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller

Residue Removal [Bulk/Mini Bulk]

To clean container before final disposal empty the remaining contents from this container into application equipment or mix tank Fill the container about 10 percent full with water Agitate vigorously or recirculate water with the pump for 2 minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times

To be used in accordance with directions for use in Princep 4L label booklet See Princep 4L label booklet for pesticide disposal instructions

For minor spills leaks etc follow all precautions indicated on this label and clean up immediately Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes In the event of a major spill fire or other emergency call 1 800 888 8372 day or night

[Bulk]

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once unopened and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product which are beyond the control of SYNGENTA CROP PROTECTION LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law, (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

Princep® and the Syngenta logo trademarks of a Syngenta Group Company

©201_ Syngenta

Manufactured for
Syngenta Crop Protection LLC
P O Box 18300
Greensboro North Carolina 27419 8300

PRINCEP 4L 526 MAS 0711 AMEND NOV2012 – bb – 11 14 12

000100 00526 20121114 PRINCEP 4L AMEND NOV2012 pdf

◀ ◊
◊
◊ ◊ ◊
◊ ◊ ◊ ◊
◊ ◊
◊ ◊ ◊
◊ ◊