



100-528

2-4-77

1025

ACCEPTED
FEB 4 1977
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 100-528

Spectrum®



Kills
Ants
Brown dog ticks
Crickets
Cutworms
Earwigs
Grubs
Millipedes
Onion maggots
Sowbugs
Wireworms

For use in lawns, gardens, and around the home.

Ready-to-use. Contains 5% Diazinon™ insecticide.


CAUTION: Keep out of reach of children.

See additional caution statements on back of container.

and others
See Directions for Use

TREATS UP TO 1,000 SQ. FT.

One Pound
Net Weight



2012

Spectracide Crawling Insect Control Granules is designed especially for outdoor control of ants and other crawling insects. This product is effective as a preplant treatment for control of cutworms and wireworms in vegetable gardens. It also controls a wide range of insects in garden plantings and around the home, and offers excellent protection against major lawn insects on both grass and dichondra lawns. See Directions for Use for specific recommendations.

Directions for Use

Lawn Protection:

Spectracide Crawling Insect Control Granules controls the following lawn pests in both grass and dichondra lawns:

Ants	Crickets	Springtails
Armyworms	Cutworms	White grubs of:
Bermudagrass mites	Earwigs	Japanese beetle
Billbugs	Fleas	European chafer
Brown dog ticks	Leafhoppers	Southern chafer
Chiggers	Millipedes	
Chinch bugs	Sod webworms	
Clover mites	(Lawn moths)	
	Sowbugs	

How to Apply:

For most insects, sprinkle contents of this canister over a 500 sq. ft. area (10' x 50'). For heavy lawns, such as St. Augustine grass, spread contents evenly over 320 sq. ft. (10' x 32'). For billbug and white grub control, spread contents over 400 sq. ft. (10' x 40').

For best results, the lawn should be dry before application. To control surface feeding insects, armyworms and sod webworms (lawn moths), apply in late afternoon. Water lawn lightly after application. To control ants, billbugs, chinch bugs, cutworms, and white grubs, water lawn thoroughly after application.

When to Apply:

Apply when insects are present. To control ants, armyworms, Bermuda mites, chinch bugs, cutworms, leafhoppers and sod webworms (lawn moths), apply at 4-6 week intervals.

To control white grubs, apply any time between July and early October. To control billbugs, apply when activity is first observed, or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

Spot Treatment:

Ants: Apply 1 1/2 teaspoons over and around each ant hill.

Digger Wasps: Apply 1 1/2 teaspoons in and around each nest opening. Apply at dusk when wasps are not active.

Band Treatment Around House Foundation:

The following pests are commonly found around or near foundations of houses as well as in lawns. Certain of these pests may enter houses.

Ants	Fleas	Springtails
Brown dog ticks	Millipedes	(Collembola)
Clover mites	Sowbugs	
Crickets	(Pillbugs)	
Earwigs		

How to Apply:

Apply a 5-foot band of Spectracide Crawling Insect Control Granules next to the house foundation. Distribute granules uniformly, using a rate of 1-5 canister for each 100 sq. ft. area (5' x 20') treated.

When to Apply:

Apply when insects are present, especially in late spring or early fall. Repeat as necessary.

Garden Insect Control:

Spectracide Crawling Insect Control Granules controls certain garden insects on crops listed below. Apply just prior to planting and immediately mix thoroughly into soil to the specified depth, except where applied in seed furrow.

Cutworms (surface and subterranean)

Broccoli	Kale	Spinach
Cabbage	Lettuce	Summer Squash
Cantaloupes	Lima Beans	Swiss Chard
Cauliflower	Muskmelons	Tomatoes
Celery	Parsley	Watermelons
Collards	Peas	Winter Squash
Cucumbers	Peppers	
Endive (Escarole)	Potatoes	
	Snap Beans	

Apply 7 to 14 oz. per 500 sq. ft. and mix into soil to a depth of 2-3 inches for surface cutworms and 3-6 inches for subterranean cutworms. For potatoes, mix into soil 4-6 inches deep.

Mole Crickets

Broccoli	Kale	Tomatoes
Cabbage	Lettuce	Turnips
Cauliflower	Mustard	
Collards	Peppers	
Endive (Escarole)	Radishes	
	Strawberries	

Apply 4 oz. per 500 sq. ft. and mix into soil to a depth of 1-2 inches.

Wireworms

Broccoli	Carrrots	Collards
Cabbage	Cauliflower	Corn
Cantaloupes	Celery	Cucumbers

Endive (Escarole)	Onions	Summer Squash
Kale	Parsley	Sweet Potatoes
Lettuce	Peas	Swiss Chard
Lima Beans	Peppers	Tomatoes
Muskmelons	Snap Beans	Watermelons
	Spinach	Winter Squash

Apply 7 to 14 oz. per 500 sq. ft. and mix into soil to a depth of 4-8 inches.

Broccoli, Cabbage, Cauliflower

Root Maggots: Apply 7 to 11 oz. per 500 sq. ft. over entire seedbed before planting or transplanting and mix into soil to a depth of 3-4 inches.

Carrots

Carrot Rust Flies: Apply 7 oz. per 500 sq. ft. in seed furrow at planting time.

Onions

Onion Maggots: Apply 4 oz. per 500 sq. ft. in seed furrow at planting time.

Radishes

Root Maggots: Sprinkle lightly at the rate of 3-4 oz. per 500 linear feet of row in seed furrow at planting time.

Sweet Corn

Corn Rootworms: Apply 4-7 oz. per 500 sq. ft. in a band over the row at planting time.

Cutworms: Apply 7-11 oz. per 500 sq. ft. Mix into soil

to a depth of 1-2 inches for surface cutworms and 3-6 inches for subterranean cutworms.

European Corn Borers: Apply 4-7 oz. per 500 sq. ft. over corn so granules fall into whorls when 75% of the plants show first-generation larvae feeding, and again 7 days later, if needed. Treat for second-generation when there is an average of 1 egg mass per plant.

Fall Armyworms: Apply 4-7 oz. per 500 sq. ft. over corn so granules fall into whorls when larvae first appear, and again for control of second-generation armyworms, if necessary.

Southwestern Corn Borers: Apply 4-7 oz. per 500 sq. ft. over corn so granules fall into whorls when second-generation larvae first appear. Repeat application one week later.

*Note: Corn may be picked immediately following last application. Do not feed fodder to livestock for ten days following last application.

Wireworms: Apply 7-15 oz. per 500 sq. ft. in a broadcast application. Mix into soil to a depth of 4-8 inches.

Sweet Potatoes (North Carolina Only)

Wireworms: Apply 11 oz. per 500 sq. ft. over the top of sweet potato plant foliage when roots begin to enlarge.

Tomatoes

Vinegar Flies (Drasophila spp.): Apply 4 oz. per 500 sq. ft. over top of plants when flies first appear. Repeat at 7-10 day intervals, or as required. Tomatoes may be picked the same day.

Turnips

Root Maggots: Apply 4 oz. per 500 sq. ft. in seed furrow at planting time.

Caution: Harmful if swallowed. Avoid contamination of food and feed. Do not store near food or feed products. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. May be absorbed through skin. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

Fish and Wildlife Caution: This product is toxic to fish and wildlife. Keep out of lakes, streams, or ponds. Apply this product only as specified on this label.

Disposal: Do not reuse empty container. Wrap container and put in trash collection.

Active Ingredient:

O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	5%
Inert Ingredients:	95%
Total:	100%

Diazinon® Spectracide® and Spectrum® trademarks of CIBA-GEIGY

EPA Reg. No. 100-528-AA

Spectrum Home and Garden Products
Agricultural Division
CIBA-GEIGY Corporation
Greensboro, North Carolina 27409

