


100-528 10FJ
RM 15

D·z·n® 6000

TREATS UP TO 5000 SQUARE FEET

lawn & garden
insect
control

This product must not be used
on golf courses and sod farms.



D·z·n kills

Ants
Armyworms
Bermudagrass mites
Billbugs
Brown dog ticks
Chiggers
Chinch bugs
Crickets
Cutworms

Earwigs
Fire ants*
Fleas
Onion maggots
Root maggots
Sod webworms
White grubs of:
 . Japanese beetle
 . European chafer
 . Southern chafer
Wireworms
and others

*Aids in control.

See Directions for Use.

Ready-to-use Granules. Contains
5% diazinon insecticide.

Active Ingredient:

Diazinon: O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	5%
Inert Ingredients:	95%
Total:	100%


KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements
on back of bag.

EPA Est.

Ten Pounds
Net Weight



In Lawns
D·z·n kills:

- Ants
- Armyworms
- Bermudagrass mites
- Billbugs
- Brown dog ticks
- Chiggers
- Chinch bugs
- Clover mites
- Crickets
- Cutworms
- Digger wasps
- Earwigs
- Fleas
- Fire ants*
- Lawn moths
- Leafhoppers
- Millipedes
- Sod webworms
- Sowbugs
- Springtails
- White grubs of:
 - . European chafer
 - . Japanese beetle
 - . Southern chafer

In Gardens
D·z·n kills:

- Armyworms
- Carrot rust flies
- Corn borers
- Cutworms
- Mole crickets
- Onion maggots
- Root maggots
- Rootworms
- Vinegar flies
- Wireworms

*Aids in control.


Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Overall Lawn Application:

D·z·n 6000 control: the following lawn insects on both grass and dichondra lawns:

- Ants . Armyworms . Bermudagrass mites . Brown dog ticks .
- Chiggers . Chinch bugs . Clover mites . Crickets . Cutworms .
- Earwigs . Fleas . Lawn moths (sod webworms) . Leafhoppers .
- Millipedes . Sowbugs . Springtails (Collembola) .



Apply D·z·n 6000 with a fertilizer spreader. See spreader setting chart on side of bag. Apply when grass is dry. Water lawn lightly after application. Repeat as necessary.

For application with other types of equipment, apply at the rate of 2 lbs. per 1000 sq. ft. (an area 20' by 50') or 3.2 oz. per 100 sq. ft. (an area 10' by 10'). For heavy lawns such as St. Augustinegrass lawns, the rate should be increased to 3 lbs. per 1000 sq. ft. (or 4.8 oz. per 100 sq. ft.).

Billbugs; White grubs of Japanese beetle, European chafer, Southern chafer, Ataenius spretulus (dung beetle):
Apply 2.5 lbs. per 1000 sq. ft. of lawn. Water grass thoroughly after application. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary. For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in spring or early summer when damage is first noticed. Water grass thoroughly with 1/4-1/2 in. of water after application. Treat again in late summer or early fall if a second generation occurs.


Hyperodes weevils (New York only):
Apply 2 lbs. per 1000 sq. ft. of grass. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

Spot Treatment:

Ants: Apply 1 1/2 teaspoons of D·z·n 6000 over and around each ant hill.

Fire ants: To aid in the control of fire ants in lawns and other recreational areas, apply 1/2 cup over, and 2 ft. around, each fire ant mound. Immediately after application, water in thoroughly. A minimum of 1 gallon of water should be applied to each mound area. Reapply as necessary. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness.

Note: For best results, apply in cool weather, 65-80°F, or in early morning or late evening hours. Treat new mounds as they appear.



U.S.P.

Digger wasps: Apply 1 1/2 teaspoons of D·z·n 6000 in and around each nest opening. Apply at dusk when wasps are not active.

Band Treatment Around House Foundation:

The following pests are commonly found around or near foundations of houses as well as in lawns. Certain of these pests may enter houses.

- | | |
|-----------------|--------------------|
| Ants | Fleas |
| Brown dog ticks | Millipedes |
| Clover mites | Sowbugs (Pillbugs) |
| Crickets | Springtails |
| Earwigs | (Collembola) |

Apply a 5 foot band of D·z·n 6000 next to the house foundation. Distribute granules uniformly, using a rate of 3.2 oz. for each 100 sq. ft. of area (5' by 20') treated. Repeat as necessary.


Garden Insect Control:

D·z·n 6000 controls certain garden insects on crops listed below. Apply just prior to planting and immediately mix thoroughly into the soil to the specified depth, except where applied in seed furrow.

Cutworms (surface and subterranean)

- | | | |
|-------------|------------|---------------|
| Broccoli | Kale | Snap Beans |
| Cabbage | Lettuce | Spinach |
| Cantaloupes | Lima Beans | Summer Squash |
| Cauliflower | Muskmelons | Swiss Chard |
| Celery | Parsley | Tomatoes |
| Collards | Peas | Watermelons |
| Cucumbers | Peppers | Winter Squash |
| Endive | Potatoes | |
| (Escarole) | | |

Apply 7 to 14 oz. per 500 sq. ft. and mix into soil to a depth of 2-3 inches for surface cutworms and 3-6 inches for subterranean cutworms. For potatoes, mix into soil 4-6 inches deep.



Mole Crickets

Broccoli	Endive (Escarole)	Peppers
Cabbage	Kale	Radishes
Cauliflower	Lettuce	Strawberries
Collards	Mustard	Tomatoes
		Turnips

Apply 4 oz. per 500 sq. ft. and mix into soil to a depth of 1-2 inches.

Wireworms

Broccoli	Endive (Escarole)	Snap Beans
Cabbage	Kale	Spinach
Cantaloupes	Lettuce	Summer Squash
Carrots	Lima Beans	Sweet Potatoes
Cauliflower	Muskmelons	Swiss Chard
Celery	Onions	Tomatoes
Collards	Parsley	Watermelons
Corn	Peas	Winter Squash
Cucumbers	Peppers	

Apply 7 to 14 oz. per 500 sq. ft. and mix into soil to a depth of 4-8 inches.

Broccoli, Cabbage, Cauliflower

Root Maggots: Apply 7 to 11 oz. per 500 sq. ft. over entire seedbed before planting or transplanting and mix into soil to a depth of 3-4 inches.

Carrots

Carrot Rust Flies: Apply 7 oz. per 500 sq. ft. in seed furrow at planting time.

Onions


Onion Maggots: Apply 4 oz. per 500 sq. ft. in seed furrow at planting time.

Radishes

Root Maggots: Sprinkle lightly at the rate of 3-4 oz. per 500 linear feet of row in seed furrow at planting time.

Sweet Corn

Corn Rootworms: Apply 4-7 oz. per 500 sq. ft. in a band over the row at planting time.



Cutworms: Apply 7-11 oz. per 500 sq. ft. Mix into soil to a depth of 1-2 inches for surface cutworms and 3-6 inches for subterranean cutworms.

European Corn Borers: Apply 4-7 oz. per 500 sq. ft. over corn so granules fall into whorls when 75% of the plants show first-generation larvae feeding, and again 7 days later, if needed. Treat for second-generation when there is an average of 1 egg mass per plant.*

Fall Armyworms: Apply 4-7 oz. per 500 sq. ft. over corn so granules fall into whorls when larvae first appear, and again for control of second-generation armyworms, if necessary.*

Southwestern Corn Borers: Apply 4-7 oz. per 500 sq. ft. over corn so granules fall into whorls when second-generation larvae first appear. Repeat application one week later.*

*Note: Corn may be picked immediately following last application. Do not feed fodder to livestock for ten days following last application.

Wireworms: Apply 7-15 oz. per 500 sq. ft. in a broadcast application. Mix into soil to a depth of 4-8 inches.

Sweet Potatoes (North Carolina only)

Wireworms: Apply 11 oz. per 500 sq. ft. over the top of sweet potato plant foliage when roots begin to enlarge.


Tomatoes

Vinegar Flies (Drosophila spp.): Apply 4 oz. per 500 sq. ft. over top of plants when flies first appear. Repeat at 7-10 day intervals, or as required. Tomatoes may be picked the same day.

Turnips

Root Maggots: Apply 4 oz. per 500 sq. ft. in seed furrow at planting time.

NOTE: The spreader settings listed below are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust spreader setting if necessary, in order to assure correct rate of application.



SPREADER SETTING NO.
(per 1000 sq. ft.)

Spreader Make and Model	Insects listed below†		White grubs, Billbugs 2 1/2 lbs.
	Grass or Dichondra Lawn 2 lbs.	St. Augustine Lawn 3 lbs.	
Central Imperial Mark No. 1662-G	6 1/2	7 1/4	7
Central Suburbia Lawn Spreader - No. 1622-0	6 1/2	7 1/2	7
Cyclone Model B1 Rotary	4 1/2	5	4 3/4
Greenfield Model 105	6 1/2	7 1/4	6 3/4
ORTHO Drop Spreader Model 3000	4 1/2	5 1/2	5
ORTHO Broadcast Spreader Model 4000	4 1/2	7	6
J. C. Penney Model 6005	7 1/4	8	7 1/2
J. C. Penney Rotary Spreader	5	7	6
Scott Model 35	4	5	4 1/2
Sears Craftsman Model 671.19198	6	7	6 1/2

†Ants
Armyworms
Bermudagrass Mites
Brown Dog Ticks
Chiggers
Chinch Bugs

Clover Mites
Crickets
Cutworms
Earwigs
Fleas
Hyperodes Weevil
(New York only)

Lawn Moths
(Sod Webworms)
Leafhoppers...
Millipedes
Sowbugs
Springtails...
(Collembola)

Pof 5

Storage and Disposal

Storage: Store in original container. Store product in an area inaccessible to children and domestic animals. To prevent cross-contamination, do not store near fertilizers, herbicides, other insecticides, seeds, food, or feed.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse empty bag. Discard bag in trash.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. Avoid contamination of food and feed. Do not store near food or feed products. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. May be absorbed through skin. Keep children and pets off treated areas until this material is washed into the soil and grass is dry. In the event of an emergency, call (919) 292-7100 day or night.

Environmental Hazards

This product is toxic to fish and wildlife. Keep out of lakes, streams, or ponds. Apply this product only as specified on this label.

D·z·n® trademark of CIBA-GEIGY for diazinon

©1988 CIBA-GEIGY Corporation

EPA Reg. No. 100-528

Agricultural Division
CIBA-GEIGY Corporation
Greensboro, North Carolina 27419

CGA A1L86C 048

.....
.....
June 20, 1983
(Initial Copy of 10 Lb. Bag)
Revised July 19, 1983
(Incorporated crane fly-SLN
for Washington State.)
Revised March 18, 1987
(cancelled SLN), revised
Storage & Disposal
Revised April 6, 1988
added do not use on golf
courses and sod farms.