

16F18
Pm 15
100-528

D·z·n® 6000

TREATS UP TO 5000 SQUARE FEET

lawn & garden
insect
control

This product must not be used
on golf courses and sod farms.

ACCEPTED

JUN 21 1989

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 100-528

D·z·n kills

Ants
Armyworms
Bermudagrass mites
Billbugs
Brown dog ticks
Chiggers
Chinch bugs
Crickets
Cutworms

Earwigs
Fire ant *
Fleas
Onion maggots
Root maggots
Sod webworms
White grubs of:
 . Japanese beetle
 . European chafer
 . Southern chafer
Wireworms
and others

*Aids in control.

See Directions for Use.

Ready-to-use Granules. Contains
5% diazinon insecticide.

Active Ingredient:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	5%
Inert Ingredients:	95%
Total:	100%

KEEP OUT OF REACH OF CHILDREN.


CAUTION

See additional precautionary statements
on back of bag.

EPA Reg. No. 100-528

EPA Est.

Ten Pounds
Net Weight



DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of CIBA-GEIGY or the Seller. All such risks shall be assumed by the Buyer.

CIBA-GEIGY warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. CIBA-GEIGY makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall CIBA-GEIGY or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. CIBA-GEIGY and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of CIBA-GEIGY.

9 3 8 8

In Lawns
D·z·n kills:

Ants
Armyworms
Bermudagrass mites
Billbugs
Brown dog ticks
Chiggers
Chinch bugs
Clover mites
Crickets
Cutworms
Digger wasps
Earwigs
Fleas
Fire ants*
Lawn moths
Leafhoppers
Millipedes
Sod webworms
Sowbugs
Springtails
White grubs of:
 . European chafer
 . Japanese beetle
 . Southern chafer

*Aids in control.

In Gardens
D·z·n kills:

Armyworms
Carrot rust flies
Corn borers
Cutworms
Mole crickets
Onion maggots
Root maggots
Rootworms
Wireworms

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Overall Lawn Application:

D·z·n 6000 controls the following lawn insects on both grass and dichondra lawns:

Ants . Armyworms . Bermudagrass mites . Brown dog ticks
Chiggers . Chinch bugs . Clover mites . Crickets . Cutworms . . .
Earwigs . Fleas . Lawn moths (sod webworms) . Leafhoppers . .
Millipedes . Sowbugs . Springtails (Collembola)

Precautions: Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds. Avoid overlapping treated areas. If waterfowl, i.e., ducks, geese, can be expected to enter the treated area after treatment (except spot treatments), water lawn with at least 0.25 inch water immediately after application, however, stop watering before puddling occurs.

Apply D·z·n 6000 with a fertilizer spreader. See spreader setting chart below. Apply when grass is dry. Repeat as necessary.

For application with other types of equipment, apply at the rate of 2 lbs. per 1000 sq. ft. (an area 20' by 50') or 3.2 oz. per 100 sq. ft. (an area 10' by 10').

Billbugs, White grubs of Japanese beetle, European chafer, Southern chafer, Ataenius spretulus (dung beetle):

Apply 2 lbs. per 1000 sq. ft. of lawn. Water grass thoroughly after application. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary. For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in Spring or early Summer when damage is first noticed. Water grass thoroughly with 1/4-1/2 in. of water after application. Treat again in late Summer or early Fall if a second generation occurs.

Hyperodes weevils:


Apply 2 lbs. per 1000 sq. ft. of grass. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

Spot Treatment

Ants: Apply 1 1/2 teaspoons of D·z·n 6000 over and around each ant hill.

Fire ants: To aid in the control of fire ants in lawns and other recreational areas, apply 1/2 cup over, and 2 ft. around, each fire ant mound. Immediately after application, water in thoroughly. A minimum of 1 gallon of water should be applied to each mound area. Reapply as necessary. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness.

Note: For best results, apply in cool weather, 65-80°F, or in early morning or late evening hours. Treat new mounds as they appear.



Digger wasps: Apply 1 1/2 teaspoons of D·z·n 6000 in and around each nest opening. Apply at dusk when wasps are not active.

Band Treatment Around House Foundation:

The following pests are commonly found around or near foundations of houses as well as in lawns. Certain of these pests may enter houses.


Ants
Brown dog ticks
Clover mites
Crickets
Earwigs

Fleas
Millipedes
Sowbugs (Pillbugs)
Springtails
(Collembola)

Apply a 5 foot band of D·z·n 6000 next to the house foundation. Distribute granules uniformly, using a rate of 3.2 oz. for each 100 sq. ft. of area (5' by 20') treated. Repeat as necessary.

Vegetables - Insect Control

D·z·n 6000 controls certain garden insects on crops listed in the following tables. Apply uniformly as directed for each crop. When a range of rates is given, use the higher rate for heavier infestations. *only*



CROP	PEST	RATE/ACRE	REMARKS
Beans (Pole, Snap or Lima) (succulent only)	Cutworms *Garden Symphylans (Centi- pedes)	7-14 ozs. per 500 sq. ft.	Broadcast just prior to planting. Incorporate into the top 2-6 inches of soil. *The high rate of application will only help to suppress garden symphylan populations.
	Lesser Cornstalk Borers	3 1/2-7 ozs. per 500 sq. ft.	Apply in a 6-inch band centered over the row when plants are 1/2-1 inch tall.
	Wireworms	10 1/2-14 ozs. per sq. ft.	Broadcast just prior to planting. Incorporate into soil 4-8 inches. NOTE: Do not use on dried bean varieties, such as pintos, dried limas, navy beans, etc.
Beets, Red (Table)	Cutworms *Garden Symphylans (Centi- pedes),	7-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into soil 4-6 inches. *This application will only suppress garden symphylan populations. NOTE: This treatment will also control mole crickets and root maggots.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Broadcast to plant beds or fields 1-2 days prior to sowing of seed when insects are present, and incorporate into the top 1-2 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast prior to planting and incorporate into the top 4-8 inches of soil.
Broccoli	Root Maggots	7-10 1/2 ozs. per 500 sq. ft.	Broadcast before planting or transplanting. Incorporate into the top 3-4 inches of soil.
	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 4-6 inches of soil.

70F18

CROP	PEST	RATE/ACRE	REMARKS
Broccoli (Cont'd)	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Make applications to plant beds 1-2 days prior to planting when insects are present or expected and incorporate into the top 1-2 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 4-8 inches of soil.
Brussels Sprouts	Root Maggots	7-10 1/2 ozs. per 500 sq. ft.	Broadcast before planting or transplanting. Incorporate into the top 3-4 inches of soil.
	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 4-6 inches of soil.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Make applications to plant beds 1-2 days prior to planting when insects are present or expected and incorporate into the top 1-2 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just before planting and incorporate into the top 4-8 inches of soil.
Cabbage	Root Maggots	7-10 1/2 ozs. per 500 sq. ft.	Broadcast before planting or transplanting. Incorporate into the top 3-4 inches of soil.
	Cutworms *Symphylans (Centi- pedes)	7-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 4-6 inches of soil. *The high rate will only suppress garden symphylan populations.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Make applications to plant beds 1-2 days prior to planting when insects are present or expected and incorporate into the top 1-2 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 4-8 inches of soil.

CROP	PEST	RATE/ACRE	REMARKS
Carrots	Cutworms *Garden Symphylans (Centi- pedes)	7-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into soil 4-6 inches. *This application will only suppress garden symphylan populations. NOTE: This treatment will also control mole crickets and root maggots.
	Carrot Rust Flies	7 ozs. per 500 sq. ft.	Apply in the seed furrow at planting time.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Broadcast to plant beds or fields 1-2 days prior to sowing of seed when insects are present and incorporate into the top 1-2 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast prior to planting and incorporate into soil 4-8 inches.
Cauliflower	Root Maggots	7-10 1/2 ozs. per 500 sq. ft.	Broadcast before planting or transplanting. Incorporate into the top 3-4 inches of soil.
	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 4-6 inches of soil.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Make applications to plant beds 1-2 days prior to planting when insects are present or expected and incorporate into the top 1-2 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just before planting and incorporate into the top 4-8 inches of soil.
Celery	Cutworm	7-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the soil 2-4 inches to control cutworms.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the soil 4-8 inches to control wireworms.

CROP	PEST	RATE/ACRE	REMARKS
Collards	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 4-6 inches of soil.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Make applications to plant beds 1-2 days prior to planting when insects are present or expected and incorporate into the top 1-2 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just before planting and incorporate into the top 4-8 inches of soil.
Cucumber	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast just prior to planting. Incorporate into the top 2-6 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just prior to planting. Incorporate into the top 4-8 inches of soil.
Endive (Escarole)	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 2-4 inches of soil.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Apply to plant beds or fields 1-2 days before planting or transplanting when insects are present or expected. Incorporate into the top 1-2 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 4-8 inches of soil.
Kale	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 4-6 inches of soil.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 4-6 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just before planting and incorporate into the top 4-8 inches of soil.

CROP	PEST	RATE/ACRE	REMARKS
Lettuce	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 2-4 inches of soil.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Apply to plant beds or fields 1-2 days before planting or transplanting when insects are present or expected. Incorporate into the top 1-2 inches of soil.
	Wireworms, *Garden Symphylans (Centipedes)	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 4-8 inches of soil. *The high rate will only suppress garden symphylan populations.
Melons (Cantaloupes, Casabas, Crenshaws, Honeydew, Muskmelons, Persians, & Hybrids of these)	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast just prior to planting. Incorporate into the top 2-6 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just prior to planting. Incorporate into the top 4-8 inches of soil.
Mustard	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Apply to plant beds or fields 1-2 days before planting or transplanting when insects are present or expected. Incorporate into top 1-2 inches of soil.
Onions (green and bulb)	Onion Maggots	7-14 ozs. per 500 sq. ft.	Broadcast just before planting and mix into the top 3-4 inches of soil by disking, harrowing, or other suitable means. NOTE: Diazinon may not control organophosphate-resistant onion maggots.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just prior to the planting. Work into the top 4-8 inches of soil immediately after application.

110F 18

CROP	PEST	RATE/ACRE	REMARKS
Parsley	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 2-4 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 4-8 inches of soil.
Peas (succulent only)	Cutworms, *Garden Symphylans (Centipedes)	7-14 ozs. per 500 sq. ft.	Broadcast just prior to planting. Incorporate into the top 2-6 inches of soil. *The high rate of application will only help to suppress garden symphylan populations.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just prior to planting. Incorporate into the top 4-8 inches of soil. NOTE: Do not use on dried pea varieties such as dried blackeyed peas.
Peppers	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast just prior to planting. Incorporate into the top 2-6 inches of soil.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Apply when insects are present. Make application to plant beds or fields 1-2 days prior to sowing of seed or transplanting and incorporate into the top 1-2 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just prior to planting. Incorporate into the top 4-8 inches of soil.
Potatoes	Cutworms (Surface & Subterranean)	7-14 ozs. per 500 sq. ft.	Just before planting, broadcast and incorporate immediately into the top 4-6 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and incorporate into the top 4-6 inches of soil before planting. NOTE: Do not use on commercially-grown potatoes which will be hand-harvested.

CROP	PEST	RATE/ACRE	REMARKS
Radish	Root Maggots	5 1/2-7 ozs. per 500 sq. ft.	Apply in the seed furrow at planting (2 1/2-2 4/5 ozs. per 1,000 linear feet of row).
	*Garden Symphylans (Centipedes)	14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 4-6 inches of soil. *This application will only suppress garden symphylan populations. NOTE: This treatment will also control mole crickets, cutworms, and wireworms.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Make application to plant beds 1-2 days prior to planting when insects are present or expected and incorporate into the top 1-2 inches of soil.
Spinach	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 2-4 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 4-8 inches of soil.
Squash, Summer and Winter	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast just prior to planting. Incorporate into the top 2-6 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just prior to planting. Incorporate into the top 4-8 inches of soil.
Sweet Corn	Cutworms	7-14 ozs. per 500 sq. ft.	For surface cutworms, broadcast just prior to planting, or, when larvae first appear, and incorporate into soil 2-4 inches. Do not apply after layby. For subterranean cutworms, apply just prior to planting and incorporate into soil 3-6 inches.

130F18

CROP	PEST	RATE/ACRE	REMARKS
Sweet Corn (Cont'd)	Seed Corn Maggots	7-14 ozs. per 500 sq. ft.	Broadcast just prior to planting. Incorporate into soil 2 inches.
	Wireworms	7-14 ozs. per 500 sq. ft.	Broadcast prior to planting. Incorporate into soil 4-6 inches.
	European Corn Borers	3 1/2-7 ozs. per sq. ft.	Apply over the corn so granules fall into whorls when 75% of the plants show first generation borers feeding. Make a second application 7 days later if needed. Treat for second generation when there is an average of 1 egg mass per plant. Do not make more than 3 applications per season.
	Fall Armyworms	3 1/2-7 ozs. 500 sq. ft.	Apply over corn so granules fall into whorls when larvae first appear and again in 7 days if necessary. Do not make more than 3 applications per season.
	South- western Corn Borers	3 1/2-7 ozs. per sq. ft.	Apply over corn so granules fall into whorls when second generation larvae first appear. Repeat application one week later. Do not make more than 3 applications per season. NOTES: Corn may be picked immediately following last application. Do not feed fodder to livestock for 10 days following last application.
Sweet Potatoes	Wireworms, Flea Beetles, White Grubs, Southern Corn Rootworms	10 1/2 ozs. per 500 sq. ft.	Broadcast just prior to planting. Incorporate into the top 4-8 inches of soil or broadcast 10 1/2 ozs. per 500 sq. ft. over the tops of vines when roots begin to enlarge (Southeastern U.S.)

CROP	PEST	RATE/ACRE	REMARKS
Swiss Chard	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 4-8 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 2-4 inches of soil.
Tomatoes	Cutworms, *Garden Symphylans (Centi-pedes)	7-14 ozs. per 500 sq. ft.	Broadcast just prior to planting. Incorporate into the top 2-3 inches of soil. *The high rate of application will only help to suppress garden symphylan populations. NOTE: This treatment will also control mole crickets, root maggots, and wireworms.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Apply when insects are present. Make application to plant beds or field 1-2 days prior to sowing seed or transplanting and incorporate into the top 1-2 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just prior to planting. Incorporate into the top 4-8 inches of soil.
Turnip	*Garden Symphylans (Centi-pedes)	7-14 ozs. per 500 sq. ft.	Broadcast just before planting. Incorporate into the top 4-6 inches of soil. *The high rate of application will only suppress garden symphylan populations. NOTE: This treatment will also control mole crickets and root maggots.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Make application to plant beds 1-2 days prior to planting when insects are present or expected and incorporate into the top 1-2 inches of soil.

CROP	PEST	RATE/ACRE	REMARKS
Turnip (Cont'd)	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast prior to planting, and incorporate into the top 4-8 inches of soil.
Watermelon	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast just prior to planting. Incorporate into the top 2-6 inches of soil.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast just prior to planting. Incorporate into the top 4-8 inches of soil.

SPREADER SETTING NO.
(per 1000 sq. ft.)

Spreader Make and Model	Insects listed below†
	Grass or Dichondra Lawn 2 lbs.
Central Imperial Mark No. 1662-G	6 1/2
Central Suburbia Lawn Spreader - No. 1622-0	6 1/2
Cyclone Model B1 Rotary	4 1/2
Greenfield Model 105	6 1/2
ORTHO Drop Spreader Model 3000	4 1/2
ORTHO Broadcast Spreader Model 4000	4 1/2
J. C. Penney Model 6005	7 1/4
J. C. Penney Rotary Spreader	5
Scott Model 35	4
Sears Craftsman Model 671.19198	6

†Ants
Armyworms
Bermudagrass Mites
Brown Dog Ticks
Chiggers
Chinch Bugs

Clover Mites
Crickets
Cutworms
Earwigs
Fleas
Hyperodes Weevil

Lawn Moths
(Gad. Webworms)
Leafhoppers
Millipedes ..
Sowbugs
(Pill Bugs)
Springtails
(Collembola)

Storage and Disposal

Storage: Store in original container in a cool, dry, well-ventilated area. Store product in an area inaccessible to children and domestic animals. Do not store near fertilizers, herbicides, other insecticides, seeds, food, or feed.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse empty bag. Discard bag in trash.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust and contact with eyes, skin or clothing. Wash thoroughly after handling and before eating or smoking. Do not store near food or feed products. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Keep children and pets off treated areas until this material is watered into the soil and grass is dry.

Statement of Practical Treatment

If swallowed, remove victim to fresh air and apply artificial respiration if indicated.

If on skin, wash promptly with soap and water. Rinse thoroughly.

If in eyes, rinse eyes with plenty of water and call a physician immediately.

Note to Physician

This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Environmental Hazards

This product is toxic to fish, birds and other wildlife. Birds, including waterfowl feeding on treated areas may be killed. Do not apply directly to water or wetlands. Do not contaminate water by disposal of equipment wash waters. Apply this product only as specified on this label.

D-z-n® trademark of CIBA-GEIGY for diazinon

©1989 CIBA-GEIGY Corporation

EPA Reg. No. 100-528

Agricultural Division
CIBA-GEIGY Corporation
Greensboro, North Carolina 27419

CGA A1L86D

June 20, 1983
(Initial Copy of 10 Lb. Bag)
Revised July 19, 1983
(Incorporated crane fly-SLN
for Washington State.)
Revised March 18, 1987
(cancelled SLN), revised
Storage & Disposal
Revised April 6, 1988
Added do not use on golf
courses and sod farms.
Revised September 13, 1988
Changed chem. name, deleted
crops not supported under
May 1, 1987 EPA Data Call-
In, reduced rates on turf to
a max. of 4 lbs. a.i./A,
added avian precautions.
Revised November 4, 1988
specified sweet corn.
Revised May 17, 1989 -
Revised rates to concur with
Data Call-In residue program,
changed format, deleted
vinegar flies from tomatoes,
added several pests to some
crops, garden symphylan
suppression only, revised
Storage and Disposal, Env.
Hazards, Precautionary
Statements

(LABELS-D.4 - DZN/6000

