

RE-100-528

D·z·n®

6000

Lawn & Garden Insect Control

This product must not be used on golf courses and sod farms.

Ready-to-use Granules. Contains 5% diazinon insecticide

Treats up to 5000 square feet.

10 Pounds Net Weight

Active Ingredient:
Diazinon: O,O-Diethyl
O-(2-isopropyl-6-
methyl-4-pyrimidinyl)
phosphorothioate. .5%

Inert Ingredients: 95%

Total: 100%

KEEP OUT OF REACH
OF CHILDREN.

CAUTION

See additional precautionary statements on back of bag.

EPA Reg. No. 10-528

D·z·n kills:

- Ants, Armyworms
- Bermudagrass mites
- Billbugs
- Brown dog ticks
- Chiggers, Chinch bugs
- Crickets, Cutworms
- Earwigs, Fire ants*
- Fleas, Onion maggots
- Root maggots
- Sod webworms
- White grubs of:
 - Japanese beetle
 - European chafer
 - Southern chafer
- Wireworms
- ...and others

*Aids in control

See Directions for Use

ACCEPTED
MAR 26 1990

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 100-528

CIBA-GEIGY

In Lawns D-z-n kills

Ants	Crickets	Sod webworms
Armyworms	Cutworms	Sowbugs
Bermudagrass mites	Digger wasps	Springtails
Billbugs	Earwigs	White grubs of:
Brown dog ticks	Fire ants*	• Japanese beetle
Chiggers	Fleas	• European chafer
Chinch bugs	Lawn moths	• Southern chafer
Clover mites	Leafhoppers	
	Millipedes	* Aids in control

In Gardens D-z-n kills

Armyworms	Mole crickets	Rootworms
Corn rust flies	Onion maggots	Vinegar flies
Corn borers	Root maggots	Wireworms
Cutworms		

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of CIBA-GEIGY or the Seller. All such risks shall be assumed by the Buyer.

CIBA-GEIGY warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. **CIBA-GEIGY makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall CIBA-GEIGY or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product.** CIBA-GEIGY and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of CIBA-GEIGY.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Overall Lawn Application

D-z-n 6000 controls the following lawn insects on both grass and dichondra lawns:

Ants • Armyworms • Bermudagrass mites • Brown dog ticks • Chiggers • Chinch bugs • Clover mites • Crickets • Cutworms • Earwigs • Fleas • Lawn moths (sod webworms) • Leafhoppers • Millipedes • Sowbugs • Springtails (Collembola)

Precautions: Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds. Avoid overlapping treated areas. If waterfowl, i.e., ducks, geese, can be expected to enter the treated area after treatment (except spot treatments), water lawn with at least 1/4 inch water immediately after application, however, stop watering before puddling occurs.

Apply D-z-n 6000 with a fertilizer spreader. See spreader setting chart below. Apply when grass is dry. Repeat as necessary.

For application with other types of equipment, apply at the rate of 2 lbs. per 1,000 sq. ft. (an area 20' by 50') or 3.2 oz. per 100 sq. ft. (an area 10' by 10').


Billbugs; White grubs of Japanese beetle, European chafer, Southern chafer, *Ataenius spretulus* (dung beetle): Apply 2 lbs. per 1,000 sq. ft. of lawn. Water grass thoroughly after application. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary. For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in Spring or early Summer when damage is first noticed. Water grass thoroughly with 1/4-1/2 in. of water after application. Treat again in late Summer or early Fall if a second generation occurs.

Hyperodes weevils

Apply 2 lbs. per 1,000 sq. ft. of grass. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

LAWN SPREADER SETTING NO. (per 1000 sq. ft.)

Spreader Make and Model	Lawns/2 lbs.
Central Imperial Mark No. 1662-G	6 1/2
Central Suburbia Lawn Spreader No. 1622-0	6 1/2
Cyclone Model B1 Rotary	4 1/2
Greenfield Model 105	6 1/2
ORTHO Drop Spreader Model 3000	4 1/2
ORTHO Broadcast Spreader Model 4000	4 1/2
J. C. Penney Model 6005	7 1/4
J. C. Penney Rotary Spreader	5
Scott Precision Flow Spreader Model PF-4	4
Sears Craftsman Model 671.19198	6



D-z-n® 6000

Spot Treatment Lawns

Ants: Apply 1 1/2 teaspoons of D-z-n 6000 over and around each ant hill.

Fire ants: To aid in the control of fire ants in lawns and other recreational areas, apply 1/2 cup over, and 2 ft. around, each fire ant mound. Immediately after application, water in thoroughly. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. A minimum of 1 gallon of water should be applied to each mound area. Reapply as necessary.

Notes: For best results, apply in cool weather, 65-80°F, or in early morning or late evening hours. Works within 24 hours. No waiting for days or weeks. Treat new mounds as they appear.

Digger wasps: Apply 1 1/2 teaspoons of D-z-n 6000 in and around each nest opening. Apply at dusk when wasps are not active.

Band Treatment Around House Foundation

The following pests are commonly found around or near foundations of houses as well as in lawns. Certain of these pests may enter houses

Ants	Crickets	Millipedes
Brown dog ticks	Earwigs	Sowbugs (Pillbugs)
Clover mites	Fleas	Springtails (Collembola)

Apply a 5 foot band of D-z-n 6000 next to the house foundation. Distribute granules uniformly, using a rate of 3.2 oz. for each 100 sq. ft. of area (5' by 20') treated. Repeat as necessary

Vegetables - Insect Control


D-z-n 6000 controls certain garden insects on crops listed in the following tables. Apply uniformly as directed for each crop. When a range of rates is given, use the higher rate only for heavier infestations

Unless directions under "Timing/Special Directions" state otherwise, applications for control of pests listed below should be thoroughly mixed into the soil during or immediately after application.

Insect	Depth of Mixing or Incorporation into Soil
Surface cutworms*	2-3 inches
Subterranean cutworms*	3-6 inches
Mole crickets	1-2 inches
Root maggots	3-4 inches
Wireworms	4-8 inches

*If type of cutworm cannot be identified, mix into soil 4-6 inches deep.


CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Beans (Pole, Snap or Lima) (succulent only)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 oz. per 500 sq. ft.	
	Lesser Cornstalk Borers	3 1/2-7 oz. per 500 sq. ft.	Apply in a 6-inch band centered over the row when plants are 1/2-1 inch tall.
NOTE: Do not use on dried bean varieties, such as pintos, dried limas, navy beans, etc.			
Beets, Red (Table)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 oz. per 500 sq. ft.	
	Mole Crickets	3 1/2 oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Broccoli	Root Maggots	7-10 1/2 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 oz. per 500 sq. ft.	
	Mole Crickets	3 1/2 oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Brussels Sprouts	Root Maggots	7-10 1/2 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 oz. per 500 sq. ft.	
	Mole Crickets	3 1/2 oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Cabbage	Root Maggots	7-10 1/2 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 oz. per 500 sq. ft.	
	Mole Crickets	3 1/2 oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.



D-z-n® 6000

Carrots	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Cauliflower	Root Maggots	7-10½ oz. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 oz. per 500 sq. ft.	
	Wireworms	10½-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Celery	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Collards	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Cucumber	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Endive (Escarole)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Kale	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.

Lettuce	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Melons (Cantaloupes, Casabas, Crenshaws, Honeydew, Muskmelons, Persians, & Hybrids of these and watermelons)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Mustard	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Onions (green and bulb)	Onion Maggots	7-14 oz. per 500 sq. ft.	Broadcast just before planting and mix into the top 3-4 inches of soil by disking, harrowing, or other suitable means. NOTE: Diazinon may not control organo-phosphate-resistant onion maggots.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Parsley	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Peas (succulent only)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
NOTE: Do not use on dried pea varieties such as dried blackeyed peas			
Peppers	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Potatoes	Cutworms (Surface & Subterranean)	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
NOTE: Do not use on commercially-grown potatoes which will be hand harvested.			



D·z·n® 6000

Radishes	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting
Spinach	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting
	Wireworms	10½-14 oz. per 500 sq. ft.	
Squash, Summer and Winter	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Sweet Corn	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Seed Corn Maggots	7-14 oz. per 500 sq. ft.	Broadcast just before planting. Mix into soil 2 inches.
	European Corn Borers	3½-7 oz. per 500 sq. ft.	When 75% of the plants show first-generation borers, feeding, apply over the corn so granules fall into whorls. Make a second application 7 days later if needed. For second-generation borers, apply as above when there is an average of 1 egg mass per plant. Do not make more than 3 applications per season.
	Fall Armyworms	3½-7 oz. per 500 sq. ft.	Apply over corn so granules fall into whorls when larvae first appear. If necessary, application may be repeated twice at 7-day intervals. Do not make more than 3 applications per season.
	Southwestern Corn Borers	3½-7 oz. per 500 sq. ft.	
NOTES. Corn may be picked immediately after last application. Do not feed fodder to livestock for 10 days following last application			

Sweet Potatoes	Wireworms Flea Beetles	10½ oz. per 500 sq. ft.	Broadcast and mix into soil just before planting Southeastern U.S. - Broadcast 10½ oz. per 500 sq. ft. over the tops of vines when roots begin to enlarge, but do not mix into soil.
Swiss Chard	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Tomatoes	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Turnips	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.


Storage and Disposal

Storage: Store in original container in a cool, dry, well-ventilated area. Store product in an area inaccessible to children and domestic animals. Do not store near fertilizers, herbicides, other insecticides, seeds, food, or feed.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash

Container Disposal: Do not reuse empty bag. Discard bag in trash

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-888-8372 day or night.



Precautionary Statements**Hazards to Humans and Domestic Animals****CAUTION**

Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust and contact with eyes, skin or clothing. Wash thoroughly after handling and before eating or smoking. May cause contact sensitization following repeated contact with skin of susceptible individuals. Avoid repeated contact with skin. If sensitization reactions result, consult a physician. Do not store near food or feed products. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets on treated areas until granules have been watered into the soil and the grass or soil is dry.

Statement of Practical Treatment

If swallowed: call a physician immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If inhaled: remove victim to fresh air and apply artificial respiration, if indicated.

If on skin: wash promptly with soap and water. Rinse thoroughly.

If in eyes: rinse eyes with plenty of water and call a physician immediately.

Note to Physician

This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-AM is also antidotal and may be administered, but only in conjunction with atropine.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping granules. Where irrigation (watering) is recommended in the directions or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply a minimum of ¼ inch of water immediately after applying this product. Stop irrigation before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries.

Do not apply directly to water or wetlands (swamp, bogs, marshes and potholes). Runoff may be hazardous to aquatic organisms in neighboring areas.

Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

D-z-n® trademark of CIBA-GEIGY for diazinon.

© 1989 CIBA-GEIGY Corporation

Agricultural Division

CIBA-GEIGY Corporation

Greensboro, North Carolina 27419

CGA A1L86D

129

