

Spectracide® Brand Rose and Floral Spray

See Directions for Use for the list of specific insects and diseases controlled and the plants on which the spray may be used.


Active Ingredients:	
Pyrethrins	0.026%
*Piperonyl Butoxide, Technical	0.256%
Rotenone	0.128%
Other Cube® Extractives	0.238%
Captan (N-trichloromethylthio- 4-cyclohexene-1,2- dicarboximide)	0.504%
*2,4-Dinitro-6-octyl** phenyl crotonate	0.146%
*2,6-Dinitro-4-octyl** phenyl crotonate Nitrooctyl** phenols (principally dinitro)	0.010%
Petroleum Distillate	0.026%
Inert Ingredients:	98.666%
Total:	100.000%

CAUTION: Keep out of reach of children. See additional caution statements on back of container.

EPA Reg. No. 100-537

15 oz.

Net Weight



*Equivalent to 0.205% of (butylcarbityl) (6-propylpiperonyl) ether and to 0.051% of related compounds.

**A mixture of 1-methylheptyl, 1-ethylhexyl, and 1-propylpentyl isomers

†Karathane—Trademark of Rohm & Haas Company

Directions for Use

Hold can at least 18 inches from plant, with orifice directed away from you. Press actuator button to release in short bursts. Contact both upper and lower leaf surfaces. Avoid visibly wetting foliage. Spray before infestation becomes severe and repeat at 7-10 day intervals or as necessary.

This ornamental spray may be used on plants as a multipurpose spray in small gardens, and for spot treatment of incipient insect infestations to prevent spreading in large gardens.

FOR THE CONTROL OF GARDEN INSECTS--For use on roses, dahlias, asters and other ornamentals to kill Japanese beetles hit by spray, as well as aphids, exposed thrips and spider mites. Also effective against lace bugs, pavement ants, and clover mites.

This spray may be used on African violets, asters, azaleas, begonias, camelias, carnations, chrysanthemums, dahlias, delphiniums, dogwood, English ivy, euonymus, fuchsia, geraniums, crassula, ornamental grape vine, Kentia palm, laurel, marigolds, rhododendron, roses, rubber plants, snapdragons, stocks, Wandering Jew and zinnias.

Do not use on maidenhair fern, lantana or cyclamen.

As a Fungicide, spray before infestation occurs for the prevention of powdery mildew and black spot on roses. Spray at 5-7 day intervals while danger of fungus infection persists.

Caution:

Keep Out of Reach of Children.

Harmful if swallowed or inhaled. Avoid contamination of feed and foodstuffs. Avoid breathing or contact with eyes or skin. Wash thoroughly after use. May cause skin irritation or sensitization.

Warning:

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130° F may cause bursting. Never throw container into fire or incinerate.

Spectracide® trademark of CIBA-GEIGY

Home and Garden Products
Agricultural Division
CIBA-GEIGY Corporation
Greensboro, North Carolina 27409