

**Insecticide for control of certain insects
on certain vegetables, fruits, ornamentals**

Directions for Use

Soil Application: To control cutworms and wireworms prior to planting of the following vegetables:

<u>Beans</u>	<u>Cauliflower</u>	<u>Lettuce</u>	<u>Peas</u>	<u>Squash</u>
<u>Broccoli</u>	<u>Corn</u>	<u>Melons</u>	<u>Peppers</u>	<u>Tomatoes</u>
<u>Cabbage</u>	<u>Cucumbers</u>	<u>Onions</u>	<u>Potatoes</u>	
<u>Carrots</u>	<u>Endive</u>	<u>Parsnips</u>	<u>Spinach</u>	

Apply evenly as a broadcast treatment over the surface of the soil. Immediately after spreading, mix thoroughly with the soil to a depth of 6-8 inches. Apply 1/2 lb. per 225 sq. ft.

Foliage Application: Apply dust at first signs of insects. Repeat as necessary. For best results, dust in early morning or late evening when temperature is lower and a mist still. Apply dust so that all parts of plants are contacted, especially upper and lower leaf surfaces. Number in parentheses, (), indicates days required between last application and harvest.

Plants	Insects
<u>Beans</u> (7)	<u>Aphids, Mexican Bean Beetles, Leafhoppers</u>
<u>Peas</u> (0)	<u>Aphids</u>
<u>Broccoli</u> (5), <u>Cabbage</u> (7), <u>Cauliflower</u> (5)	<u>Aphids, Diamondback Moths, Imported Cabbageworm</u>
<u>Cucumbers</u> (7), <u>Melons: Cantaloupe, Muskmelon, Watermelon</u> (3), <u>Squash: Summer</u> (7), <u>Winter</u> (3)	<u>Aphids, Leafhoppers, Serpentine Leafminers, Mites, Thrips</u>
<u>Peppers</u> (5)	<u>Aphids, Serpentine Leafminers</u>
<u>Tomatoes</u> (3)	<u>Aphids, Dipterous Leafminers, Banded Cucumber Beetles, Fall Armyworm, Southern Armyworm</u>
<u>Sweet Corn</u> (2)	<u>Corn Earworm</u>
<u>Strawberries</u> (5)	<u>Aphids, Twospotted Mites</u>
<u>Roses, Flowers, Shrubs, Ornamentals</u>	<u>Aphids, Mites, Leafhoppers, Mealybugs, Whiteflies, Scale Crawlers, Lacebugs</u>

Note: Do not use on Ferns, Poinsettias, Hibiscus and Gardenias since injury may occur.

*Apply to silks as soon as they appear. Make applications at 3 to 4 day intervals as infestation warrants. Re-apply immediately after rain.

Active Ingredient:	
0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	4%
Inert Ingredients:	96%
Total:	100%



Caution:
Keep out of reach of children.
Harmful if swallowed. Avoid breathing dust. May be absorbed through skin. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and food-stuffs.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Apply this product only as specified on

EPA Reg. No. 100-538
EPA Est. No.

Diazinon - trademark of CIBA-GEIGY

Spectracide - trademark of CIBA-GEIGY



protects:
tomatoes
vegetables
fruit
roses
flowers
shrubs
as specified
in label directions

<u>Squash: Summer (7), Winter (3)</u>	<u>Aphids, Serpentine Leafminers</u>
<u>Peppers (5)</u>	<u>Aphids, Dipterous Leafminers, Banded Cucumber Beetles, Fall Armyworm, Southern Armyworm</u>
<u>Tomatoes (3)</u>	<u>Corn Earworm</u>
<u>Sweet Corn (2)</u>	<u>Aphids, Twospotted Mites</u>
<u>Strawberries (5)</u>	<u>Aphids, Mites, Leafhoppers, Mealybugs, Whiteflies, Scale Crawlers, Lacebugs</u>
<u>Roses, Flowers, Shrubs, Ornamentals</u>	

Note: Do not use on Ferns, Poinsettias, Hibiscus and Gardenias since injury may occur.

*Apply to silks as soon as they appear. Make applications at 3 to 4 day intervals as infestation warrants. Re-apply immediately after rain.

Active Ingredient:	
0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	4%
Inert Ingredients:	96%
Total:	100%

EPA Reg. No. 100-538
EPA Est. No.

Diazinon® trademark of CIBA-GEIGY

Spectracide® trademark of CIBA-GEIGY

Home and Garden Products
Agricultural Division
CIBA-GEIGY Corporation
Greensboro, North Carolina 27409


CGA A1L14 094

Caution:
Keep out of reach of children.
Harmful if swallowed. Avoid breathing dust. May be absorbed through skin. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and food-stuffs.


This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse container. Destroy when empty.



protects:
tomatoes
vegetables
fruit
roses
flowers
shrubs
as specified
in label directions



Comments Contains Diazinon®

CAUTION:
Keep out of reach of children.
See additional caution statements on back of container

Net Wt.
**One
Pound**

ACCEPTED WITH COMMENTS