


NOV 2 1974

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. 100-539

# Spectracide

## rose and flower spray



controls  
both insects  
and diseases:  
black spot  
powdery mildew  
aphids  
spider mites  
thrips  
leafhoppers

**Ready-To-Use  
Contains Diazinon Insecticide**

Spectracide Rose and Flower Spray is a highly effective, broad spectrum pressurized insecticide which kills many chewing and sucking insect pests and spider mites. It also controls the plant diseases of Black Spot, Powdery Mildew, Septoria Leaf Spot and Alternaria Leaf Spot. The product is non-phytotoxic to most plants when used as directed. For best results, read the following instructions carefully. Note: It is not necessary to wet plant foliage or the insect being sprayed. Excessive spraying only results in wasted material.

**Instructions for Use:**

Hold container upright. Point valve opening away from user and press down. Container should be held 18-24 inches from foliage. Spray with a slow sweeping motion, covering approximately 1 sq. ft. per second. Upper and lower leaf surfaces should be treated. Avoid wetting foliage. Care should be taken when spraying young plants with tender foliage and new growth. For best results spray when the air is still. Do not spray plants when temperature exceeds 90° F.

Over-all treatments may be made to smaller plants where spray mist will reach all parts of the plant. Larger plants may be spot treated using 1-second bursts to control exposed localized insect or mite infestations. Repeat as necessary.

Do not use on Ferns, Poinsettias, Hibiscus, Papayas, Pileas, Gardenias and Orchids since plant injury may occur.

Spectracide Rose & Flower Spray has no unpleasant odor.

**Insect and Mite Control**

Problem Areas	Pests	Notes
Azaleas	Aphids	Repeat as necessary to control exposed and active stages. Avoid wetting foliage. Hold container 18-24 inches from foliage. Spray with slow sweeping motion.
Camellias	Leafhoppers	
Carnations	Thrips	
Chrysanthemums	Spider Mites	
Gladioli		
Junipers		
Lilacs		
Rhododendron		
Roses		

**Plant Disease Control**

Problem Areas	Pests	Notes
Roses	Black Spot Powdery Mildew	For plant disease control begin when disease first appears and repeat at 7-10 day intervals. Avoid wetting foliage. Hold container 18-24 inches from foliage. Spray with slow sweeping motion.
Chrysanthemums	Septoria Leaf Spot	
Iris	Didymellina Leaf Spot	
Carnations	Rust Alternaria Leaf Spot	
Asters, Phlox	Alternaria Leaf Spot	
Chrysanthemums	Powdery Mildew	
Marigolds	Anthracnose	
Snap Dragons		
Zinnias		


**Active Ingredients:**  
 0,0-diethyl 0-(2-iso-propyl-6-methyl-4-pyrimidinyl) phosphorothioate ..... 0.500%  
 pyrethrins ..... 0.025%  
 technical piperonyl butoxide (equivalent to 0.200% [butyl-carbityl] [6-propyl-piperonyl] ether and 0.050% other related compounds) ..... 0.250%  
 Folpet (n-trichloromethyl thioph-

1.275%

**Caution:** Keep out of reach of children. Harmful if swallowed. Avoid contact with eyes. Avoid breathing spray mist and skin contact; wash thoroughly after using. Do not smoke while using. Do not use on household pets and humans. Avoid contamination of feed and foodstuffs. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. **Warning: Contents under pressure. Do**

NELEIN

# flower spray



**controls  
both insects  
and diseases:  
black spot  
powdery mildew  
aphids  
spider mites  
thrips  
leafhoppers**

as specified in label directions

## CAUTION:

Keep out of reach of children.  
See additional caution state-  
ments on back of container.

Net Wt.  
**15 oz.**

Problem Areas	Pests
Azaleas	Aphids
Camellias	Leafhoppers
Carnations	Thrips
Chrysanthemums	Spider Mites
Gladiah	
Junipers	
Lilacs	
Rhododendron	
Roses	

**Notes**  
Repeat as necessary to control  
exposed and active stages. Avoid  
wetting foliage. Hold container 18-24  
inches from foliage. Spray with slow  
sweeping motion.

### Plant Disease Control

Problem Areas	Pests	Notes
Roses	Black Spot Powdery Mildew	For plant disease control begin when disease first appears and repeat at 7-10 day intervals. Avoid wetting foliage. Hold container 18-24 inches from foliage. Spray with slow sweeping motion.
Chrysanthemums	Septoria Leaf Spot	
Iris	Didymellina Leaf Spot	
Carnations	Rust Alternaria Leaf Spot	
Asters, Phlox	Alternaria Leaf Spot	
Chrysanthemums	Powdery Mildew	
Marigolds	Anthracoise	
Snap Dragons		
Zinnias		

Active Ingredients:	1.275%
0,0-diethyl 0-(2-iso- propyl-6-methyl-4- pyrimidinyl) phos- phorothioate . . . . .	0.500%
pyrethrins . . . . .	0.025%
technical piperonyl butoxide (equivalent to 0.200% [butyl- carbityl] [6-propyl- piperonyl] ether and 0.050% other related compounds) . . . . .	0.250%
Folpet (n-trichloro- methyl thioph- thalimide) . . . . .	0.500%
Inert Ingredients:	98.725%
<b>Total:</b>	<b>100.000%</b>

**Caution:** Keep out of reach of children.  
Harmful if swallowed. Avoid contact  
with eyes. Avoid breathing spray mist  
and skin contact; wash thoroughly  
after using. Do not smoke while using.  
Do not use on household pets and  
humans. Avoid contamination of feed  
and foodstuffs.

This product is highly toxic to bees  
exposed to direct treatment or resi-  
dues on crops. Protective information  
may be obtained from your Coopera-  
tive Agricultural Extension Service

**Warning:** Contents under pressure. Do  
not puncture. Do not use or store near  
heat or open flame. Exposure to  
temperatures above 130°F may cause  
bursting. Never throw container into  
fire or incinerator.

Diazinon® and Spectracide® trade-  
marks of CIBA-GEIGY

Home and Garden Products  
Agricultural Division  
CIBA-GEIGY Corporation  
Greensboro, North Carolina 27409  
CGA A1L13

RECEIVED  
1977/12/16