

SPECIMEN LABEL

Net Weight: 5 lb

USDA Registration No. 8192-12

# Preforan<sup>®</sup>

(C-6989) 50 WP HERBICIDE

PREFORAN is a selective herbicide for preemergence weed control in soybeans to be harvested as a seed crop only. Do not use treated crop for food, feed or oil purposes. Do not graze treated areas or feed treated vegetation to livestock.

ACCEPTED

100-556

U.S. DEPARTMENT OF AGRICULTURE  
 OFFICE OF PEST CONTROL REGISTRATION  
 WASHINGTON, D.C. 20250  
 REG. NO. 8192-12

**Active Ingredient:**

p-nitrophenyl 2-nitro-4-(trifluoromethyl) phenyl ether. . . 50%

Inert Ingredients . . . . . 50%

Total 100%

**WARNING:** KEEP OUT OF REACH OF CHILDREN AND ANIMALS. May be harmful if swallowed. May cause eye irritation. Avoid contact of spray to eyes. Wear safety goggles when handling. In case of contact flush with water and get medical attention. Avoid inhaling dust. Avoid repeated or prolonged contact with the skin. Wear clean clothes and wash clothing before reusing. Wash hands with soap before eating, drinking, or smoking. Do not contaminate food or feed. Do not reuse container. Destroy empty containers by burning. Do not inhale smoke from burning containers.

**NOTE:** Toxic to fish. Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes and containers.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. In no case shall CIBA or the seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. CIBA and the seller offer this product, and the buyer and user accept it, subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of CIBA.

C I B A

**CIBA AGROCHEMICAL COMPANY**  
 Division of CIBA Corporation  
 556 Morris Ave., Summit, New Jersey 07091

SPECIMEN LABEL

**GENERAL INFORMATION**

PREFORAN is effective for the control of certain annual grass and broadleaf weeds when applied preemergence to soybeans. PREFORAN may be applied as a broadcast or as a band treatment to the surface of the soil at planting or at any time preemergence to the crops and weeds.

Use only one application to the crop during any one year. Do not follow the treated crop within the same year with susceptible crops, such as cucurbits or tomatoes.

Do not soil incorporate PREFORAN (C-6989) 50 WP Herbicide.

**WEED CONTROL**

PREFORAN is effective for controlling certain annual grass and broadleaf weeds in soybeans, such as:

**Annual Grass Weeds**

Brachiaria  
Crabgrass  
Giant Foxtail  
Goosegrass  
Green Foxtail  
Yellow Foxtail

**Annual Broadleaf Weeds**

Carpetweed  
Common Ragweed  
Lambsquarters  
Pigweeds  
Purslane  
Smartweeds

**APPLICATION DIRECTIONS**

**Broadcast Treatment** Apply 7.5 to 9.0 pounds of PREFORAN 50 WP in 20 to 40 gallons of water per acre to the surface of the soil with conventional power sprayers. PREFORAN may be applied at the time of planting soybeans to be harvested as a seed crop only, or at any time preemergence to the crop and weeds.

**Band Treatment** - A band treatment of PREFORAN is made in the same manner as a broadcast treatment. The following formula may be used to calculate the amount of PREFORAN per acre for band treatment:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Rate per acre for broadcast treatment} = \text{Amount needed for band treatment}$$

**MIXING INSTRUCTIONS** MAKE CERTAIN SPRAY EQUIPMENT HAS BEEN THOROUGHLY CLEANED BEFORE USING THIS PRODUCT. PREFORAN 50 WP should be thoroughly mixed as a paste or thick slurry in a separate container before adding to the spray tank. The spray tank should contain 1/4 to 1/2 the total amount of water to be used before adding the slurry. Avoid leaving spray mixture in tank without constant agitation. If bypass agitation is used, it should terminate at the bottom of the tank to minimize foaming. Screen openings should be 50 mesh. Use Tee Jet nozzle tips 8002 to 8006 or equivalent for broadcast treatments or 8002E to 8006E or equivalent for band applications.

Calibrate sprayer accurately, and operate at a constant speed. If sprayer is slowed or stopped, shut off spray, or overdosage may result in crop injury.

CIBA AGROCHEMICAL COMPANY Division of CIBA Corporation, 550, Morris Avenue, Summit, N.J. 07991

4/72