

DRAFT LABEL

**RESTRICTED USE PESTICIDE**

FOR RETAIL SALE TO AND USE ONLY BY  
 CERTIFIED APPLICATORS OR PERSONS  
 UNDER THEIR DIRECT SUPERVISION, AND ONLY  
 FOR THOSE USES COVERED BY THE  
 CERTIFIED APPLICATORS CERTIFICATION

Curacron® 6E

Insecticide - Miticide

For insect and mite control in cotton

Active Ingredient:

Profenofos: Q-(4-bromo-2-chlorophenyl) Q-ethyl S-	
<u>propyl phosphorothioate</u>	59.7%
<u>Inert Ingredients<sup>1</sup>:</u>	40.3%
<u>Total:</u>	100.0%

<sup>1</sup>contains petroleum distillates

2 1/2 Gallons  
U. S Standard Measure

Curacron 6E contains 6 lbs. active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN

WARNING

See additional precautionary statements and directions for use inside booklet.

Curacron® trademark of CIBA-GEIGY for profenofos

EPA Reg. No. 100-599  
EPA Est. 100-AL-1

©1991 CIBA-GEIGY Corporation


Agricultural Division  
CIBA-GEIGY Corporation  
Greensboro, North Carolina 27419

CGA 55L6 (DRAFT) 08/5/91

ACCEPTED

SEP 10 1991

100-599



DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

**IMPORTANT:** Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.


---

Conditions of Sale and Warranty

---

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of CIBA-GEIGY or the Seller. All such risks shall be assumed by the Buyer.

CIBA-GEIGY warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. CIBA-GEIGY makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall CIBA-GEIGY or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. CIBA-GEIGY and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of CIBA-GEIGY.



General Information

Curacron 6E is an emulsifiable insecticide-miticide which is effective for control of cotton bollworm, tobacco budworm, certain other insects, and mites on cotton.

Best results can be expected from fields under Pest Management/Scouting programs.

Note: When using Curacron 6E as a larvicide, best results will be obtained when application is timed to control first to third instar larvae.

Mixing Instructions

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Agitate thoroughly before and during application. Good bypass agitation is sufficient to keep Curacron 6E in emulsion.

Curacron 6E Alone: Add 2/3 of the required amount of water to the spray tank with the agitator running, then add the desired amount of Curacron 6E. Continue agitation while adding the remainder of the water.

Curacron 6E + Tank Mixtures: Add 2/3 of the required amount of water to the spray tank. With the agitator running if possible, add the desired amount of the other products recommended on this label. Then add the desired amount of Curacron 6E. Continue agitation while adding the remainder of water.

Note: Do not let spray mixture stand overnight in the spray tank. Rinse spray tank at the end of the day.

Chemigation: Do not apply this product through any type of irrigation system.


DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW ALL DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR CONTROL, AND/OR ILLEGAL RESIDUES.

Cotton

Apply Curacron 6E at the specified rates below in a minimum of 5 gallons of water per acre with ground equipment or in a minimum



of 1 gallon of water per acre by air. For better coverage of late season "rank" cotton increase gallonage. To avoid spray drift, do not apply under windy conditions. Avoid spray overlap because crop injury may result.

If the need to control boll weevil becomes apparent while using Curacron, Guthion® or methyl parathion may be tank mixed with Curacron. Refer to the Guthion or methyl parathion labels for directions, limitations, and precautions.

Aphids (All cotton growing states): Apply 2/3 pt. of Curacron 6E per acre at 5-7 day intervals as needed starting when economic thresholds are reached. Economic thresholds will vary by locality. Consult your pest control advisor or cotton consultant to determine economic thresholds in your area.

Plant bugs/lygus bugs\* (All cotton growing states): Apply 1/3-2/3 pt. of Curacron 6E per acre at 5-7 day intervals as needed starting when economic thresholds are reached. Economic thresholds will vary by locality. Consult your pest control advisor or cotton consultant to determine economic thresholds in your area.

\*In Arizona and California, Curacron 6E offers suppression of lygus bugs.

Cotton bollworm, tobacco budworm (All cotton growing states except Arizona and California):

Contact Ovicide Activity: For contact ovicidal activity, apply 1/6-1/3 pt. of Curacron 6E per acre when egg lay exceeds thresholds. Continue at 5-7 day intervals as needed. If larvae survive, use higher rates listed below.

Larvae Control: To control larvae only, apply 2/3 to 1 pt. of Curacron 6E for light to moderate infestations. Apply 1 to 1 1/3 pts. Curacron 6E for moderate to heavy infestations. Apply at 5-7 day intervals as needed beginning when infestations first appear.

Cotton bollworm, tobacco budworm (Arizona and California): Apply Curacron 6E at 5-7 day intervals as needed beginning when infestations first appear. Use 2/3 to 1 pt. per acre for light to moderate infestations and 1-1 1/3 pts. per acre for moderate to heavy infestations.

Cotton leafperforator, beet armyworm (All cotton growing states): Apply Curacron 6E at 5-7 day intervals as needed starting when larvae first appear. Use 2/3 - 1 pt. per acre for light to moderate infestations and 1-1 1/3 pts. per acre for moderate to heavy infestations.

Fall armyworm (All cotton growing states): Apply 1-1 1/3 pts. of Curacron 6E per acre at 5-7 day intervals as needed starting

when larvae first appear. For best results, application should be made to young larvae still feeding on leaf surfaces and before migration to blooms and bolls has occurred.

Mites (All cotton growing states): Apply 2/3 - 1 1/3 pts. of Curacron 6E per acre at 5-7 day intervals as needed starting when mites first appear.

Whitefly suppression (All cotton growing states): To aid in the reduction of whitefly populations, apply 1-1 1/3 pts. of Curacron 6E at 5-7 day intervals as needed starting when whiteflies first appear.

Curacron 6E + Tank Mixtures (All cotton growing states): To broaden the spectrum of insect control and make a more useful spray application, if multiple pests occur on cotton, it may be advantageous to tank mix Curacron 6E with other EPA-registered insecticides.

Curacron 6E has been found to be compatible in tank mixtures with the products listed below. When using Curacron 6E in tank mixtures, observe all limitations, precautions and rate recommendations which appear on the labels for these products: Ambush®, Ammo®, Baythroid® 2, Capture® 2EC, Comite®, Cymbush®, Dipel® 2X, Guthion®, Karate®, Lannate® L, Larvin® 3.2, methomyl, methyl parathion, Orthene® 75S, Pay-Off®, Pounce®, Pydrin® and Thiodan® 3EC.

Note: Applications to cotton may cause spotting or reddening of leaves.

Note: To avoid illegal residues: 1) Do not apply more than 4 qts. of Curacron 6E per acre per season. 2) Do not apply Curacron 6E within 14 days of harvest. 3) Do not allow livestock to graze treated cotton. 4) Do not feed gin trash or foliage from treated cotton plants to livestock.

Fields treated with Curacron may be rotated to other crops.

#### Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container disposal: Triple rinse (or equivalent). Then puncture container and dispose of in a sanitary landfill, or incinerate. The container may be disposed of by open burning if allowed by state and local authorities. Stay out of smoke from burning container.

Emergency Response

In the event of a major spill, fire, or other emergency, call 1-800-888-8372.

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING

May be fatal if absorbed through skin. Harmful if swallowed or inhaled. Causes moderate eye irritation. Do not get in eyes, on skin, or on clothing. Avoid breathing vapors or spray mist. Wear protective clothing and rubber gloves when handling. Wash thoroughly with soap and water before eating or using tobacco. Remove contaminated clothing and wash before reuse.

Statement of Practical Treatment

If in eyes: immediately flush with plenty of water. Get medical attention.

If on skin: wash with plenty of soap and water. Remove contaminated clothing. Get medical attention.

If swallowed: call a physician or poison control center. May present aspiration hazard. DO NOT induce vomiting. Drink large quantities of water. Get medical attention immediately.

If inhaled: remove victim to fresh air.

Note to Physician: If symptoms of cholinergic poisoning occur, atropine plus 2-PAM may be helpful. If swallowed, lavage stomach, taking care to avoid aspiration of stomach contents into the lungs.

Environmental Hazards

Due to the hazard to endangered fish, the application of this product is prohibited in Reeves County, Texas and within one mile of the Dexter National Fish Hatchery in New Mexico. This pesticide is toxic to fish and wildlife. Use with care when applying to areas frequented by wildlife or adjacent to any body of water. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply with aircraft within 300 feet

upwind of impounded water. Do not apply when weather conditions favor drift from treated areas. Do not apply with aircraft when wind speed is greater than 10 mph. Apply this pesticide only as specified on this label. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply this product or allow it to drift to crops or weeds on which bees are foraging. Additional information may be obtained from your Cooperative Agricultural Extension Service.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

Ambush®, Cymbush and Karate® trademarks of ICI Americas, Inc.

Ammo®, Capture®, Pounce® and Thiodan® trademarks of FMC Corporation

Baythroid® trademark of Mobay Corporation

Comite® trademark of Uniroyal Chemical Company

Curacron® trademark of CIBA-GEIGY Corporation  
U.S. Patent No. 3,992,533

Dipel® trademark of Abbott Laboratory

Guthion® trademark of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen

Lannate® trademark of E. I. DuPont DeNemours & Co.

Larvin® trademark of Rhone-Poulenc & Company

Pay-Off® trademark of American Cyanamid

Pydrin® trademark of Shell Chemical Company


© 1991 CIBA-GEIGY Corporation

Agricultural Division  
CIBA-GEIGY Corporation  
Greensboro, North Carolina 27419

CGA 55L6 (DRAFT)

[GANNONC.LABELC]CURACRON6E\_DRAFT - 08/5/91

Product ID: 2.5 gal. 5A320



**RESTRICTED USE PESTICIDE**

FOR RETAIL SALE TO AND USE ONLY BY  
 CERTIFIED APPLICATORS OR PERSONS  
 UNDER THEIR DIRECT SUPERVISION, AND ONLY  
 FOR THOSE USES COVERED BY THE  
 CERTIFIED APPLICATORS CERTIFICATION

Curacron® 6E

Insecticide - Miticide

For insect and mite control in cotton

Active Ingredient:

Profenofos: O-(4-bromo-2-chlorophenyl) O-ethyl S-	
propyl phosphorothioate .....	59.7%
Inert Ingredients <sup>1</sup> :	40.3%
Total:	100.0%

<sup>1</sup>contains petroleum distillates

2 1/2 Gallons  
U. S Standard Measure

EPA Reg. No. 100-599  
EPA Est. No. 100-AL-1

Curacron 6E contains 6 lbs. active ingredient per gallon.

See directions for use inside booklet.

KEEP OUT OF REACH OF CHILDREN

WARNING

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING

May be fatal if absorbed through skin. Harmful if swallowed or inhaled. Causes moderate eye irritation. Do not get in eyes, on skin, or on clothing. Avoid breathing vapors or



spray mist. Wear protective clothing and rubber gloves when handling. Wash thoroughly with soap and water before eating or smoking. Remove contaminated clothing and wash before reuse.

Statement of Practical Treatment

If in eyes: immediately flush with plenty of water. Get medical attention.

If on skin: wash with plenty of soap and water. Remove contaminated clothing. Get medical attention.

If swallowed: call a physician or poison control center. May present aspiration hazard. DO NOT induce vomiting. Drink large quantities of water. Get medical attention immediately.

If inhaled: remove victim to fresh air.

Note to Physician: If symptoms of cholinergic poisoning occur, atropine plus 2-PAM may be helpful. If swallowed, lavage stomach, taking care to avoid aspiration of stomach contents into the lungs.

Environmental Hazards

Due to the hazard to endangered fish, the application of this product is prohibited in Reeves County, Texas and within one mile of the Dexter National Fish Hatchery in New Mexico. This pesticide is toxic to fish and wildlife. Use with care when applying to areas frequented by wildlife or adjacent to any body of water. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply with aircraft within 300 feet upwind of impounded water. Do not apply when weather conditions favor drift from treated areas. Do not apply with aircraft when wind speed is greater than 10 mph. Apply this pesticide only as specified on this label. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply this product or allow it to drift to crops or weeds on which bees are foraging. Additional information may be obtained from your Cooperative Agricultural Extension Service.

Physical or Chemical Hazards

Do not use or store near open flame.

Do not apply this product through any type of irrigation system.

Curacron® trademark of CIBA-GEIGY Corporation for profenofos  
U.S. Patent No. 3,992,533

©1991 CIBA-GEIGY Corporation

Agricultural Division  
CIBA-GEIGY Corporation  
Greensboro, North Carolina 27419

CGA 55L6 (DRAFT)

Product ID: 54320

[GANJONC.LABELC]CURACRON6E\_DRAFT - 08/5/91

