

B A

Princep[®]

Caliber[™] 90

Herbicide

For weed control
in certain crops,
ornamental plant-
ings, on industrial
sites, and
around the farm

Ten Pounds
Net Weight

Active Ingredient:
Simazine: 2-chloro-4,
6-bis (ethylamino)-
s-triazine 90%
Inert Ingredients: 10%
Total: 100%

EPA Est. 100-LA-1
EPA Reg. No. 100-603

Princep Caliber 90 is a
water dispersible
granule

Keep Out of Reach
of Children.

CAUTION:

See additional pre-
cautionary statements
on back of bag.

See directions for use
on panel below.

CIBA-GEIGY

ACCEPTED

JUL 17 1979

Under the Federal Insecticide,
Fungicide and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 100-603

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

Conditions of Sale and Warranty

The Directions for Use of this product are the property of Ciba-Geigy and are based on field use and tests. The directions are given for the use of this product under the best of conditions. However, it is impossible to make a statement as to the results to be obtained with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application of this product which are beyond the control of Ciba-Geigy or the Seller. All such risks shall be assumed by the Buyer.

Ciba-Geigy warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes intended to be used in the Directions for Use subject to the inherent risks referred to above. **Ciba-Geigy makes no other express or implied warranty of fitness or merchantability or any other express or implied warranty. In no case shall Ciba-Geigy or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product.** Ciba-Geigy and the Seller offer this product and the Buyer and user accept it subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of Ciba-Geigy.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

General Information

Apply this herbicide before weeds emerge or after removal of weed growth. It controls a wide variety of annual broad leaf and grass weeds when used at selective rates in agricultural crops and ornamental plantings. When used at higher, nonselective rates in noncrop areas, it also controls many perennial broad leaf and grass weeds.

Where a range of application rates is given, use the low rate on coarser textured soil and soil lower in organic matter; use the high rate on finer textured soil and soil higher in organic matter.

Since this product enters weeds mainly through the roots, moisture is needed to move it into the root zone. Very dry soil conditions and lack of rainfall following application may necessitate shallow cultivation.

This product is noncorrosive to equipment, nonflammable, and has low electrical conductivity.

FAILURE TO FOLLOW PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR WEED CONTROL AND/OR ILLEGAL RESIDUES.

Annual Weeds Controlled

allyssum	Florida cantabush	ramo, fescue
annual bluegrass	Florida pusley	redmaids
annual morningglory	foxtails	Russian thistle
annual ryegrass	goosegrass	shepherdspurse
barnyardgrass (watergrass)	groundsel	sorghum
burclover	hairy wood	sunflower spp.
carelessweed	juglone	silver chaffgrass
carpetweed	knawel (German moss)	smartweed
common chickweed	common lamb's quarters	spartanweed
crabgrass (<i>D. glabra</i> spp.)	nightshade	speedwell
downy brome (cheatgrass)	pepperweed	tanysmum
fall panicum	powerweed	wild mustard
foxtail	pineappleweed	wild oats
fraxinella	prickly lettuce	wiregrass
freeweed	common purslane	witchgrass
greenhook bassia	ragweed	yellow flower
		pepperweed

Equipment

Ground Application: Use conventional spray equipment with fan type nozzles. Screens in nozzles as well as those in suction and in line strainers should be no finer than 50-mesh.

Use hydraulic or mechanical agitation during mixing and application to maintain a uniform suspension.

Use a pump with capacity to maintain 35-40 psi at the nozzles. If hydraulic agitation is used, the pump should also provide sufficient agitation in the tank to keep the mixture in suspension.

Aerial Application: Use aerial application only where specified in the use directions. Screens in nozzles as well as those in suction and in line strainers should be no finer than 50-mesh.

Avoid application under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Avoid application directly to animals or humans. Although it is unnecessary for fragmen or loaders to wear protective clothing or equipment, they should avoid inhalation of dust or spray mist and prolonged contact with skin. Fragmen and loaders should wash thoroughly before eating and at the end of each day's operation.

Mixing Instructions

Mixing procedure: (1) Be sure sprayer is clean and not contaminated with 2,4-D, residual oil, or other materials. (2) Fill tank half to two-thirds full with clean water. (3) Start agitation. (4) Pour product directly from bag into tank. Let it wet and settle into water. (5) Finish filling tank with water. Continue agitation during mixing and application. (6) Wash sprayer thoroughly with clean water immediately after use.

For ground application where the amount of water is not specified, apply this product in a minimum of 20 gals. of spray mixture per acre.

For aerial application, use a minimum of 20 gals. of water per acre plus 1 lb. of product per acre of spray mixture per acre.

Band Treatment

The amount of product to be applied in band treatment is as indicated by the following formula:

$$\text{Rate (lb./acre)} = \frac{\text{Rate (lb./acre)} \times \text{Band width (ft.)}}{\text{Row spacing (ft.)}}$$

Rotational Crops and Perennial Crop Replanting

To avoid crop injury, observe the following precautions:

1. Do not treat in the year following application, plant only, unless other wise stated in this label. 2. If replanting perennial crops or if rotating to 3rd crops other than corn, do not apply this product in the year preceding planting these crops.

Fruit and Nut Crops

Apply the spray to the orchard or vineyard floor avoiding contact with fruit, foliage, or stems. Apply only to orchards or groves where trees have been established one year or more unless specified differently. Make only one application per year, except as noted otherwise. Recommended rates are based on broadcast treatment. For band applications or spot applications around trees in fruit or nut plantings, reduce the broadcast rate of this product and water per acre in proportion to the area actually sprayed.

Almonds and Peaches (CA only)

Apply 1-2 2 lbs. per treated acre in a 2-4 ft. band on each side of the tree row. Apply before weeds emerge in late fall or early winter. Weeds controlled by 1-1 lbs. include burclover, common chickweed, wild mustard, and shepherdspurse. Apply only once per year.

Precautions: 1. Do not treat trees established in the grove less than 3 years. 2. Do not treat the Mission (Texas) variety of almonds. 3. Do not apply to almond trees propagated on plum rootstocks. 4. Do not replant almonds or peaches in treated soil for 12 months after treatment. 5. Do not apply on gravel, sand, or loamy sand soil or on soil with less than 1% organic matter. 6. Do not treat areas where water will accumulate.

Apples, Pears, Sour Cherries

Apply 2-4 4 lbs. per acre

Avocados

CA and FL only. Apply 2-4 4 lbs. per acre after final preparation of grove.

Precaution: Do not apply on gravelly sand or loamy sand soil.

Blueberries and Caneberries (blackberries, boysenberries, loganberries, raspberries)

Apply 2-4 4 lbs. per acre in the spring or apply a split application of 2 2 lbs. per acre in the spring plus 2 2 lbs. per acre in the fall. Apply in a minimum of 40 gals. of water per acre.

On plantings less than 6 months old, use 1/2 the above rate.

To control quackgrass, apply 4 4 lbs. per acre in the fall or split the application applying 2 2 lbs. per acre in the fall plus 2 2 lbs. per acre in the spring, when quackgrass is actively growing.

Note: Do not apply when fruit is present.

Cranberries

MA. Apply up to 4 4 lbs. per acre either before spring growth begins or in the fall after harvest.

Other areas. Apply 2 2 lbs. per acre before spring growth begins.

Filberts

Apply 2-4 4 lbs. per acre in the fall or apply a split application of 2 2 lbs. per acre in the fall plus 2 2 lbs. per acre in the spring.

Precautions: 1. If trees are planted on a hillside, excessive soil erosion may result from the elimination of weeds. 2. Do not apply on gravelly sand or loamy sand soil.

Note: Do not apply when nuts are on the ground during the harvest period.

Grapes

Apply 2-5 3 lbs. per acre any time between harvest and early spring.

Precaution: Do not use in vineyards established less than three years. Do not apply on gravelly sand or loamy sand soil.

Grapefruit, Lemons, Oranges

AZ (Lemons and Oranges only). Apply a split application of 1 75 lbs. per acre in the spring plus 1 75 lbs. per acre in the fall.

CA. In grapefruit, apply 2-4 4 lbs. per acre. In lemons and oranges, apply 2-4 4 lbs. per acre in a single application or apply 2 2 lbs. per acre in the fall and 2 2 lbs. per acre in the spring. Do not use in the Imperial, Coachella, or Palo Verde v.

FL (Grapefruit and Oranges only). Apply only once per year. For control of weeds listed under General Information, apply 7-10 6 lbs. per acre in the spring, to also control nutweed vine, use 8-9-10 6 lbs. per acre either preemergence or before vines are 12 inches tall.

TX (Grapefruit and Oranges only). Apply 4-5 3 lbs. per acre.

Precautions (All areas): 1. Do not use in nurseries. 2. Do not apply to bedded grapefruit, lemons, or oranges. 3. To avoid possible injury, do not apply to trees under stress from freeze damage for one year after the freeze.

BEST DOCUMENT AVAILABLE

[illegible]

2. 1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100. 101. 102. 103. 104. 105. 106. 107. 108. 109. 110. 111. 112. 113. 114. 115. 116. 117. 118. 119. 120. 121. 122. 123. 124. 125. 126. 127. 128. 129. 130. 131. 132. 133. 134. 135. 136. 137. 138. 139. 140. 141. 142. 143. 144. 145. 146. 147. 148. 149. 150. 151. 152. 153. 154. 155. 156. 157. 158. 159. 160. 161. 162. 163. 164. 165. 166. 167. 168. 169. 170. 171. 172. 173. 174. 175. 176. 177. 178. 179. 180. 181. 182. 183. 184. 185. 186. 187. 188. 189. 190. 191. 192. 193. 194. 195. 196. 197. 198. 199. 200. 201. 202. 203. 204. 205. 206. 207. 208. 209. 210. 211. 212. 213. 214. 215. 216. 217. 218. 219. 220. 221. 222. 223. 224. 225. 226. 227. 228. 229. 230. 231. 232. 233. 234. 235. 236. 237. 238. 239. 240. 241. 242. 243. 244. 245. 246. 247. 248. 249. 250. 251. 252. 253. 254. 255. 256. 257. 258. 259. 260. 261. 262. 263. 264. 265. 266. 267. 268. 269. 270. 271. 272. 273. 274. 275. 276. 277. 278. 279. 280. 281. 282. 283. 284. 285. 286. 287. 288. 289. 290. 291. 292. 293. 294. 295. 296. 297. 298. 299. 300. 301. 302. 303. 304. 305. 306. 307. 308. 309. 310. 311. 312. 313. 314. 315. 316. 317. 318. 319. 320. 321. 322. 323. 324. 325. 326. 327. 328. 329. 330. 331. 332. 333. 334. 335. 336. 337. 338. 339. 340. 341. 342. 343. 344. 345. 346. 347. 348. 349. 350. 351. 352. 353. 354. 355. 356. 357. 358. 359. 360. 361. 362. 363. 364. 365. 366. 367. 368. 369. 370. 371. 372. 373. 374. 375. 376. 377. 378. 379. 380. 381. 382. 383. 384. 385. 386. 387. 388. 389. 390. 391. 392. 393. 394. 395. 396. 397. 398. 399. 400. 401. 402. 403. 404. 405. 406. 407. 408. 409. 410. 411. 412. 413. 414. 415. 416. 417. 418. 419. 420. 421. 422. 423. 424. 425. 426. 427. 428. 429. 430. 431. 432. 433. 434. 435. 436. 437. 438. 439. 440. 441. 442. 443. 444. 445. 446. 447. 448. 449. 450. 451. 452. 453. 454. 455. 456. 457. 458. 459. 460. 461. 462. 463. 464. 465. 466. 467. 468. 469. 470. 471. 472. 473. 474. 475. 476. 477. 478. 479. 480. 481. 482. 483. 484. 485. 486. 487. 488. 489. 490. 491. 492. 493. 494. 495. 496. 497. 498. 499. 500. 501. 502. 503. 504. 505. 506. 507. 508. 509. 510. 511. 512. 513. 514. 515. 516. 517. 518. 519. 520. 521. 522. 523. 524. 525. 526. 527. 528. 529. 530. 531. 532. 533. 534. 535. 536. 537. 538. 539. 540. 541. 542. 543. 544. 545. 546. 547. 548. 549. 550. 551. 552. 553. 554. 555. 556. 557. 558. 559. 560. 561. 562. 563. 564. 565. 566. 567. 568. 569. 570. 571. 572. 573. 574. 575. 576. 577. 578. 579. 580. 581. 582. 583. 584. 585. 586. 587. 588. 589. 590. 591. 592. 593. 594. 595. 596. 597. 598. 599. 600. 601. 602. 603. 604. 605. 606. 607. 608. 609. 610. 611. 612. 613. 614. 615. 616. 617. 618. 619. 620. 621. 622. 623. 624. 625. 626. 627. 628. 629. 630. 631. 632. 633. 634. 635. 636. 637. 638. 639. 640. 641. 642. 643. 644. 645. 646. 647. 648. 649. 650. 651. 652. 653. 654. 655. 656. 657. 658. 659. 660. 661. 662. 663. 664. 665. 666. 667. 668. 669. 670. 671. 672. 673. 674. 675. 676. 677. 678. 679. 680. 681. 682. 683. 684. 685. 686. 687. 688. 689. 690. 691. 692. 693. 694. 695. 696. 697. 698. 699. 700. 701. 702. 703. 704. 705. 706. 707. 708. 709. 710. 711. 712. 713. 714. 715. 716. 717. 718. 719. 720. 721. 722. 723. 724. 725. 726. 727. 728. 729. 730. 731. 732. 733. 734. 735. 736. 737. 738. 739. 740. 741. 742. 743. 744. 745. 746. 747. 748. 749. 750. 751. 752. 753. 754. 755. 756. 757. 758. 759. 760. 761. 762. 763. 764. 765. 766. 767. 768. 769. 770. 771. 772. 773. 774. 775. 776. 777. 778. 779. 780. 781. 782. 783. 784. 785. 786. 787. 788. 789. 790. 791. 792. 793. 794. 795. 796. 797. 798. 799. 800. 801. 802. 803. 804. 805. 806. 807. 808. 809. 810. 811. 812. 813. 814. 815. 816. 817. 818. 819. 820. 821. 822. 823. 824. 825. 826. 827. 828. 829. 830. 831. 832. 833. 834. 835. 836. 837. 838. 839. 8

As a result of the 1995-1996 season, the 1996-1997 season was the best for the 1996-1997 season.

the fact that the two different types of cells are not found in the same place. A cell of one type is found in the same place as a cell of the other type. A cell of one type is found in the same place as a cell of the other type.

[illegible]

| Soil texture | Rate per acre |
|--|---------------|
| Sand, damp sand | 20-25 lbs |
| Sandy loam | 20 lbs |
| Loam or clay soil
low in organic matter | 22-25 lbs |
| Clay, high in
organic matter | 30-44 lbs |

Precautions: Do not use west of the Rocky River in TX or NM, AZ, or CA as no hawk occur. Do not make approximations to transparent trees that have been established less than two years in the grove as they may not

Note: Do not apply when nuts are on the ground. Do not allow animals to graze treated areas.

For control of chinchweed, quailgrass, mustard and shepherdspurse, apply broadcast 1-1.5 lb per acre in fields where overhead irrigation is used to irrigate. If product is applied after harvest at time of bed preparation in fields where overhead irrigation is not available, apply during late October through November.

Precautions: Make only one application per growing season. Do not apply within 4 months after transplanting. Do not exceed 11.4 g/ha (1 lb/acre) per application. Do not use on sand.

App. 2 2 4 4 lbs per acre

and following operations and application of explosives, etc., in the process of construction.

Note: Do not apply when nuts are on the ground.

Take mixture with Elix-BOW for grapefruit and oranges.

Use in grapefruit and oranges in FL and TX for control of insects listed in the following directions and on the label as follows: Apply 1.5 oz. per tree when at or near 30 days of water stress (fruiting) for grapefruit and orange but avoid contact with citrus fruit. In grapefruit, for control of scale, when only Apply 1.5 oz. per tree has returned after frost in the spring. In orange, for control of scale, exceed 1.5 oz. per tree (except as noted for Elix-BOW). Apply 1.5 oz. per tree per year. To avoid injury, do not use in nurseries or around trees that have been established in the grove less than two years or where trees are under stress from freeze damage the year after the freeze. Do not apply within 10 days before harvest of citrus grapefruit or oranges. Observe all precautions and limitations of the labeling of Elix-BOW.

Fl. Ridge Area: Apply 4.4 lbs. of this product plus 5 lbs. of E.L. 80W per acre controls no-weed vine when applied either pre-emergence or on vines up to 24 inches tall.

Full Bedded Areas: Apply 3¹/₂ lbs. of this product plus 6 lbs. of E-180W per acre controls balsam poplar when applied either preemergence or on vines up to 12 inches tall.

Tx. Apr. 3 5 4 lbs. of this product plus 2 3 lbs. of E. = 80% per acre plus 2 qts. nonionic surfactant such as xxx per 100 gals. of spray mixture add surfactant to spray tank as controls buffelgrass, manzanita, iron tree, Texas panicum, sandbar, and sunflower and gives top 4 of oniongrass use 4 lbs. of this product per acre for bare residual control use 5 lbs. of E. = 80% per acre for control of heavy weed growth and for top kill of oniongrass use the higher rate of both products for control of manzanita

This product tank mixed with Paraquat CL is effective in the fruit and nut crops listed in the following table for all of existing vegetation and for residual control of the annual broadleaf and grass weeds listed under General Information, except where noted in the following use directions. This combination is also effective for top kill and suppression of perennial weeds.

Apply the appropriate rates in 50-100 gals./30-50 gals. for pecans, of water per acre to the orchard floor avoiding contact with fruit, foliage or stems. Add a nonionic surfactant such as X-77 at 1 pt. per 100 gals. of spray. Apply when weeds are succulent and new growth is 1-6 inches tall. For mature wood, weeds or difficult to control perennial weeds, treat or spot treat with Paraquat CL if regrowth occurs.

Add this product to the spray tank first, refer to Mix and Instructions section of this label, then add Paraquat CL and add the surfactant as instructed. Provide constant agitation during mixing and application to keep the mixture in suspension.

NEST DOCUMENT AVAILABLE

[illegible][illegible][illegible]

1. The A. A. C. is a "non-profit" corporation, organized under the laws of the State of New York, and is not a charitable organization as defined by the Internal Revenue Code.

[illegible][illegible]

| Soil texture | Rate per acre |
|------------------------------------|---------------|
| Finest sand, low in organic matter | 40 |
| Hard sand | 45 to 50 |
| Coarse sand, low in organic matter | 45 to 55 |
| Clays | 50 to 55 |

24x5d¹⁰er¹⁰g

For the first three years, the average number of children per female application officer was 1.5, after 3.5 years per male officer, and 2.5 after 3.5 years per female officer.

[illegible]

Preemergence Apply to seed and/or emerge (use the appropriate rate in the table) **POA**

Preplant All seeds were stored at 4°C in the dark until the day of sowing. After sowing, the seeds were kept in the dark at 25°C until germination. The seeds were then stored at 4°C until the day of sowing. The seeds were then stored at 4°C until the day of sowing.

* 1990-1991, 1992-1993, 1994-1995, 1996-1997, 1998-1999, 2000-2001, 2002-2003, 2004-2005, 2006-2007, 2008-2009, 2010-2011, 2012-2013, 2014-2015, 2016-2017, 2018-2019, 2020-2021, 2022-2023, 2024-2025, 2026-2027, 2028-2029, 2030-2031, 2032-2033, 2034-2035, 2036-2037, 2038-2039, 2040-2041, 2042-2043, 2044-2045, 2046-2047, 2048-2049, 2050-2051, 2052-2053, 2054-2055, 2056-2057, 2058-2059, 2060-2061, 2062-2063, 2064-2065, 2066-2067, 2068-2069, 2070-2071, 2072-2073, 2074-2075, 2076-2077, 2078-2079, 2080-2081, 2082-2083, 2084-2085, 2086-2087, 2088-2089, 2090-2091, 2092-2093, 2094-2095, 2096-2097, 2098-2099, 2100-2101, 2102-2103, 2104-2105, 2106-2107, 2108-2109, 2110-2111, 2112-2113, 2114-2115, 2116-2117, 2118-2119, 2120-2121, 2122-2123, 2124-2125, 2126-2127, 2128-2129, 2130-2131, 2132-2133, 2134-2135, 2136-2137, 2138-2139, 2140-2141, 2142-2143, 2144-2145, 2146-2147, 2148-2149, 2150-2151, 2152-2153, 2154-2155, 2156-2157, 2158-2159, 2160-2161, 2162-2163, 2164-2165, 2166-2167, 2168-2169, 2170-2171, 2172-2173, 2174-2175, 2176-2177, 2178-2179, 2180-2181, 2182-2183, 2184-2185, 2186-2187, 2188-2189, 2190-2191, 2192-2193, 2194-2195, 2196-2197, 2198-2199, 2200-2201, 2202-2203, 2204-2205, 2206-2207, 2208-2209, 2210-2211, 2212-2213, 2214-2215, 2216-2217, 2218-2219, 2220-2221, 2222-2223, 2224-2225, 2226-2227, 2228-2229, 2230-2231, 2232-2233, 2234-2235, 2236-2237, 2238-2239, 2240-2241, 2242-2243, 2244-2245, 2246-2247, 2248-2249, 2250-2251, 2252-2253, 2254-2255, 2256-2257, 2258-2259, 2260-2261, 2262-2263, 2264-2265, 2266-2267, 2268-2269, 2270-2271, 2272-2273, 2274-2275, 2276-2277, 2278-2279, 2280-2281, 2282-2283, 2284-2285, 2286-2287, 2288-2289, 2290-2291, 2292-2293, 2294-2295, 2296-2297, 2298-2299, 2300-2301, 2302-2303, 2304-2305, 2306-2307, 2308-2309, 2310-2311, 2312-2313, 2314-2315, 2316-2317, 2318-2319, 2320-2321, 2322-2323, 2324-2325, 2326-2327, 2328-2329, 2330-2331, 2332-2333, 2334-2335, 2336-2337, 2338-2339, 2340-2341, 2342-2343, 2344-2345, 2346-2347, 2348-2349, 2350-2351, 2352-2353, 2354-2355, 2356-2357, 2358-2359, 2360-2361, 2362-2363, 2364-2365, 2366-2367, 2368-2369, 2370-2371, 2372-2373, 2374-2375, 2376-2377, 2378-2379, 2380-2381, 2382-2383, 2384-2385, 2386-2387, 2388-2389, 2390-2391, 2392-2393, 2394-2395, 2396-2397, 2398-2399, 2400-2401, 2402-2403, 2404-2405, 2406-2407, 2408-2409, 2410-2411, 2412-2413, 2414-2415, 2416-2417, 2418-2419, 2420-2421, 2422-2423, 2424-2425, 2426-2427, 2428-2429, 2430-2431, 2432-2433, 2434-2435, 2436-2437, 2438-2439, 2440-2441, 2442-2443, 2444-2445, 2446-2447, 2448-2449, 2450-2451, 2452-2453, 2454-2455, 2456-2457, 2458-2459, 2460-2461, 2462-2463, 2464-2465, 2466-2467, 2468-2469, 2470-2471, 2472-2473, 2474-2475, 2476-2477, 2478-2479, 2480-2481, 2482-2483, 2484-2485, 2486-2487, 2488-2489, 2490-2491, 2492-2493, 2494-2495, 2496-2497, 2498-2499, 2500-2501, 2502-2503, 2504-2505, 2506-2507, 2508-2509, 2510-2511, 2512-2513, 2514-2515, 2516-2517, 2518-2519, 2520-2521, 2522-2523, 2524-2525, 2526-2527, 2528-2529, 2530-2531, 2532-2533, 2534-2535, 2536-2537, 2538-2539, 2540-2541, 2542-2543, 2544-2545, 2546-2547, 2548-2549, 2550-2551, 2552-2553, 2554-2555, 2556-2557, 2558-2559, 2560-2561, 2562-2563, 2564-2565, 2566-2567, 2568-2569, 2570-2571, 2572-2573, 2574-2575, 2576-2577, 2578-2579, 2580-2581, 2582-2583, 2584-2585, 2586-2587, 2588-2589, 2590-2591, 2592-2593, 2594-2595, 2596-2597, 2598-2599, 2600-2601, 2602-2603, 2604-2605, 2606-2607, 2608-2609, 2610-2611, 2612-2613, 2614-2615, 2616-2617, 2618-2619, 2620-2621, 2622-2623, 2624-2625, 2626-2627, 2628-2629, 2630-2631, 2632-2633, 2634-2635, 2636-2637, 2638-2639, 2640-2641, 2642-2643, 2644-2645, 2646-2647, 2648-2649, 2650-2651, 2652-2653, 2654-2655, 2656-2657, 2658-2659, 2660-2661, 2662-2663, 2664-2665, 2666-2667, 2668-2669, 2670-2671, 2672-2673, 2674-2675, 2676-2677, 2678-2679, 2680-2681, 2682-2683, 2684-2685, 2686-2687, 2688-2689, 2690-2691, 2692-2693, 2694-2695, 2696-2697, 2698-2699, 2700-2701, 2702-2703, 2704-2705, 2706-2707, 2708-2709, 2710-2711, 2712-2713, 2714-2715, 2716-2717, 2718-2719, 2720-2721, 2722-2723, 2724-2725, 2726-2727, 2728-2729, 2730-2731, 2732-2733, 2

| Soil texture | Broadcast rate per acre |
|---|-------------------------|
| Coarse textured (sand, sand loam, loam, silty loam, silty clay loam, silty clay, clay loam, clay) | 4-5 ts |
| Medium textured (silt, silt loam, silty clay, silty clay loam, silty clay, clay) | 4-5 ts |
| Fine textured (silt, silty clay, silty clay loam, silty clay, clay) | 4-5 ts |
| Organics (peat, muck, and other organic soils) | 4-5 ts |

*For calculation of land hearing rate, see the Land Hearing Information section.

[illegible]

Note: It needs to be pointed out that, under the above conditions, a standard cultivation will generally result in better weeds control.

[illegible]

[illegible]

Paragard CL is a registered trademark of Paragard, Inc. All other trademarks are the property of their respective owners. © 2007 Paragard, Inc. All rights reserved.

ALL INFORMATION CONTAINED HEREIN IS UNCLASSIFIED DATE 01-11-2001 BY 60322 UCBAW/STP

$\Delta \rho_{\text{eff}} = \rho_{\text{eff}} - \rho_{\text{eff}}^0$ is the difference between the effective density of the polymer solution and the effective density of the pure polymer, ρ_{eff}^0 , at the same temperature and concentration. ρ_{eff}^0 is calculated from the densities of the pure components and the composition of the polymer solution. $\Delta \rho_{\text{eff}}$ is a function of the concentration of the polymer solution and the temperature. The temperature dependence of $\Delta \rho_{\text{eff}}$ is shown in Figure 1. The temperature dependence of $\Delta \rho_{\text{eff}}$ is very weak in the range of 20–40 °C. The temperature dependence of $\Delta \rho_{\text{eff}}$ is very weak in the range of 20–40 °C.

the design of the system, the general idea of the system, and the design of the system.

Figure 4.4-15. Population of the dominant species of the 10 most common weeds in
South America, 1950-1970. The data are based on the number of individuals per 10
square meters of the following weeds:

For a long time, I have been interested in the question of the effect of the environment on the development of the individual. I have been particularly interested in the question of the effect of the environment on the development of the individual's personality. I have been particularly interested in the question of the effect of the environment on the development of the individual's personality. I have been particularly interested in the question of the effect of the environment on the development of the individual's personality.

Grasses Grown for Seed (Pacific Northwest Only)

Grasses Grown for Seed (Pacific Northwest Only)

Perennial ryegrass, highland or Astoria perennegrasses, orchardgrass, tall fescue and fine fescues such as tree fescue, Pennsylvanian Chewings and related species for fine swards of broadleaf weeds and annual grasses, including annual ryegrass, Italian ryegrass, slender ryegrass and lawn from wheat, apply 2.2 lbs in a minimum of 1 gal. of water per acre as soon as a rain starts. Early application is important.

From J. C. Anderson - 10 grasses from which at least one species has been cul-

4. The A cow 30 days between last application and grazing of dairy, and beef cattle and sheep, and a cow 60 days between the last application and milking for dairy.

Audience: Christians, non-Christians, Jews, Muslims, Buddhists, Hindus, etc.

Nurseries (see list below)
Age: 2, 3, 4 yrs. at least 25 gal. of water per acre in a spring at least one year after transplanting

Christmas Tree Plantings and Shelterbelts (see list below)
 We have seed grain before application. Apply 2.2-4.4 lbs. of urea or 20 lbs. of water per acre after
 application. Use the same rate for annual maintenance applications.

transplanting, use the same rate for annual maintenance applications.
 For broadcast applications, apply 4.4 lb per acre in the fall or apply a split application of 2.2 lbs per acre in the
 fall and 2.2 lbs per acre in the spring after lucerngrass begins growth.

13. Seedlings: Do not use or seedbeds or cutting beds. In CA, OR, and WA do not apply to stressed trees or seedlings younger than one year after transplanting in riparian areas. In riparian areas stressed trees or shrubs smaller than spurs less than three years old. In riparian areas more than one year except as directed for riparian areas.

As a consequence, the following results are obtained:

Comments

| Conifers | Deciduous |
|-----------------|-----------------|
| Arbutus tree | Bayberry shrub |
| Australian pine | Black spruce |
| Bishop pine | White spruce |
| Jack pine | Salisbury white |
| Red pine | Virginia pine |
| Scots pine | Fraser fir |
| Montezuma pine | White fir |
| Mugho pine | Hemlock |
| Red pine | Red cedar |
| Swedish pine | White cedar |
| Scotch pine | Yellow cedar |
| White pine | Yellow cedar |
| Blue spruce | Yellow cedar |

Page 151

[illegible]

- 400 C.A.U.

Free information on: www.elsevier.com/locate/locate/locate

Apply in the spring before weeds emerge and before new growth appears. If applying after transplanting, do not apply to the top of tree seedlings before new growth appears. If new growth is present, do not spray the sprouts and foliage contact. Do not use on seedling sprouts of 1/4" or more. Apply once per year.

The following rates are based on the standard treatment of 2,000 gals. of water per acre. For bands or spot applications, reduce the broadcast rate proportionally to the water actually sprayed.

App. 2 2 4 4 5

App. 2 2 4 4 cs per acre

156255-5183

Use of certain symbols on the Form, App. 44-6 is required.

use for es'ale

4.0 General Information: See Broad Spectrum for general information. See also General Information section for applicable weed control and maintenance areas and its production.

SECRET

BEST DOCUMENT AVAILABLE