



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

November 10, 2021

Cherilyn Moore
Regulatory Product Manager
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, NC 27419

Subject: Registration Review Label Amendments for Simazine Incorporating Mitigation Measures from the Interim Decision and the Technical Registrants' Commitments for the Endangered Species Act (ESA) Biological Evaluation
Product Name: PRINCEP CALIBER 90 HERBICIDE
EPA Registration Number: 100-603
Application Date: 11/20/2020
Decision Number: 568105

Dear Ms. Moore:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Simazine Interim Decision and with the technical registrants' commitments for the ESA Biological Evaluation. The Agency has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.


A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only

Page 2 of 2
EPA Reg. No. 100-603
Decision No. 568105

distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Anna Senninger at Senninger.Anna@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Julie Javier", is written over a light yellow rectangular background.

Julie Javier, Team Leader
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

Enclosure

[Master Label]

Sale, use, and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.

Not for use in the states of Hawaii or Alaska or in the U.S. territories (Puerto Rico, Guam, American Samoa, the U.S. Virgin Islands, and the North Mariana Islands).

SIMAZINE	GROUP	5	HERBICIDE
-----------------	--------------	----------	------------------

Princep® Caliber 90® Herbicide

For weed control in certain crops

Active Ingredient:

Simazine: 2-chloro-4,6-bis(ethylamino)-s-triazine 90.0%

Other Ingredients: 10.0%

Total: 100.0%

Princep Caliber 90 is a water-dispersible granule.

KEEP OUT OF REACH OF CHILDREN.


CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-603

EPA Est.

Net Contents



FIRST AID	
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

Personal Protective Equipment (PPE)

Mixers and loaders supporting groundboom applications must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥14 mils, or Viton® ≥14 mils
- Chemical-resistant footwear plus socks
- Chemical-resistant apron
- A minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R, or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters

All other mixers, loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥14 mils, or Viton ≥14 mils

Applications made by mechanically pressurized handguns to strawberries are restricted to either spot treatment only or mixer/loader/applicators must wear coveralls over long-sleeved shirt and long pants in addition to the PPE specified above.

Mixer/loader/applicators for backpack application to grapefruit and oranges must wear coveralls over long-sleeved shirt and long pants in addition to the PPE specified above.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Engineering Control Statements

When handlers use enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-5)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

Simazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Simazine has been found in ground water. Users are advised not to apply simazine to sand and loamy sand soils where the water table (ground water) is close to the surface and where these soils are very permeable; i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.

Product must not be mixed or loaded within 50 feet of intermittent streams and rivers, natural or impounded lakes and reservoirs. Product must not be applied within 66 feet of points where field surface water runoff enters perennial or intermittent streams and rivers or within 200 feet of natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66 foot buffer or setback from runoff entry points must be planted to crop, or seeded with grass or other suitable crop.

Product must not be mixed or loaded, or used within 50 feet of all wells, including abandoned wells, drainage wells, and sink holes. Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited, unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rain water that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment of 100% of the capacity of the largest pesticide container or application

equipment on the pad. Containment capacities as described above shall be maintained at all times. The above-specified minimum containment capacities do not apply to vehicles when delivering pesticide to the mixing/loading sites.

Additional State imposed requirements regarding well-head setbacks and operational area containment must be observed.

One of the following restrictions must be used in applying simazine to tile-outletted terraced fields containing standpipes:

- Do not apply within 66 feet of standpipes in tile-outletted terraced fields.
- Apply this product to the entire tile-outletted terraced field and immediately incorporate it to a depth of 2-3 inches in the entire field.
- Apply this product to the entire tile-outletted terraced field under a no-till practice only when a high crop residue management practice is practiced. High crop residue management is described as a crop management practice where little or no crop residue is removed from the field during and after crop harvest.

NON-TARGET ORGANISM ADVISORY STATEMENT

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

ANY USE OF THIS PRODUCT IN AN AREA WHERE USE IS PROHIBITED IS A VIOLATION OF FEDERAL LAW. Before using this product, you must consult the Simazine Watershed Information Center (SWIC) to determine whether the use of this product is prohibited in your watershed. SWIC can be accessed through www.simazine-watershed.info or 1-888-365-2874. If the SWIC indicates that use of this product is prohibited in your watershed, you may return this product to your point of purchase or contact Syngenta Crop Protection, LLC for a refund.

Endangered Species

It is a Federal offense to use any pesticide in a manner that results in an unauthorized “take” (e.g., kill or otherwise harm) of an endangered species under the Endangered Species Act section 9. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the area in which you are applying the product. You must obtain a Bulletin no earlier than six months before using this product. To obtain Bulletins, consult <http://www.epa.gov/espp/>, call 1-844-447-3813, or email ESPP@epa.gov. You must use the Bulletin valid for the month in which you will apply the product.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

For Christmas trees, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours. For all other crops and use patterns, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves made of any waterproof material
- Shoes plus socks

Applications made by mechanically pressurized handguns are restricted to spot treatment only for the following uses

- Citrus (Grapefruit, Oranges, Lemons)
- Pome Fruits (Apples, Pears)
- Stone Fruits (Cherries [sweet and tart], peaches, Plums, Nectarines)
- Tree Nuts (Pecans, Walnuts, Filberts, Almonds, Macadamia Nuts)
- Berry and Small Fruit (Blueberries, Blackberries, Loganberries, Raspberries, Grapes, Lowbush Blueberries, Cranberries)
- Tropical and Sub-tropical Fruits (Avocado, Olive)
- Nursery/Ornamentals
- Sweet Corn

Applications made by mechanically pressurized handguns to strawberries are restricted to either spot treatment only or mixer/loader/applicators must wear coveralls over long-sleeved shirt and long pants, shoes plus socks, and gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others (including children or pets) to enter treated areas until sprays have dried.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR WEED CONTROL, AND/OR ILLEGAL RESIDUES.

USE RESTRICTIONS

Sale, use, and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.

Not for use in the states of Hawaii or Alaska or in the U.S. territories (Puerto Rico, Guam, American Samoa, the U.S. Virgin Islands, and the North Mariana Islands).

Do not apply this product through any type of irrigation system.

For use on warm season turf grass **only** (e.g., Bermudagrass, Centipedegrass, St. Augustinegrass, and Zoysiagrass).

All shelterbelt and forestry uses, except "Christmas Tree Plantings", are prohibited.

This product is not for use in formulating end-use products for aquatic use.

PRODUCT INFORMATION

Aerial application is prohibited.

When tank mixing or sequentially applying simazine or products containing simazine, the total pounds of simazine applied must not exceed the specific maximum rate per calendar year as noted in the use directions.

Apply this herbicide before weeds emerge or after removal of weed growth. Princep Caliber 90 controls a wide variety of annual broadleaf and grass weeds when used at selective rates in agricultural crops.

Where a range of application rates is given, use the low rate on coarse-textured soil and soil lower in organic matter; use the high rate on fine-textured soil and soil higher in organic matter.

Since this product enters weeds mainly through their roots, moisture is needed to move it into the root zone. Very dry soil conditions and lack of rainfall following application may necessitate shallow cultivation.

This product is noncorrosive to equipment, nonflammable, and has low electrical conductivity.

ANNUAL WEEDS CONTROLLED

alyssum	common lambsquarters
annual bluegrass	nightshade
annual morningglory	pepperweed
annual ryegrass	pigweed
barnyardgrass (watergrass)	pineappleweed
burclover	prickly lettuce
carelessweed	common purslane
carpetweed	ragweed
common chickweed	rattail fescue
crabgrass (<i>Digitaria</i> spp.)	redmaids
downy brome (cheatgrass)	Russian thistle
fall panicum	shepherdspurse
fiddleneck	signalgrass (<i>Brachiaria</i> spp.)
filaree	silver hairgrass
fireweed	smartweed
fivehook bassia	spanishneedles
Flora's paintbrush	speedwell
Florida pusley	tansymustard
foxtails	wild mustard
goosegrass	wild oats
groundsel	witchgrass
henbit	yellow flower pepperweed
jungerice	
knawel (German moss)	

WEED RESISTANCE MANAGEMENT

SIMAZINE	GROUP 5	HERBICIDE
-----------------	----------------	------------------

Simazine, the active ingredient in this product, is a Group 5 herbicide based on the mode of action classification system of the Weed Science Society of America. Any weed population may contain plants naturally resistant to Group 5 herbicides. Such

resistant weed plants may not be effectively managed using Group 5 herbicides but may be effectively managed utilizing another herbicide alone or in mixtures from a different group and/or by using cultural or mechanical practices. However, an herbicide mode of action classification by itself may not adequately address specific weeds that are resistant to specific herbicides. Consult your local company representative, state cooperative extension service, professional consultants or other qualified authorities to determine appropriate actions for treating specific resistant weeds.

Scout and know your field

- Know weed species present in the field to be treated through scouting and field history. An understanding of weed biology is useful in designing a resistance management strategy. Ensure the weed management program will control all weeds present.
- Fields should be scouted prior to application to determine species present and growth stage. Always apply this herbicide at the full labeled rate and correct timing for the weeds present in the field.

Utilize non-herbicidal practices to add diversity

- Use diversified management tactics such as cover crops, mechanical weed control, harvest weed seed control, and crop rotation as appropriate.

Use good agronomic practices, start clean and stay clean

- Use good agronomic practices that enhance crop competitiveness.
- Plant into weed-free fields utilizing tillage or an effective burndown herbicide for control of emerged weeds.
- Sanitize farm equipment to avoid spreading seed or vegetative propagules prior to leaving fields.

Difficult to control weeds

- Fields with difficult to control weeds should be planted in rotation with crops that allow the use of herbicides with an alternative mode of action or different management practices.
- Difficult to control weeds may require sequential applications, such as a broad spectrum preemergence herbicide followed by one or more postemergence herbicide applications. Utilize herbicides containing different modes of action effective on the target weeds in sequential applications.

Do not overuse the technology

- Do not use more than two applications of this or any other herbicide with the same mode of action in a single growing season unless mixed with an herbicide with a different mode of action which provides overlapping spectrum for the difficult to control weeds.

Scout and inspect fields following application

- Prevent an influx of weeds into the field by controlling weeds in field borders.
- Scout fields after application to verify that the treatment was effective.
- Suspected- herbicide resistant weeds may be identified by these indicators
 - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - A spreading patch of non-controlled plants of a particular weed species; and
 - Surviving plants mixed with controlled individuals of the same species.
- Report non-performance of this product to your Syngenta retailer, Syngenta representative, or call 1-866-Syngent(a) (866-796-4368). If resistance is suspected ensure weed escapes are controlled using an herbicide with an effective mode of action and/or use non-chemical means to prevent further seed production.

Prevent weed escapes before, during, and after harvest

- Do not allow weed escapes to produce seed or vegetative structures such as tubers or stolons which contribute to spread and survival. Consider harvest weed seed management and control weeds post-harvest to prevent seed production.

Resistant Weeds

- Contact your local Syngenta representative, retailer, crop advisor or extension agent to determine if weeds resistant to this mode of action are present in your area. If resistant biotypes have been reported, use the full labeled rate of this product, apply at the labeled timing, and tank-mix with a different mode of action product so there are multiple effective modes of application for each suspected resistant weed.

PERENNIAL CROP REPLANTING

If replanting perennial crops, do not apply Princep Caliber 90 in the year preceding planting these crops, unless otherwise stated on this label, or crop injury may occur.

APPLICATION PROCEDURES

Ground application: Use conventional ground sprayers equipped with nozzles that provide accurate and uniform application. Be certain that nozzles are uniformly spaced and the same size. Calibrate sprayer before use and recalibrate at the start of each season and when changing carriers. Unless otherwise specified, use a minimum of 20 gallons of spray mixture per acre.

Use a pump with capacity to: (1) maintain 35-40 psi at nozzles, (2) provide sufficient agitation in tank to keep mixture in suspension, and (3) to provide a minimum of 20% bypass at all times. Use centrifugal pumps which provide propeller shear action for dispersing and mixing this product. The pump should provide a minimum of 10 gallons/minute/100 gallons tank size circulated through a correctly positioned sparger tube or jets.

Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on suction side of pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles. Check nozzle manufacturer's recommendations.

For band applications, calculate amount to be applied per acre as follows:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

MANDATORY SPRAY DRIFT MANAGEMENT

Ground Boom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a coarse or coarser droplet size (ASABE S572)
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.
- User must maintain a 15 foot (4.6 meter) in-field downwind buffer (in the direction in which the wind is blowing) from the edge of streams and rivers, as well as high-tide line for all estuarine/marine environments.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If

a greater spray volume is needed, consider using a nozzle with a higher flow rate.

- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce spray drift.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

WIND

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.** Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Handheld Technology Applications

Take precautions to minimize spray drift.

MIXING PROCEDURES

All Uses: (1) Be sure sprayer is clean and not contaminated with any other materials, as crop injury or sprayer clogging may result. (2) Fill tank $\frac{1}{4}$ full with clean water, nitrogen solution, or complete fluid fertilizer. (3) Start agitation. (4) Be certain that the

agitation system is working properly and creates a rippling or rolling action on the liquid surface. (5) Pour product directly from bag into tank. (6) Continue filling tank until 90% full. Increase agitation if necessary to maintain surface action. (7) Add tank mix herbicide(s) after this product is thoroughly suspended. (8) Finish filling tank. Maintain agitation to avoid separation of materials. (9) Empty tank as completely as possible before refilling to prevent buildup of emulsifiable concentrate residue from possible tank mix herbicides. (10) If an emulsifiable concentrate film starts to build up in tank, drain it and clean with strong detergent solution or solvent. (11) Clean sprayer thoroughly immediately after use by flushing system with water containing a detergent.

Compatibility test: To determine the tank mix compatibility of Princep Caliber 90 with liquid fertilizer, crop oil, spreaders, or recommended pesticides, use this test method. Nitrogen solutions or complete fluid fertilizers may replace all or part of the water in the spray in corn. Since liquid fertilizers can vary, even within the same analysis, check compatibility each time before use. Be especially careful when using **complete** suspension or fluid fertilizers, as serious compatibility problems are more apt to occur. Commercial application equipment may improve compatibility in some instances. Check compatibility using this procedure:

1. Add 1 pt. of fertilizer to each of two 1-qt. glass jars with tight lids.
2. To **one** of the jars, add ¼ teaspoon of a compatibility agent approved for this use (¼ teaspoon is equivalent to 2 pt./100 gallons spray). Cap and shake until mixed. Examples of compatibility agents include Compex® and Unite®.
3. To **both** jars, add the appropriate amount of herbicide(s). If more than one herbicide is used, add them separately with dry herbicides first, flowables next, and emulsifiable concentrates last. After each addition, cap and shake until thoroughly mixed. The appropriate amount of herbicides for this test follows (assuming a spray volume of 25 gallons/A):

Dry herbicides: **For each pound per acre**, add 1.5 teaspoons to each jar. Fluff up wettable powder products before measuring.

Liquid herbicides: **For each pint per acre**, add 0.5 teaspoon to each jar.

For a spray volume other than 25 gallons/A, change the teaspoons added to each jar as follows:

$$\frac{25 \text{ gallons} \times \text{number teaspoons given above}}{\text{desired gallons spray volume per acre}} = \text{teaspoons to add to each jar}$$

4. After adding all ingredients, put lids on and tighten. Shake jars vigorously one minute. Let the mixtures stand 15 minutes and then look for separation, large flakes, precipitates, grease, gels, medium to heavy oily film on the jar, or other

signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the 2 jars. If either mixture separates, but can be remixed readily, the mixture probably can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility: (A) slurry the dry herbicide(s) in water before addition, or (B) add ½ of the compatibility agent to the fertilizer and the other ½ to the emulsifiable concentrate or flowable herbicide before addition to the mixture. If still incompatible, do not use the materials mixed in the same spray tank.

CROP USE DIRECTIONS

FRUIT AND NUT CROPS

Apply the spray to the orchard or vineyard floor, avoiding contact with fruit, foliage, or stems. Specified rates are based on broadcast treatment. For band applications or spot applications around fruit or nut trees, reduce the broadcast rate of this product and water per acre in proportion to the area actually sprayed.

Precautions (All fruit and nut crops): To avoid crop injury, (1) Apply only to orchards or groves where trees have been established 12 months or more, unless specified differently. (2) Make only one application per year, except as noted otherwise. (3) Do not use on gravelly, sand, or loamy sand soil. (4) Immediately following application, limit overhead sprinkler irrigation to ½ inch.

Almonds, Peaches, and Nectarines (CA Only)

Princep Caliber 90 must be applied in a 2-4 foot band on each side of the tree row at a rate of 1.1-2.2 lb. Princep Caliber 90 per treated acre. Apply before weeds emerge in late fall or early winter. Weeds controlled by 1.1 lb. include burclover, common chickweed, wild mustard, and shepherdspurse. Do not apply more than 2.2 lb. Princep Caliber 90 (2.2 lb. a.i./treated acre) per calendar year. Do not apply more than once per calendar year.

Precautions: To avoid crop injury, (1) Do not treat trees established in the grove less than 3 years. (2) Do not treat the Mission (TX) variety of almonds. (3) Do not apply to almond trees propagated on plum rootstocks. (4) Do not replant almonds, peaches, or nectarines in treated soil for 12 months after treatment. (5) Do not apply to soil with less than 1% organic matter. (6) Do not treat areas where water will accumulate.

Apples, Pears, and Tart Cherries

Apply 2.2-4.4 lb./A Princep Caliber 90. Do not apply more than 4.4 lb./A Princep Caliber 90 (4 lb. a.i./A) per calendar year. Do not apply more than once per calendar year. For apples, the pre-harvest interval is 150 days.

Avocados

CA and FL Only: Apply 2.2-4.4 lb./A Princep Caliber 90 after final preparation of grove. Do not apply more than 4.4 lb. Princep Caliber 90 (4 lb. a.i./A) per calendar year. Do not apply more than once per calendar year.

Blueberries and Caneberries (Blackberries, Boysenberries, Loganberries, Raspberries)

Apply 2.2-4.4 lb./A Princep Caliber 90 in the spring or apply a split application of 2.2 lb./A Princep Caliber 90 in the spring plus 2.2 lb./A Princep Caliber 90 in the fall. Apply in a minimum of 40 gallons of water per acre.

On plantings less than 6 months old, use ½ the above rate.

To control quackgrass, apply 4.4 lb./A Princep Caliber 90 in the fall or split the application applying 2.2 lb./A Princep Caliber 90 in the fall plus 2.2 lb./A Princep Caliber 90 in the spring, when quackgrass is actively growing.

Do not apply more than 4.4 lb./A Princep Caliber 90 (4 lb. a.i./A) per calendar year. Do not apply more than twice per calendar year.

Note: Do not apply when fruit is present, or illegal residues may result.

Filberts

Apply 2.2-4.4 lb./A Princep Caliber 90 in the fall or apply a split application of 2.2 lb./A Princep Caliber 90 in the fall plus 2.2 lb./A Princep Caliber 90 in the spring. Do not apply more than 4.4 lb./A Princep Caliber 90 (4 lb. a.i./A) per calendar year. Do not apply more than twice per calendar year.

Precaution: If trees are planted on a hillside, excessive soil erosion may result from the elimination of weeds.

Note: Do not apply when nuts are on the ground during the harvest period, or illegal residues may result.

Grapes

Apply 2.2-4.4 lb./A Princep Caliber 90 any time between harvest and early spring. Do not apply more than 4.4 lb./A Princep Caliber 90 (4 lb. a.i./A) per calendar year. Do not apply more than once per calendar year.

Precaution: Do not use in vineyards established less than 3 years, or crop injury may occur.

Grapefruit, Lemons, and Oranges

AZ (Lemons and Oranges Only): Apply a split application of 1.77 lb./A Princep Caliber 90 in the spring plus 1.77 lb./A Princep Caliber 90 in the fall. Do not apply more than 3.54 lb./A Princep Caliber 90 (3.2 lb. a.i./A) per calendar year. Do not apply more than 1.77 lb./A Princep Caliber 90 (1.6 lb. a.i./A) per application.

CA (Grapefruit, Lemons, and Oranges): Apply 2.2-4.4 lb./A Princep Caliber 90 in a single application, or apply 2.2 lb./A Princep Caliber 90 in the fall and 2.2 lb./A Princep Caliber 90 in the spring. Do not apply more than 4.4 lb./A Princep Caliber 90 (4 lb. a.i./A) per calendar year. Do not apply more than twice per calendar year.

Precaution: Do not use in the Imperial, Coachella, or Palo Verde valleys, or crop injury may occur.

FL (Grapefruit and Oranges Only)

Broadcast Applications: Apply up to 4.4 lb. of Princep Caliber 90/A per application to weed-free soil once during the spring and/or once during the fall to control weeds expected to emerge during these periods. Apply prior to emergence of weeds or if weeds have emerged, apply in tank mixture with a contact herbicide. Use caution to keep the treatment off the foliage, fruit, or trunk of citrus trees. Do not apply more than 4.4 lb. Princep Caliber 90 per acre (4 lb. a.i./A) per application. Do not apply more than 8.8 lb. Princep Caliber 90 per acre (8 lb. a.i./A) per calendar year.

Banded Applications: For control of difficult species, such as balsamapple and spanishneedles, and partial control of honeyvine milkweed, apply no more than 8.8 lb. of Princep Caliber 90 (8 lb. a.i./A) of Princep Caliber 90 as a single application in the spring using a 50% band application. The amount of Princep Caliber 90 applied per acre of field must not exceed 4.4 lb. (4 lb. a.i./A). Apply once in the spring growing season between January and April. Do not make a fall Princep Caliber 90 application if this treatment was used in the spring. When emerged weeds are present, apply Princep Caliber 90 in tank mixture with a recommended contact herbicide. Follow all directions, precautions, limitations, etc. on the tank mix product.

TX (Grapefruit and Oranges Only): Apply 2.2 - 4.4 lb./A Princep Caliber 90. Do not apply more than 4.4 lb. of Princep Caliber 90 per acre (4 lb. a.i./A) per calendar year. Do not apply more than twice per calendar year.

Precautions (All areas): To avoid crop injury, (1) Do not use in nurseries. (2) Do not apply to bedded grapefruit, lemons, or oranges (except for FL grapefruit and oranges). (3) Do not apply to trees under stress from freeze damage for one year after the freeze. (4) Do not exceed 4.4 lb./acre/year of Princep Caliber 90, or crop injury may occur.
Note: Dark-red grapefruit hybrids, e.g. "Star Ruby", have a higher risk of crop injury than non-dark-red types.

Macadamia Nuts

Apply 2.2-4.4 lb. Princep Caliber 90 in 50 gallons of water per acre before harvest and just prior to weed emergence. Do not apply more than 4.4 lb./A Princep Caliber 90 (4 lb. a.i./A) per calendar year. Do not apply more than once per calendar year.

Note: Do not apply when nuts are on the ground during the harvest period, as illegal residues may result.

Olives

Apply 2.2-4.4 lb./A Princep Caliber 90 following grove preparation in the fall. Repeat annually in midwinter. Do not apply more than 4.4 lb./A Princep Caliber 90 (4 lb. a.i./A) per calendar year. Do not apply more than once per calendar year.

Peaches, Plums, and Sweet Cherries

Apply 1.77-4.4 lb./A Princep Caliber 90. Apply in late fall to early spring prior to weed emergence. Do not apply more than 4.4 lb./A Princep Caliber 90 (4 lb. a.i./A) per calendar year. Do not apply more than once per calendar year.

Precautions: To avoid crop injury, (1) Peaches: use only in AR, LA, MO, OK, TX, and states east of the Mississippi River. For CA, see specific directions in the Almonds, Peaches, and Nectarines (CA only) section. (2) Plums and sweet cherries: use only in MO and states east of the Mississippi River, except TN.

Pecans

Apply 2.2-4.4 lb./A Princep Caliber 90 before weeds emerge in the spring. Do not apply more than 4.4 lb./A Princep Caliber 90 (4 lb. a.i./A) per calendar year. Do not apply more than once per calendar year.

Precautions: To avoid crop injury, (1) Do not use west of the Pecos River in TX or in AZ, CA, or NM. (2) Do not make applications to transplanted trees that have been established less than 2 years in the grove.

Note: Do not apply when nuts are on the ground, or illegal residues may result. Do not allow animals to graze treated areas.

Strawberries

OR and WA Only: For control of chickweed, groundsel, mustard, and shepherdspurse, apply broadcast 1.1 lb./A Princep Caliber 90. In fields where overhead irrigation is used to activate this product, apply after harvest at time of bed renovation. In fields where overhead irrigation is not available, apply during early October through November. Do

not apply more than 1.1 lb./A Princep Caliber 90 (1 lb. a.i./A) per calendar year. Do not apply more than once per calendar year.

Precautions: To avoid crop injury, (1) Make only one application per growing season. (2) Do not apply within 4 months after transplanting.

Walnuts

Apply 2.2-4.4 lb./A Princep Caliber 90. Leveling and furrowing operations after application will lessen effectiveness of weed control. Do not apply more than 4.4 lb./A Princep Caliber 90 (4 lb. a.i./A) per calendar year. Do not apply more than once per calendar year.

Note: Do not apply when nuts are on the ground, or illegal residues may result.

Tank Mixture with Bromacil 80W for Grapefruit and Oranges (FL Only)

Use in grapefruit and oranges in FL for control of balsamapple, black nightshade, carpetweed, crabgrass, cudweed, dayweed, Florida pusley, horseweed, pepperweed, pigweed, poorjoe, ragweed, rattlebox, spanishneedles and sandbur, and for partial control of bermudagrass, bahiagrass, pangolagrass, paragrass, and torpedograss. Apply 4.4 lb. of Princep Caliber 90 plus 3-4 lb. of Bromacil 80W per acre beneath trees in a minimum of 40 gallons of water per acre before or soon after weed growth begins. Use caution to keep the spray off the foliage, fruit, or trunk of citrus trees. Temporary yellowing of citrus leaves may occur following treatment.

Follow use and precautionary directions on the Bromacil label.

Precautions: To avoid crop injury, (1) Do not use in nurseries or where trees are under stress from freeze damage for one year after the freeze. (2) Do not use on soil with less than 1% organic matter or on poorly drained soil. (3) Do not treat trees planted in irrigation furrows. (4) Do not treat diseased trees, such as those with foot rot. (5) Do not use in groves interplanted with other trees or desirable plants, nor in home grapefruit or orange plantings, or in areas where roots of other valuable plants or trees may extend. (6) Treated areas may be planted to citrus trees one year after application. (7) Do not rotate to other crops within 2 years after application.

Note: Apply only once per year and avoid contact with foliage and fruit with spray or mist, or illegal residues may result.

Tank Mixture with Gramoxone Inteon®

This tank mix is effective in the following fruit and nut crops for kill of existing vegetation and for residual control of the annual broadleaf and grass weeds claimed for this product applied alone. This combination is also effective for top kill and suppression of perennial weeds. In FL, this mixture may be applied in spring or fall to emerged weeds.

Use this tank mix on these crops:

almonds (CA only)	olives
apples	oranges ²
avocados (CA and FL ¹ only)	peaches ³
cherries (tart and sweet ⁴)	pears
filberts	pecans
grapefruit (CA and TX only)	plums ⁴
grapes	walnuts
lemons (AZ and CA only)	
macadamia nuts	

Apply the rate given under the appropriate crop on this label plus the appropriate labeled rate of Gramoxone Inteon in 50-200 gallons (30-50 gallons for pecans) of water per acre to the orchard floor, avoiding contact with fruit, foliage, or stems. Add a nonionic surfactant, such as X-77®, at 0.5 pt./100 gallons of spray. Apply when weeds are succulent and new growth is 1-6 inches tall. For mature woody weeds or difficult to control perennial weeds, re-treat or spot treat with Gramoxone Inteon if regrowth occurs.

Add this product to the spray tank first (refer to the **Mixing procedures** section of this label), then add Gramoxone Inteon, and add the surfactant last. Provide constant agitation during mixing and application to keep the mixture in suspension.

Precautions: To avoid crop injury, (1) Apply the tank mix only once per year. (2) Use a shield for young trees or vines.

Refer to the labels of both products for further directions, specific weeds controlled, and precautions and limitations on each crop.

¹In avocados in FL, this tank mix also controls balsamapple vine, rattail amaranth, and at the higher specified rate of each herbicide, it suppresses coral vine.

²In oranges in FL, apply 4.4 lb. of this product per acre per application during spring or fall. Do not exceed 8.8 lb. of Princep Caliber 90 per calendar year.

³Limited to AR, CA, LA, MO, OK, TX, and states east of the Mississippi River. As specified, refer to the sections **Almonds, Peaches, and Nectarines (CA only) or Peaches, Plums, and Sweet Cherries** for rate of this product and other information.

⁴Limited to MO and states east of the Mississippi River, except TN.

Tank Mixture with Roundup® or Touchdown® Brand Herbicides

This tank mixture is effective in grape vineyards and in the following bearing or nonbearing tree crops for control of existing vegetation and for residual control of the annual broadleaf and grass weeds claimed for this product applied alone. This combination is also effective for partial control of perennial weeds contacted by the spray mixture during application.

Use this tank mixture on these crops:

almonds** (CA only)	macadamia nuts**
apples	oranges*** (AZ, CA, FL, and TX)
avocados* (CA and FL only)	peaches**** ¹
cherries (tart and sweet**** ²)	pears*
filberts**	pecans**
grapefruit*** (CA, FL, and TX)	plums**** ²
grapes*	walnuts**
lemons* (AZ and CA)	

Use the specified rate given elsewhere on this label for Princep Caliber 90 applied alone to the crop being treated. Add to the spray tank the specified labeled rate of a Roundup or Touchdown brand product per acre, depending on weeds present and their growth stage. Also add an agriculturally approved nonionic surfactant at 0.5% by volume of spray solution. Apply the mixture in 10-40 gallons of water per acre as a postemergence spray to the weeds at the appropriate weed growth stage given on the Roundup label.

Add Princep Caliber 90 to the spray tank first, then add the Roundup or Touchdown brand product. Provide constant agitation during mixing and application to keep the mixture in suspension. Refer to the **Application Procedures** section of this label for further directions.

Precaution: To avoid crop injury, take extreme care to avoid contact of herbicide solution, spray, drift, or mist with foliage or green bark of trunk, branches, suckers, fruit, or other parts of trees or vines. Observe precautions on both the Princep Caliber 90 and Roundup or Touchdown labels for each crop involved.

Always refer to the labels of both herbicides for further directions, specific weeds controlled, precautions, and limitations on each crop.

*Allow a minimum of 14 days between last application and harvest.

**Allow a minimum of 21 days between last application and harvest of these crops.

***Allow a minimum of 1 day between last application and harvest of these crops.

****Allow a minimum of 17 days between last application and harvest of these crops;

¹Limited to AR, CA, LA, MO, OK, TX, and states east of the Mississippi River. For CA, see specific directions in the **Almonds, Peaches, and Nectarines (CA only)** section.

²Plums and sweet cherries: limited to MO and states east of the Mississippi River, except TN.

Tank Mixture with Solicam®

For improved control of such weeds as clover, cutleaf eveningprimrose, dandelion, henbit, horseweed or marestalk, lambsquarters, and puncturevine, apply Princep Caliber 90 in tank mixture with Solicam DF on these crops:

oranges
grapefruit
lemons

Apply 4.4 lb./A* of Princep Caliber 90 plus 4-5 lb. of Solicam DF Herbicide (78.6% active ingredient) in 20-100 gallons of water per acre. Princep Caliber 90 + Solicam may be applied in tank mixture with Gramoxone Inteon or Roundup.

*Maximum 1.77 lbs/A in Arizona.

Follow the labeled directions and restrictions of use on labels of the herbicides used in mixtures with Princep Caliber 90.

Precaution: Keep Princep Caliber 90 + Solicam mixtures from contacting foliage, fruits, and stems of citrus trees during spraying.

Tank Mixture with Surflan®

Use this tank mix in the following crops for preemergence control of all weeds claimed on both labels:

almonds	lemons
apples	oranges
avocados	peaches
caneberries	pears
cherries	pecans
filberts	plums
grapefruit	English walnuts
grapes	

Apply the Princep Caliber 90 rate given under the appropriate crop on this label plus 2.67-5.33 lb. of Surflan 75W or 2-4 qt. of Surflan A.S. in 20-40 gallons of water per acre. Refer to the Surflan 75W or Surflan A.S. label for complete tank mix directions. Observe all precautions and limitations on the Princep Caliber 90 and Surflan labels.

FIELD AND FORAGE CROPS

Corn (Field and Sweet)

Apply a maximum of 2.2 lb. Princep Caliber 90 per acre (2.0 lb. a.i./A) as a single preemergence application on soils that are not highly erodible or on highly erodible soils, as defined by the Natural Resources Conservation Service, if at least 30% of the soil is covered with plant residues. If a second treatment is required following an earlier herbicide application, the total simazine applied may not exceed 2.5 lb. a.i./A per calendar year.

Apply a maximum of 1.77 lb. Princep Caliber 90 per acre (1.6 lb. a.i./A) as a single preemergence application on highly erodible soils if <30% of the soil is covered with plant residues.

Pre-grazing/pre-harvest interval for field corn: 60 days.

Pre-grazing/pre-harvest interval for sweet corn: 45 days.

Nitrogen solutions or complete fluid fertilizers may replace all or part of the water as a carrier in the spray. Determine the physical compatibility of this product with fertilizer before use. Refer to the **Product Information** section of this label for a compatibility test. Do not apply after corn has emerged, as there is danger of liquid fertilizers causing crop injury.

Preemergence: Apply before weeds and corn emerge. Use the specified rate as stated above.

Preplant: Apply in the spring after plowing at the specified rate as stated above. Apply before, during, or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation. Best results will be obtained when this product is applied within 2 weeks before planting. Under dry weather conditions, preplant applications may give better weed control. If weeds develop, particularly under relatively dry conditions, a shallow cultivation will generally result in better weed control.

Winter Annual Broadleaf Control - Preemergence Fall Application

For preemergence control of winter annual weeds, such as common chickweed, henbit, shepherdspurse, tansymustard, wild mustard, annual bluegrass, downy brome, and others, broadcast 1.1 lb./A of Princep Caliber 90 after harvest of the preceding crop and prior to weed emergence on land to be planted to corn the following year. A tillage operation may precede the application. Do not apply to frozen ground. If Princep Caliber 90 is used in the fall corn weed control program, do not exceed 1.67 lb. of Princep Caliber 90 preemergence in the spring.

Precautions: (1) Do not apply more than 2.77 lb./A Princep Caliber 90 to corn per year. (2) Do not rotate to any crop except corn until the following year, or injury may occur. (3) After harvest, plow and thoroughly till the soil in fall or spring to minimize possible injury to spring-seeded rotational crops, regardless of the rate used. (4) Do not apply preplant incorporated in corn in the High Plains and Intermountain areas of the West (including central and western KS, western NE, western OK, and the Panhandle of TX) where rainfall is sparse and erratic or where irrigation is required. (5) In the High Plains and Intermountain areas of the West where rainfall is sparse and erratic or where irrigation is required, use this product to control weeds in corn only when corn is to follow corn or when a crop of untreated corn is to precede another rotational crop. (6) Injury may occur to soybeans planted in north-central and northwest IA, south-central and southwest MN, northeast NE, southeast SD, and other areas the year following application on soils having a calcareous surface layer. (7) Do not plant sugar beets, tobacco, vegetables (including dry beans), spring-seeded small grains, or small-seeded legumes and grasses the year after an application, or injury may occur.

Note: Do not graze treated areas, or illegal residues may result.

Tank Mixtures on Corn

AAtrex®: Use this product in a preplant or preemergence tank mixture with AAtrex Nine-O®, or AAtrex 4L for control of many annual weeds, including carpetweed, crabgrass, fall panicum, foxtail, lambsquarters, morningglory, pigweed, ragweed, and velvetleaf.

When tank mixing or sequentially applying simazine and/or products containing either active ingredient to corn, the total pounds of simazine and/or atrazine applied must not exceed 2.5 lbs a.i./acre per calendar year.

Refer to the AAtrex label for complete directions, and use 1.1 lb. of this product for each qt. of Princep 4L. Also refer to the AAtrex label and the **Corn** section of this label for precautions and limitations.

Gramoxone Inteon: Use in a tank mixture where corn will be planted directly in a cover crop, established sod, or previous crop residues. This combination controls existing vegetation and provides residual control of the annual broadleaf and grass weeds listed under the **Product Information** section.

Add this product to the spray tank, mix thoroughly with water, and then add Gramoxone Inteon and a nonionic surfactant. Provide constant agitation during mixing and application to keep the mixture in suspension.

Apply a maximum of 1.77 lb. Princep Caliber 90/A (1.6 lbs. a.i./A) as a single preemergence application on highly erodible soils if <30% of the soil is covered with plant residues.

Apply up to 2.2 lb. of this product plus the specified labeled rate of Gramoxone Inteon in 20-60 gallons of water per acre as a broadcast spray either before or after planting, but before corn emerges. Add a nonionic surfactant at the rate of 0.5 pt./100 gallons of spray volume.

For further information, see product information, caution and warning statements, precautions, and notes on the Princep Caliber 90 and the Gramoxone Inteon labels.

CHRISTMAS TREE PLANTINGS

Remove weed growth before application. Apply 2.2-4.4 lb. Princep Caliber 90 in at least 25 gallons of water per acre after transplanting. Use the same rate for annual maintenance applications.

For quackgrass control, apply 4.4 lb./A Princep Caliber 90 in the fall or apply a split application of 2.2 lb./A Princep Caliber 90 in the fall plus 2.2 lb./A Princep Caliber 90 in early spring, after quackgrass begins growth.

Do not apply more than 4.4 lb./A Princep Caliber 90 (4 lb. a.i./A) per calendar year. Do not apply more than twice per calendar year.

Precautions: To avoid tree injury, (1) Do not use on seedbeds or cutting beds. (2) In CA, OR, and WA, do not apply to Christmas trees sooner than one year after transplanting. In other areas, do not apply to Christmas trees transplants less than 2 years of age. (3) Do not use until soil is firmly settled around roots. (4) Do not apply more than once a year, except as directed for quackgrass control.

Apply to these species of trees and shrubs, as specified above:

Conifers

arborvitae
Austrian pine
Bishop pine*
knobcone pine*
lodgepole pine (shore pine)
Monterey pine*
Mugho pine
red pine (Norway pine)
Scotch pine
white pine
blue spruce
Norway spruce
red spruce
white spruce
balsam fir
Douglas fir
Fraser fir
white fir
hemlock
juniper
red cedar
white cedar
yew (*Taxus* spp.)

*For CA only

Surflan Tank Mix: On Christmas tree plantings, use this tank mix for preemergence control of weeds listed on this label and the Surflan 75W (or Surflan A.S.) label. Use on field grown conifer species listed on the labels for each herbicide plus grand fir, alpine fir, Engelmann spruce, black spruce, Colorado blue spruce, Coulter pine, giant redwood, and Veitchi fir. Broadcast the mixture as a directed spray to the soil surface or as an ovetop spray, using 2.2-4.4 lb. of Princep Caliber 90 and 2.67-5.33 lb. of Surflan 75W (or 2-4 qt. of Surflan A.S.). Apply in sufficient water per acre to uniformly treat the area. Follow ovetop sprays with sprinkler irrigation to move the herbicide from leaf surfaces to the soil. Remove weed growth before application. Mix weed residues, prunings, or trash into the soil, or remove them before treatment. Soil should be in good tilth and free of clods at time of application. Shallow cultivation (1-2 inches) after treatment will not reduce weed control. Observe all precautions and limitations on the Princep Caliber 90 and Surflan labels.

Note: Length of weed control may be reduced when continuous wet soil conditions follow herbicide application.

Precautions: To avoid plant injury, (1) Do not use on seedbeds or on unrooted cuttings. (2) Do not use in greenhouses or other enclosed areas.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in a dry place.

Pesticide Disposal

Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling

Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

AAtrex®, AAtrex® Nine-O®, Gramoxone Inteon®, Princep® Caliber 90®, Solicam®, Touchdown®, the ALLIANCE FRAME, the SYNGENTA Logo, and the PURPOSE ICON are Trademarks of a Syngenta Group Company

Compex® trademark of KALO Agricultural Chemicals, Inc.

Roundup® trademark of Monsanto Company

Surflan® trademark of Dow AgroSciences

Unite® trademark of HACO, Inc.

X-77® trademark of Loveland Industries, Inc.

Viton® is a trademark of E. I. duPont de Nemours and Co.

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

©20XX Syngenta

Manufactured for:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300

Princep Caliber 90 603 MAS 1112 AMEND-D NOV2020-CL - di 11/8/2021
000100-00603.20201120D.PRINCEPCALIBER90_AMEND_1120-CL